



planetmath.org

Math for the people, by the people.

finite changes in convergent series

Canonical name	FiniteChangesInConvergentSeries
Date of creation	2013-03-22 19:03:10
Last modified on	2013-03-22 19:03:10
Owner	pahio (2872)
Last modified by	pahio (2872)
Numerical id	9
Author	pahio (2872)
Entry type	Theorem
Classification	msc 40A05
Related topic	SumOfSeriesDependsOnOrder
Related topic	RiemannSeriesTheorem
Related topic	RatioTestOfDAlembert

The following theorem means that at the beginning of a convergent series, one can remove or attach a finite amount of terms without influencing on the convergence of the series – the convergence is determined alone by the infinitely long “tail” of the series. Consequently, one can also freely change the of a finite amount of terms.

Theorem. Let k be a natural number. A series $\sum_{n=1}^{\infty} a_n$ converges iff the series $\sum_{n=k+1}^{\infty} a_n$ converges. Then the sums of both series are with

$$\sum_{n=k+1}^{\infty} a_n = \sum_{n=1}^{\infty} a_n - \sum_{n=1}^k a_n. \quad (1)$$

Proof. Denote the k th partial sum of $\sum_{n=1}^{\infty} a_n$ by S_k and the n th partial sum of $\sum_{n=k+1}^{\infty} a_n$ by S'_n . Then we have

$$S'_n = \sum_{n=k+1}^{k+n} a_n = S_{k+n} - S_k. \quad (2)$$

1°. If $\sum_{n=1}^{\infty} a_n$ converges, i.e. $\lim_{n \rightarrow \infty} S_n := S$ exists as a finite number, then (2) implies

$$\lim_{n \rightarrow \infty} S'_n = \lim_{n \rightarrow \infty} S_{k+n} - \lim_{n \rightarrow \infty} S_k = S - S_k.$$

Thus $\sum_{n=k+1}^{\infty} a_n$ converges and (1) is true.

2°. If we suppose $\sum_{n=k+1}^{\infty} a_n$ to be convergent, i.e. $\lim_{n \rightarrow \infty} S'_n := S'$ exists as finite, then (2) implies that

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} S_{k+n} = \lim_{n \rightarrow \infty} (S_k + S'_n) = S_k + S'.$$

This means that $\sum_{n=1}^{\infty} a_n$ is convergent and $S = S_k + S'$, which is (1), is in .