



AN IDENTITY FOR THE SUM OF INVERSES OF ODD DIVISORS OF n IN TERMS OF THE NUMBER OF REPRESENTATIONS OF n AS A SUM OF SQUARES

SUMIT KUMAR JHA

Let $c_r(n)$ be the number of representations of a positive integer n as a sum of r squares, where representations with different orders and different signs are counted as distinct. We prove a combinatorial identity relating this quantity to the sum of inverses of odd divisors of n :

$$\sum_{\text{odd } d \mid n} \frac{1}{d} = \frac{1}{2} \sum_{r=1}^n \frac{(-1)^{n+r}}{r} \binom{n}{r} c_r(n).$$

1. Main result

Let n be a positive integer. We derive here the following identity for the sum of inverses of odd divisors of n , in terms of the the number $c_r(n)$ of representations of n as a sum of r squares (r ranging over the positive integers, and representations with different orders or different signs being counted as distinct).

$$(1) \quad \sum_{\text{odd } d \mid n} \frac{1}{d} = \frac{1}{2} \sum_{r=1}^n \frac{(-1)^{n+r}}{r} \binom{n}{r} c_r(n).$$

Recall that the $c_r(n)$ occur in the expansion of the r -th power of the function

$$\theta(q) := \prod_{j=1}^{\infty} \frac{1-q^j}{1+q^j} = \sum_{i=-\infty}^{\infty} (-1)^i q^{i^2} \quad (|q| < 1)$$

(see [2, 7.324], for example). More precisely, we have

$$\theta(q)^r = \sum_{n=0}^{\infty} c_r(n) (-1)^n q^n.$$

Let $B_{n,k} \equiv B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ be the partial Bell polynomials [1, p. 134], defined by

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n, \ell_i \in \mathbb{N} \\ \sum_{i=1}^n i \ell_i = n \\ \sum_{i=1}^n \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!} \right)^{\ell_i}.$$

2020 AMS *Mathematics subject classification*: 11A05, 11P99.

Keywords and phrases: sum of divisors, sum of squares, partial Bell polynomials, Faà di Bruno's formula, Jacobi's theta function.

Received by the editors on July 5, 2020, and in revised form on October 23, 2020.

Lemma 1. For all positive integers n , we have

$$(2) \quad 2 \sum_{\text{odd } d \mid n} = \frac{1}{n!} \sum_{k=1}^n (-1)^k (k-1)! B_{n,k}(\theta'(0), \theta''(0), \dots, \theta^{(n-k+1)}(0)),$$

Proof. It is easy to see that

$$\begin{aligned} \log \theta(q) &= \sum_{j=1}^{\infty} \log(1 - q^j) - \sum_{j=1}^{\infty} \log(1 + q^j) = - \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \frac{q^{lj}}{l} + \sum_{j'=1}^{\infty} \sum_{l'=1}^{\infty} \frac{q^{l'j'}(-1)^{l'}}{l'} \\ &= - \sum_{n=1}^{\infty} q^n \left(\sum_{d \mid n} \frac{1 - (-1)^d}{d} \right). \end{aligned}$$

Let $f(q) = \log q$. Using Faà di Bruno's formula (see [1, p. 137] or [3]), we have

$$(3) \quad \frac{d^n}{dq^n} f(\theta(q)) = \sum_{k=1}^n f^{(k)}(\theta(q)) \cdot B_{n,k}(\theta'(q), \theta''(q), \dots, \theta^{(n-k+1)}(q)).$$

Since $f^{(k)}(q) = \frac{(-1)^{k-1} (k-1)!}{q^k}$ and $\theta(0) = 1$, letting $q \rightarrow 0$ in (3) gives the lemma. \square

Lemma 2. We have, for positive integers n, k ,

$$(4) \quad B_{n,k}(\theta'(0), \theta''(0), \dots, \theta^{(n-k+1)}(0)) = (-1)^n \frac{n!}{k!} \sum_{r=1}^k (-1)^{k-r} \binom{k}{r} c_r(n).$$

Proof. We start with the generating function for the partial Bell polynomials [1, Equation (3a') on p. 133], and proceed as follows to conclude (4):

$$\begin{aligned} \sum_{n=k}^{\infty} B_{n,k}(\theta'(0), \theta''(0), \dots, \theta^{(n-k+1)}(0)) \frac{q^n}{n!} &= \frac{1}{k!} \left(\sum_{j=1}^{\infty} \theta^{(j)}(0) \frac{q^j}{j!} \right)^k = \frac{1}{k!} (\theta(q) - 1)^k \\ &= \frac{1}{k!} \sum_{r=0}^k (-1)^{k-r} \binom{k}{r} \theta(q)^r \\ &= \frac{1}{k!} \sum_{r=0}^k (-1)^{k-r} \binom{k}{r} \sum_{n=0}^{\infty} (-1)^n c_r(n) q^n \end{aligned} \quad \square$$

Proof of Equation (1). We combine (2) and (4) to obtain

$$\begin{aligned} 2 \sum_{\text{odd } d \mid n} \frac{1}{d} &= (-1)^n \sum_{k=1}^n \frac{1}{k} \sum_{r=1}^k (-1)^r \binom{k}{r} c_r(n) = (-1)^n \sum_{r=1}^n (-1)^r c_r(n) \sum_{k=r}^n \frac{1}{k} \binom{k}{r} \\ &= (-1)^n \sum_{r=1}^n \frac{(-1)^r}{r} \binom{n}{r} c_r(n), \end{aligned}$$

having used that $\sum_{k=r}^n \frac{1}{k} \binom{k}{r} = \frac{1}{r} \binom{n}{r}$, a consequence of Pascal's formula $\binom{k}{r-1} = \binom{k+1}{r} - \binom{k}{r}$. \square

References

- [1] L. Comtet, *Advanced combinatorics: the art of finite and infinite expansions*, D. Reidel, Dordrecht, 1974.
- [2] N. J. Fine, *Basic hypergeometric series and applications*, Mathematical Surveys and Monographs **27**, American Mathematical Society, Providence, RI, 1988.
- [3] W. P. Johnson, “The curious history of Faà di Bruno’s formula”, *Amer. Math. Monthly* **109**:3 (2002), 217–234.

SUMIT KUMAR JHA: kumarjha.sumit@research.iiit.ac.in

International Institute of Information Technology, Hyderabad, India