

$$\alpha_s = \frac{g^2}{(4\pi)^2} \quad (0.1)$$

$$v^{\iota\alpha\beta\gamma} = (2\eta^{\iota\beta}\eta^{\alpha\gamma} - \eta^{\iota\alpha}\eta^{\beta\gamma} - 2\eta^{\iota\gamma}\eta^{\alpha\beta}) \quad (0.2)$$

d dim integrals from P&S

$$\int \frac{d^d l}{(2\pi)^d} \frac{1}{(l^2 - \Delta)^n} = \frac{(-1)^n i}{(4\pi)^{d/2}} \frac{\Gamma(n - \frac{d}{2})}{\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n - \frac{d}{2}} \quad (0.3)$$

$$\int \frac{d^d l}{(2\pi)^d} \frac{l^2}{(l^2 - \Delta)^n} = \frac{(-1)^{n-1} i}{(4\pi)^{d/2}} \frac{d}{2} \frac{\Gamma(n - \frac{d}{2} - 1)}{\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n - \frac{d}{2} - 1} \quad (0.4)$$

$$\int \frac{d^d l}{(2\pi)^d} \frac{l^\mu l^\nu}{(l^2 - \Delta)^n} = \frac{(-1)^{n-1} i}{(4\pi)^{d/2}} \frac{\eta^{\mu\nu}}{2} \frac{\Gamma(n - \frac{d}{2} - 1)}{\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n - \frac{d}{2} - 1} \quad (0.5)$$

$$\int \frac{d^d l}{(2\pi)^d} \frac{l^\mu l^\nu l^\rho l^\omega}{(l^2 - \Delta)^n} = \frac{(-1)^n i}{(4\pi)^{d/2}} \frac{E^{\mu\nu\rho\omega}}{4} \frac{\Gamma(n - \frac{d}{2} - 2)}{\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n - \frac{d}{2} - 2} \quad (0.6)$$

$$\int \frac{d^d k}{(2\pi)^d} \frac{k^{2a}}{(k^2 - \Delta)^b} = (-1)^{a-b} i \frac{1}{(4\pi)^{\frac{d}{2}}} \frac{\Gamma(a + \frac{d}{2}) \Gamma(b - a - \frac{d}{2})}{\Gamma(b) \Gamma(\frac{d}{2})} \left(\frac{1}{\Delta}\right)^{b - a - \frac{d}{2}} \quad (0.7)$$

last formula (B.35) in [https://www.uni-muenster.de/imperia/md/content/physik\\_tp/lectures/ss2017/standard\\_model/Vorlesung/app\\_b.pdf](https://www.uni-muenster.de/imperia/md/content/physik_tp/lectures/ss2017/standard_model/Vorlesung/app_b.pdf) for two variable integrals(derived on my own)

$$\int d^d x \int d^d y \frac{1}{(-ax^2 - by^2 + m^2)^n} = -a^{-\frac{d}{2}} b^{-\frac{d}{2}} \pi^d \frac{\Gamma(n - d)}{\Gamma(n)} (m^2)^{d-n} \quad (0.8)$$

$$\int d^d x \int d^d y \frac{x^\mu x^\nu}{(-ax^2 - by^2 + m^2)^n} = a^{-\frac{d}{2}-1} b^{-\frac{d}{2}} \pi^d \frac{\eta^{\mu\nu}}{2} \frac{\Gamma(n - d - 1)}{\Gamma(n)} (m^2)^{d-n+1} \quad (0.9)$$

$$\int d^d x \int d^d y \frac{x^\mu x^\nu y^\alpha y^\beta}{(-ax^2 - by^2 + m^2)^n} = -a^{-\frac{d}{2}-1} b^{-\frac{d}{2}-1} \pi^d \frac{\eta^{\mu\nu} \eta^{\alpha\beta}}{4} \frac{\Gamma(n - d - 2)}{\Gamma(n)} (m^2)^{d-n+2} \quad (0.10)$$

$$\int d^d x \int d^d y \frac{x^\mu x^\nu x^\theta x^\phi}{(-ax^2 - by^2 + m^2)^n} = -a^{-\frac{d}{2}-2} b^{-\frac{d}{2}} \pi^d \frac{E^{\mu\nu\theta\phi}}{4} \frac{\Gamma(n - d - 2)}{\Gamma(n)} (m^2)^{d-n+2} \quad (0.11)$$

$$\int d^d x \int d^d y \frac{x^\mu x^\nu y^\alpha y^\beta y^\gamma y^\delta}{(-ax^2 - by^2 + m^2)^n} = a^{-\frac{d}{2}-1} b^{-\frac{d}{2}-2} \pi^d \frac{E^{\alpha\beta\gamma\delta} \eta^{\mu\nu}}{8} \frac{\Gamma(n - d - 3)}{\Gamma(n)} (m^2)^{d-n+3} \quad (0.12)$$

$$\int d^d x \int d^d y \frac{y^\alpha y^\beta y^\gamma y^\delta y^\mu y^\nu}{(-ax^2 - by^2 + m^2)^n} = a^{-\frac{d}{2}-3} b^{-\frac{d}{2}} \pi^d \frac{E_6^{\alpha\beta\gamma\delta\mu\nu}}{16} \frac{\Gamma(n - d - 3)}{\Gamma(n)} (m^2)^{d-n+3} \quad (0.13)$$

Crucial formulae:

$$\frac{1}{A_1^{m_1} \dots A_n^{m_n}} = \int_0^1 dx_1 \dots dx_n \delta\left(\sum x_i - 1\right) \frac{\prod x_i^{m_i-1}}{(\sum x_i A_i)^{\sum m_i}} \frac{\Gamma(\sum m_i)}{\prod \Gamma(m_i)} \quad (0.14)$$

$$\overline{\psi(x)\psi(0)} = G(x) = \frac{i\not{x}}{2\pi^2 x^4} = \frac{i\Gamma(\frac{d}{2})\not{x}}{2\pi^2 (x^2)^{\frac{d}{2}}} \quad (0.15)$$

$$= \frac{\Gamma(\frac{D+1}{2})}{2\pi^{\frac{D+1}{2}}} \frac{i\not{x}}{(-x^2 + i\epsilon)^{\frac{D+1}{2}}} \quad (0.16)$$

$$= \frac{\Gamma(\frac{d}{2})}{2\pi^{\frac{d}{2}}} \frac{i\not{x}}{(-x^2 + i\epsilon)^{\frac{d}{2}}} \quad (0.17)$$

$$CHECK : \overline{A^{\mu,a}(x)} A^{\nu,b}(0) = \frac{-i}{4\pi^2} \frac{\eta^{\mu\nu}}{x^2} (\text{must be wrong}) \quad (0.18)$$

$$= \frac{-i}{4\pi^2} \frac{\Gamma(\frac{d}{2} - 1) \eta^{\mu\nu} \delta^{ab}}{(x^2)^{\frac{d}{2}-1}} \quad (0.19)$$

$$= \frac{-1}{4\pi^{\frac{d}{2}}} \frac{\Gamma(\frac{d}{2} - 1) \eta^{\mu\nu} \delta^{ab}}{(-x^2 + i\epsilon)^{\frac{d}{2}-1}} \quad (0.20)$$

$$\not{A} = A^\mu \gamma_\mu = A^{\mu,a} \gamma_\mu t^a \quad (0.21)$$

$$t^a t^a = C_2 \mathbb{1} = C_F \mathbb{1} \quad (0.22)$$

$$\not{x}\not{x} = x^2 \quad (0.23)$$

$$\not{a}\not{b}\not{a} = -\not{b}a^2 + 2(a \cdot b)\not{a} \quad (0.24)$$

$$\gamma_\mu \gamma^\nu \gamma^\mu = -(d-2)\gamma^\nu \quad (0.25)$$

$$\gamma^\mu \gamma_\mu = d \quad (0.26)$$

work in gauges (q: quantum, c: classical fields)

$$x_\mu A^{\mu,c} = 0 \quad (0.27)$$

$$D_\mu^c A^{\mu,q} = 0 \quad (0.28)$$

$$G^{(1)}(x) = \int dz \frac{i(\not{x} - \not{z})}{2\pi^2(x-z)^4} i g A(z) \frac{i\not{z}}{2\pi^2 z^4} \quad (0.29)$$

$$= \frac{-3ig}{2\pi^4} \int dz \int_0^1 du u \bar{u} \frac{(\bar{u}\not{x} - \not{z}) \gamma_\mu (u\not{x} + \not{z})}{(z^2 + u\bar{u}x^2)^4} A^\mu(z + ux) \quad (0.30)$$

To get there use formula eq. (0.14) and shift  $z \rightarrow z + ux$ . Next we should expand the field  $A$  assuming small  $z$ :  $A^\mu(x+z) = A^\mu(x) + z^\alpha \partial_\alpha A^\mu(x) + z^\beta z^\alpha \partial_\alpha \partial_\beta A^\mu(x)$  Then i get

$$G^{(1)}(x) = \frac{-3ig}{2\pi^4} \int dz \int_0^1 du u \bar{u} \frac{(\bar{u}\not{x} - \not{z}) \gamma_\mu (u\not{x} + \not{z})}{(z^2 + u\bar{u}x^2)^4} (A^\mu(ux) + z^\alpha \partial_\alpha A^\mu(ux) + z^\beta z^\alpha \partial_\alpha \partial_\beta A^\mu(ux)) \quad (0.31)$$

I am actually not sure whether the expansion should be only applied to field  $A$ , or also the left (fraction) term. I could also shift derivatives by integration by parts. I start with the easiest term:  $z = 0$ : look at the

numerator:

$$(\bar{u}\not{x} - \not{x})\gamma_\mu(u\not{x} + z) = \bar{u}u\not{x}\gamma_\mu\not{x} + \bar{u}\not{x}\gamma_\mu\not{x} + u\not{x}\gamma_\mu\not{x} + \not{x}\gamma_\mu\not{x} \quad (0.32)$$

$$= 2\bar{u}u x_\mu - \bar{u}u\gamma_\mu x^2 + \bar{u}\not{x}\gamma_\mu\not{x} + u\not{x}\gamma_\mu\not{x} + 2z_\mu - \gamma_\mu z^2 \quad (0.33)$$

for  $A=\text{const}$  i have only the contributions terms 1, 2 and 6 (from left to right) due to symmetry in  $z$ .

$$T_0 = \frac{-3ig}{2\pi^4} \int_0^1 du u \bar{u} \left( \frac{i\pi^2}{6\bar{u}u} \frac{2x_\mu\not{x} - \gamma_\mu x^2}{x^4} + \frac{-2i\pi^2 - \gamma_\mu}{6u\bar{u}} \frac{1}{x^2} \right) A^\mu(ux) \quad (0.34)$$

$$= \frac{-g}{4\pi^2} \int_0^1 du \left( \frac{2x_\mu\not{x}}{x^4} + \frac{\gamma_\mu}{x^2} \right) A^\mu(ux) \quad (0.35)$$

This expression i can also find in Vladimirs paper for the first  $x^{-4}$  term. HOWEVER i do not see the other term in  $x^{-2}$ .

## I Intro

### A diagram a

flavour nonsinglet string operator and wilson line

$$Op. = \bar{\psi}(x)\lambda^a\not{x}[x, 0]\psi(0) \quad (1.1)$$

$$[x, y] = P \exp \left[ ig \int_0^1 du (x - y)_\mu A^\mu(xu + y(1 - u)) \right] \quad (1.2)$$

From diagram i get a similar (but not the same) expression. Also i should think about what this diagram is - is it sort of "VEV" of operator, with not vacuum but quark states?

Brauns expression:

$$\int_0^1 du \bar{\psi}_c(x) i x_\mu g A_q^\mu \not{x} \lambda^a \psi_q(0) \left[ ig \int dz \bar{\psi}_q(z) A_q(z) \psi(z) \right] \quad (1.3)$$

$$= \int_0^1 du \bar{\psi}_c(x) i x_\mu g A_q^\mu \not{x} \lambda^a \psi_q(0) \overbrace{ig \int dz \bar{\psi}_q(z) A_q(z) \psi(z)} \quad (1.4)$$

$$= -\frac{i\alpha_s}{2\pi} C_F \bar{\psi}_c(x) \lambda^a \not{x} \int_0^1 du \int dz \frac{\Gamma(d/2)\Gamma(d/2-1)\not{x}\not{z}}{(-z^2)^{\frac{d}{2}} \left( (-ux - z)^2 \right)^{\frac{d}{2}-1}} \psi_c(z) \quad (1.5)$$

I need to verify this by looking up propagators for d dimensional spacetime, and hereby check it. going on from this point we use integration tricks(Denominator + shift):

$$-\frac{i\alpha_s}{2\pi} C_F \bar{\psi}_c(x) \lambda^a \not{x} \int_0^1 du \int_0^1 dv v^{\frac{d}{2}-2} \bar{v}^{\frac{d}{2}-1} \int dz \frac{\Gamma(d-1)(\not{x} + uv\not{z})\not{z}}{(-z^2 - v\bar{v}u^2x^2)^{d-1}} \psi_c(z + uvx) \quad (1.6)$$

Go on with again d dim integral formulas therefore we have to expand the field psi so the numerator structure is

$$\not{x} (\not{x}\not{x} + uvx^2) (\psi_c(uvx) + z^\alpha \partial_\alpha \psi_c(uvx) + \dots) \quad (1.7)$$

$$= (-x^2 \not{x} + 2(x \cdot z) \not{x} + uvx^2 \not{x}) (\psi_c(uvx) + z^\alpha \partial_\alpha \psi_c(uvx) + \dots) \quad (1.8)$$

numerator(dropping asym terms):

$$uvx^2 \psi_c(uvx) + (-x^2 \not{x} + 2(x \cdot z) \not{x}) z^\alpha \partial_\alpha \psi_c(uvx) \quad (1.9)$$

This means, combined with the denomoinator i have the three terms

$$1: \int dz \frac{uvx^2 \not{x} \psi_c}{(-z^2 - v\bar{v}u^2 x^2)^{d-1}} \quad (1.10)$$

$$= v^{2-\frac{d}{2}} \bar{v}^{1-\frac{d}{2}} u^{3-d} v i \pi^2 \frac{\Gamma(d/2 - 1)}{\Gamma(d-1)} \frac{\not{x} \psi_c}{(-x^2)^{\frac{d}{2}-2}} \quad (1.11)$$

$$2: \int dz \frac{-x^2 \gamma_\beta z^\beta z^\alpha \partial_\alpha \psi_c}{(-z^2 - v\bar{v}u^2 x^2)^{d-1}} \quad (1.12)$$

$$= (v\bar{v})^{2-\frac{d}{2}} \frac{i}{2} \pi^2 \frac{\Gamma(\frac{d}{2} - 2)}{\Gamma(d-1)} \frac{x^2 \gamma^\alpha \partial_\alpha \psi_c}{(-u^2 x^2)^{\frac{d}{2}-2}} \quad (1.13)$$

$$3: \int dz \frac{2x_\beta z^\beta z^\alpha \partial_\alpha \psi_c}{(-z^2 - v\bar{v}u^2 x^2)^{d-1}} \quad (1.14)$$

$$= (v\bar{v})^{2-\frac{d}{2}} i \pi^2 \frac{\Gamma(\frac{d}{2} - 2)}{\Gamma(d-1)} \frac{x^\alpha \partial_\alpha \psi_c}{(-u^2 x^2)^{\frac{d}{2}-2}} \quad (1.15)$$

All terms are to be multiplied with

$$-\frac{i\alpha_s}{2\pi} C_F \bar{\psi}_c(x) \lambda^a \int_0^1 du \int_0^1 dv v^{\frac{d}{2}-2} \bar{v}^{\frac{d}{2}-1} \Gamma(d-1) \quad (1.16)$$

Since the 2nd term is  $x^{-2}$  - power, it is dropped already.(?)

Remaining:

$$\frac{\alpha_s \pi^2}{2\pi} C_F \bar{\psi}_c(x) \lambda^a \int_0^1 du \int_0^1 dv \left( \frac{\bar{v} \Gamma(\frac{d}{2} - 2)}{(-u^2 x^2)^{\frac{d}{2}-2}} x^\alpha \partial_\alpha \psi_c(uvx) + \frac{\Gamma(\frac{d}{2} - 1)}{(-x^2)^{\frac{d}{2}-2}} u^{3-d} \not{x} \psi_c(uvx) \right) \quad (1.17)$$

This agrees with Vladimir Braun. Then we go on, like it is written in the paper. We drop the 2nd term because it is UV divergent for  $u \rightarrow 0$  in  $d = 4$  dimensions. With the first term we can manipulate  $x^\alpha \partial_\alpha \psi_c(uvx) = \frac{d}{du} \psi_c(uvx)$  (typo in Brauns Paper(?)) and integrate by parts: Have (for  $d = 4$ ):

$$\frac{\alpha_s \pi^2}{2\pi} C_F \bar{\psi}_c(x) \lambda^a \int_0^1 du \int_0^1 dv \frac{\bar{v} \Gamma(\frac{d}{2} - 2)}{(-x^2)^{\frac{d}{2}-2}} u^{d-4} \frac{d}{du} \psi_c(uvx) \quad (1.18)$$

$$= \frac{\alpha_s \pi^2}{2\pi} C_F \bar{\psi}_c(x) \lambda^a \int_0^1 dv \frac{\bar{v} \Gamma(\frac{d}{2} - 2)}{(-x^2)^{\frac{d}{2}-2}} (\psi_c(vx) - \psi_c(0)) \quad (1.19)$$

## B diagram b

Now the second matrix element

$$\int_0^1 du \int dz \bar{\psi}_c(z) i g A_q(z) \psi_q(z) \bar{\psi}_q(0) i g x_\mu A^\mu(\bar{u}x) \not{x} \lambda^a \psi_c(-x) \quad (1.20)$$

$$= \frac{-\alpha_s}{2\pi^3} \int_0^1 du \int_0^1 dv \int dz \bar{\psi}_c(z + \bar{u}vx) \lambda^a \Gamma(d-1) (v\bar{v})^{\frac{d}{2}-2} \bar{v} \frac{\not{x}(\not{x} + \bar{u}v\not{x})\not{x}}{(z^2 + v\bar{v}u^2x^2)^{d-1}} \psi_c(-x) \quad (1.21)$$

Expand field psi + Numerator analysis:

$$(\bar{\psi}_c(\bar{u}vx) + z^\alpha \partial_\alpha \bar{\psi}_c(\bar{u}vx)) \not{x} (\not{x} + \bar{u}v\not{x}) \not{x} = (\bar{\psi}_c(\bar{u}vx) + z^\alpha \partial_\alpha \bar{\psi}_c(\bar{u}vx)) (-\not{x}^2 + 2(z \cdot x)\not{x} + \bar{u}vx^2\not{x}) \quad (1.22)$$

$$= \bar{\psi}_c(\bar{u}vx) \bar{u}vx^2\not{x} + z^\alpha \partial_\alpha \bar{\psi}_c(\bar{u}vx) (-\not{x}^2 + 2(z \cdot x)\not{x}) \quad (1.23)$$

these terms yield

$$1: \int dz \frac{\bar{\psi}_c \bar{u}vx^2\not{x}}{(z^2 + v\bar{v}u^2x^2)^{d-1}} \quad (1.24)$$

$$= -v^{2-\frac{d}{2}} \bar{v}^{1-\frac{d}{2}} \bar{u}u^{2-d} v i \pi^2 \frac{\Gamma(d/2-1)}{\Gamma(d-1)} \frac{\not{x}\psi_c}{(-x^2)^{\frac{d}{2}-2}} \quad (1.25)$$

$$2: \int dz \frac{-x^2 z^\beta z^\alpha \partial_\alpha \bar{\psi}_c \gamma_\beta}{(z^2 + v\bar{v}u^2x^2)^{d-1}} \quad (1.26)$$

$$= \text{dropped}(1/x^2) \quad (1.27)$$

$$3: \int dz \frac{2x_\beta z^\beta z^\alpha \partial_\alpha \bar{\psi}_c}{(z^2 + v\bar{v}u^2x^2)^{d-1}} \quad (1.28)$$

$$= (v\bar{v})^{2-\frac{d}{2}} i \pi^2 \frac{\Gamma(\frac{d}{2}-2)}{\Gamma(d-1)} \frac{x^\alpha \partial_\alpha \bar{\psi}_c}{(-u^2x^2)^{\frac{d}{2}-2}} \quad (1.29)$$

merge it together, using only the third term since 2nd is of higher power and first term UV divergent ( $u \rightarrow 0$ ).

$$\frac{-i\alpha_s}{2\pi} \int_0^1 du \int_0^1 dv x^\alpha \partial_\alpha \bar{\psi}_c(\bar{u}vx) \lambda^a \bar{v}u^{4-d} \frac{\Gamma(\frac{d}{2}-2)}{(-x^2)^{\frac{d}{2}-2}} \psi_c(-x) \quad (1.30)$$

$$= \frac{-i\alpha_s}{2\pi} \int_0^1 dv (\bar{\psi}_c(0) - \bar{\psi}_c(-vx)) \lambda^a \bar{v} \frac{\Gamma(\frac{d}{2}-2)}{(-x^2)^{\frac{d}{2}-2}} \psi_c(-x) \quad (1.31)$$

factor of i missing somewhere in the beginning. why? every propagator gives one,  
then from loop integral 1, 2 from the start, 5 in total?

shifting back the arguments of the functions by +x, receive:

$$\frac{-i\alpha_s}{2\pi} \int_0^1 dv (\bar{\psi}_c(x) - \bar{\psi}_c(\bar{v}x)) \lambda^a \bar{v} \frac{\Gamma(\frac{d}{2}-2)}{(-x^2)^{\frac{d}{2}-2}} \psi_c(0) \quad (1.32)$$

## C diagram c

start with formula:

$$[D : c] = \int d^d z \int d^d y i g \bar{\psi}(z) \not{A}(z) \psi(z) \bar{\psi}(x) \lambda^a \not{x} \psi(0) i g \bar{\psi}(y) \not{A}(y) \psi(y) \quad (1.33)$$

$$= -i^5 g^2 \frac{\Gamma^2\left(\frac{d}{2}\right) \Gamma\left(\frac{d}{2} - 1\right)}{\left(2\pi^{\frac{d}{2}}\right)^2 4\pi^{\frac{d}{2}}} \delta^{ab} \eta^{\mu\nu} \int d^d z \int d^d y \quad (1.34)$$

$$\bar{\psi}(z) \lambda^a t^a t^b \gamma_\mu \frac{\not{z} - \not{x}}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}}} \not{x} \frac{-\not{y}}{(-y^2 + i\epsilon)^{\frac{d}{2}}} \gamma_\nu \frac{1}{(-(y-z)^2 + i\epsilon)^{\frac{d}{2}-1}} \psi(y) \quad (1.35)$$

$$= \frac{ig^2}{8\pi^{\frac{3d}{2}}} C_F \int d^d z \int d^d y \bar{\psi}(z) \lambda^a \gamma_\mu (\not{z} - \not{x}) \not{x} \not{y} \gamma^\mu \psi(y) \quad (1.36)$$

$$\int [d\alpha d\beta d\gamma] \frac{\alpha^{\frac{d}{2}-1} \beta^{\frac{d}{2}-1} \gamma^{\frac{d}{2}-2} \Gamma\left(\frac{3d}{2} - 1\right)}{(-\alpha(z-x)^2 - \beta y^2 - \gamma(y-z)^2 + i\epsilon)^{\frac{3d}{2}-1}} \quad (1.37)$$

$$\int [d\alpha d\beta d\gamma] = \int_0^1 d\alpha \int_0^1 d\beta \int_0^1 d\gamma \delta(\alpha + \beta + \gamma) \quad (1.38)$$

Start denominator algebra with *Mathematica* (shift y,z by y,z plus ux,wx and solve u and w such that there is no mix term xz, xy). Find the following shifts:

$$z \rightarrow z + \frac{\alpha\beta + \gamma\alpha}{\Lambda} x \quad (1.39)$$

$$y \rightarrow y + \frac{\gamma\alpha}{\Lambda} x \quad (1.40)$$

$$\Lambda = \alpha\beta + \beta\gamma + \gamma\alpha \quad (1.41)$$

Transforms the denominator to

$$(-\alpha(z-x)^2 - \beta y^2 - \gamma(y-z)^2 + i\epsilon) \quad (1.42)$$

$$\rightarrow \left( \alpha y^2 + \beta z^2 + 2\gamma yz - \frac{\alpha\beta\gamma}{\Lambda} x^2 + i\epsilon \right) \quad (1.43)$$

for the numerator (gamma structure) we only take terms which give us the correct power/dimension, meaning no  $x^2$ !

$$\gamma_\mu (\not{z} - \not{x}) \not{x} \not{y} \gamma^\mu \rightarrow \gamma_\mu \not{x} \not{x} \not{y} \gamma^\mu \quad (1.44)$$

Now we shift such that we can first integrate over  $y$  (isolate  $y^2$ ) and then over  $z$ . now shift not by  $x$  but by  $z$ ! again *Mathematica*

$$y \rightarrow y - \frac{\gamma}{\alpha} z \quad (1.45)$$

$$\left( \alpha y^2 + \beta z^2 + 2\gamma yz - \frac{\alpha\beta\gamma}{\Lambda} x^2 + i\epsilon \right) \rightarrow \left( \frac{\Lambda}{\alpha} z^2 - \alpha y^2 - \frac{\alpha\beta\gamma}{\Lambda} x^2 + i\epsilon \right) \quad (1.46)$$

$$= (-\alpha) \left( y^2 - \left( \frac{\Lambda}{\alpha^2} z^2 - \frac{\beta\gamma}{\Lambda} x^2 \right) + i\epsilon \right) \quad (1.47)$$

at the same time the numerator:

$$\gamma_\mu \not{x} \not{y} \gamma^\mu \rightarrow \gamma_\mu \not{x} \not{x} \left( \psi - \frac{\gamma}{\alpha} \not{x} \right) \gamma^\mu \quad (1.48)$$

$$\rightarrow -\frac{\gamma}{\alpha} \gamma_\mu \not{x} \not{x} \not{x} \gamma^\mu \quad (1.49)$$

$$= -\frac{\gamma}{\alpha} \gamma_\mu \gamma_\tau \not{x} \gamma_\eta \gamma^\mu z^\tau z^\eta \quad (1.50)$$

So i have the following relevant integral(if i made no mistakes the numerator structure is not relevant.):

$$(-\alpha)^{1-\frac{3d}{2}} \int d^d y \frac{1}{\left( y^2 - \left( \frac{\Lambda}{\alpha^2} z^2 - \frac{\beta\gamma}{\Lambda} x^2 \right) + i\epsilon \right)^{\frac{3d}{2}-1}} = i\pi^{\frac{d}{2}} \alpha^{1-\frac{3d}{2}} \frac{\Gamma(d-1)}{\Gamma(\frac{3d}{2}-1)} \frac{1}{\left( \frac{\Lambda}{\alpha^2} z^2 - \left( \frac{\beta\gamma}{\Lambda} x^2 - i\epsilon \right) \right)^{d-1}} \quad (1.51)$$

Now pull out coefficient of  $z^2$  and perform same integral again, now watch numerator!

$$\int d^d z \frac{-\frac{\gamma}{\alpha} \left( \frac{\Lambda}{\alpha^2} \right)^{1-d} \gamma_\mu \gamma_\tau \not{x} \gamma_\eta \gamma^\mu z^\tau z^\eta}{\left( z^2 - \left( \frac{\alpha^2 \beta\gamma}{\Lambda^2} x^2 - i\epsilon \right) \right)^{d-1}} = +\frac{\gamma}{\alpha} \left( -\frac{\Lambda}{\alpha^2} \right)^{1-d} \gamma_\mu \gamma_\tau \not{x} \gamma^\tau \gamma^\mu \frac{i\pi^{\frac{d}{2}} \Gamma(\frac{d}{2}-2)}{2 \Gamma(d-1)} \frac{1}{\left( \frac{\alpha^2 \beta\gamma}{\Lambda^2} x^2 - i\epsilon \right)^{\frac{d}{2}-2}} \quad (1.52)$$

At this point realize that all the previous stuff is pathological, since i forgot to expand the fields. Bad. !!!!!!!!!!! Collect :

$$[D : c] = \frac{ig^2}{8\pi^{\frac{3d}{2}}} C_F \int [d\alpha d\beta d\gamma] \alpha^{\frac{d}{2}-1} \beta^{\frac{d}{2}-1} \gamma^{\frac{d}{2}-2} \Gamma\left(\frac{3d}{2}-1\right) \quad (1.53)$$

$$\int d^d z \int d^d y \bar{\psi}\left(z + \frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) \lambda^a \frac{\gamma_\mu \left( \not{x} - \frac{\beta\gamma}{\Lambda} \not{x} \right) \not{x} \left( \psi + \frac{\gamma\alpha}{\Lambda} \not{x} \right) \gamma^\mu}{\left( \alpha y^2 + \beta z^2 + 2\gamma yz - \frac{\alpha\beta\gamma}{\Lambda} x^2 + i\epsilon \right)^{\frac{3d}{2}-1}} \psi\left(y + \frac{\gamma\alpha}{\Lambda} x\right) \quad (1.54)$$

$$= \frac{ig^2}{8\pi^{\frac{3d}{2}}} C_F \int [d\alpha d\beta d\gamma] \alpha^{-d} \beta^{\frac{d}{2}-1} \gamma^{\frac{d}{2}-2} \Gamma\left(\frac{3d}{2}-1\right) (-1)^{\frac{3d}{2}-1} \quad (1.55)$$

$$\int d^d z \int d^d y \bar{\psi}\left(z + \frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) \lambda^a \frac{\gamma_\mu \left( \not{x} - \frac{\beta\gamma}{\Lambda} \not{x} \right) \not{x} \left( \psi - \frac{\gamma}{\alpha} \not{x} + \frac{\gamma\alpha}{\Lambda} \not{x} \right) \gamma^\mu}{\left( y^2 - \left( \frac{\Lambda}{\alpha^2} z^2 - \frac{\beta\gamma}{\Lambda} x^2 - i\epsilon \right) \right)^{\frac{3d}{2}-1}} \psi\left(y - \frac{\gamma}{\alpha} z + \frac{\gamma\alpha}{\Lambda} x\right) \quad (1.56)$$

the numerator analysis, again: first expand the fields

$$\psi\left(y - \frac{\gamma}{\alpha} z + \frac{\gamma\alpha}{\Lambda} x\right) = \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) + y^\rho \partial_\rho \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) - \frac{\gamma}{\alpha} z^\rho \partial_\rho \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) + \dots \quad (1.57)$$

$$\bar{\psi}\left(z + \frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) = \bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) + z^\rho \partial_\rho \bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) + \dots \quad (1.58)$$

Numerator:

$$\left( \bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) + z^\rho \partial_\rho \bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) + z^\rho z^\zeta \partial_\rho \partial_\zeta \bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda} x\right) \right) \lambda^a \gamma_\mu \left( \not{x} - \frac{\beta\gamma}{\Lambda} \not{x} \right) \not{x} \left( \psi - \frac{\gamma}{\alpha} \not{x} + \frac{\gamma\alpha}{\Lambda} \not{x} \right) \gamma^\mu \quad (1.59)$$

$$\left( \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) + y^\sigma \partial_\sigma \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) - \frac{\gamma}{\alpha} z^\sigma \partial_\sigma \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) + (y^\sigma z^\xi + y^\sigma y^\xi + z^\sigma z^\xi) \partial_\sigma \partial_\xi \psi\left(\frac{\gamma\alpha}{\Lambda} x\right) \right) \quad (1.60)$$

this are in total  $3 \cdot 2 \cdot 3 \cdot 4 = 72$  terms. Then some immediately cancel by symmetry. Still a lot terms. I will

not write down all of them. now select only following terms This restriction only approves the term with constant fields...

$$\bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda}x\right)\lambda^a\gamma_\mu\not{x}\not{\beta}\frac{\gamma}{\alpha}\not{x}\gamma^\mu\psi\left(\frac{\gamma\alpha}{\Lambda}x\right) \quad (1.61)$$

which corresponds to the previous calculation where i did not expand the fields. Do it again: have the calculation

$$[D : c] = \frac{ig^2}{8\pi^{\frac{3d}{2}}}C_F \int [d\alpha d\beta d\gamma] \alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}\Gamma\left(\frac{3d}{2}-1\right)(-1)^{\frac{3d}{2}-1} \quad (1.62)$$

$$\int d^dz \int d^dy \bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda}x\right)\lambda^a \frac{-\frac{\gamma}{\alpha}\gamma_\mu\not{x}\not{\beta}\not{x}\gamma^\mu}{\left(y^2 - \left(\frac{\Lambda}{\alpha^2}z^2 - \frac{\beta\gamma}{\Lambda}x^2 - i\epsilon\right)\right)^{\frac{3d}{2}-1}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right) \quad (1.63)$$

$$= \frac{ig^2}{8\pi^{\frac{3d}{2}}}C_F \int [d\alpha d\beta d\gamma] \alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}\Gamma\left(\frac{3d}{2}-1\right)(-1)^{\frac{3d}{2}-1} \quad (1.64)$$

$$\int d^dz \frac{(-1)^{\frac{3d}{2}-1}i\pi^{\frac{d}{2}}\Gamma(d-1)}{\Gamma\left(\frac{3d}{2}-1\right)}\bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda}x\right)\lambda^a \frac{-\frac{\gamma}{\alpha}\gamma_\mu\not{x}\not{\beta}\not{x}\gamma^\mu}{\left(\frac{\Lambda}{\alpha^2}z^2 - \left(\frac{\beta\gamma}{\Lambda}x^2 + i\epsilon\right)\right)^{d-1}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right) \quad (1.65)$$

$$= \frac{ig^2}{8\pi^{\frac{3d}{2}}}C_F \int [d\alpha d\beta d\gamma] \alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}\Gamma\left(\frac{3d}{2}-1\right)(-1)^{\frac{3d}{2}-1}\left(\frac{\Lambda}{\alpha^2}\right)^{1-d} \quad (1.66)$$

$$\int d^dz \frac{(-1)^{\frac{3d}{2}-1}i\pi^{\frac{d}{2}}\Gamma(d-1)}{\Gamma\left(\frac{3d}{2}-1\right)}\bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda}x\right)\lambda^a \frac{-\frac{\gamma}{\alpha}\gamma_\mu\gamma_\tau\not{x}\gamma_\theta\gamma^\mu z^\tau z^\theta}{\left(z^2 - \left(\frac{\alpha^2\beta\gamma}{\Lambda^2}x^2 + i\epsilon\right)\right)^{d-1}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right) \quad (1.67)$$

$$= \frac{ig^2}{8\pi^{\frac{d}{2}}}C_F \int [d\alpha d\beta d\gamma] \alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}\left(\frac{\Lambda}{\alpha^2}\right)^{1-d} \quad (1.68)$$

$$\frac{(-1)^{4d-1}i^2\Gamma\left(\frac{d}{2}-2\right)}{2}\bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda}x\right)\lambda^a \frac{-\frac{\gamma}{\alpha}\gamma_\mu\gamma_\theta\not{x}\gamma^\theta\gamma^\mu}{\left(\frac{\alpha^2\beta\gamma}{\Lambda^2}x^2 + i\epsilon\right)^{\frac{d}{2}-2}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right) \quad (1.69)$$

$$= \frac{+ig^2}{16\pi^{\frac{d}{2}}}C_F \int [d\alpha d\beta d\gamma] \Lambda^{-3}\alpha^1\beta^1\gamma^1 \quad (1.70)$$

$$(-1)^{4d-1}\Gamma\left(\frac{d}{2}-2\right)\bar{\psi}\left(\frac{\alpha\beta + \gamma\alpha}{\Lambda}x\right)\lambda^a \frac{(d-2)^2\not{x}}{(x^2 + i\epsilon)^{\frac{d}{2}-2}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right) \quad (1.71)$$

$$= \frac{+ig^2}{16\pi^{\frac{d}{2}}}C_F \int [d\alpha' d\beta' d\gamma'] \quad (1.72)$$

$$(-1)^{4d-1}\Gamma\left(\frac{d}{2}-2\right)\bar{\psi}(\alpha'x)\lambda^a \frac{(d-2)^2\not{x}}{(x^2 + i\epsilon)^{\frac{d}{2}-2}}\psi(\beta'x) \quad (1.73)$$



I can also take the term (as an example)

$$-\frac{\alpha\beta\gamma^2}{\Lambda^2}\bar{\psi}\lambda^a\gamma_\mu\not{x}\not{x}\not{x}\gamma^\mu\psi\rightarrow\quad (1.74)$$

$$[D : c] = \frac{ig^2}{8\pi^{\frac{3d}{2}}}C_F\int[d\alpha d\beta d\gamma]\alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}\Gamma\left(\frac{3d}{2}-1\right)(-1)^{\frac{3d}{2}-1}\quad (1.75)$$

$$\int d^dz\int d^dy\bar{\psi}\left(\frac{\alpha\beta+\gamma\alpha}{\Lambda}x\right)\lambda^a\frac{-\frac{\alpha\beta\gamma^2}{\Lambda^2}\gamma_\mu x^2\not{x}\gamma^\mu}{\left(y^2-\left(\frac{\Lambda}{\alpha^2}z^2-\frac{\beta\gamma}{\Lambda}x^2-i\epsilon\right)\right)^{\frac{3d}{2}-1}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right)\quad (1.76)$$

$$= \frac{ig^2}{8\pi^{\frac{3d}{2}}}C_F\int[d\alpha d\beta d\gamma]\alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}(-1)^{3d}\left(\frac{\Lambda}{\alpha^2}\right)^{1-d}\quad (1.77)$$

$$\int d^dz i\pi^{\frac{d}{2}}\Gamma(d-1)\bar{\psi}\left(\frac{\alpha\beta+\gamma\alpha}{\Lambda}x\right)\lambda^a\frac{-\frac{\alpha\beta\gamma^2}{\Lambda^2}\gamma_\mu x^2\not{x}\gamma^\mu}{\left(z^2-\left(\frac{\alpha^2\beta\gamma}{\Lambda^2}x^2+i\epsilon\right)\right)^{d-1}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right)\quad (1.78)$$

$$= \frac{ig^2}{8\pi^{\frac{d}{2}}}C_F\int[d\alpha d\beta d\gamma]\alpha^{-d}\beta^{\frac{d}{2}-1}\gamma^{\frac{d}{2}-2}(-1)^{4d-1}\left(\frac{\Lambda}{\alpha^2}\right)^{1-d}\quad (1.79)$$

$$i^2\Gamma\left(\frac{d}{2}-1\right)\bar{\psi}\left(\frac{\alpha\beta+\gamma\alpha}{\Lambda}x\right)\lambda^a\frac{-\frac{\alpha\beta\gamma^2}{\Lambda^2}\gamma_\mu x^2\not{x}\gamma^\mu}{\left(\frac{\alpha^2\beta\gamma}{\Lambda^2}x^2+i\epsilon\right)^{\frac{d}{2}-1}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right)\quad (1.80)$$

$$= \frac{-ig^2}{8\pi^{\frac{d}{2}}}C_F\int[d\alpha d\beta d\gamma]\alpha^1\beta^1\gamma^1(-1)^{4d-1}\Lambda^{-3}\quad (1.81)$$

$$\Gamma\left(\frac{d}{2}-1\right)\bar{\psi}\left(\frac{\alpha\beta+\gamma\alpha}{\Lambda}x\right)\lambda^a\frac{(d-2)\not{x}}{(x^2+i\epsilon)^{\frac{d}{2}-2}}\psi\left(\frac{\gamma\alpha}{\Lambda}x\right)\quad (1.82)$$

$$= \frac{-ig^2}{8\pi^{\frac{d}{2}}}C_F\int[d\alpha'd\beta'd\gamma'](-1)^{4d-1}\quad (1.83)$$

$$\Gamma\left(\frac{d}{2}-1\right)\bar{\psi}(\bar{\alpha}'x)\lambda^a\frac{(d-2)\not{x}}{(x^2+i\epsilon)^{\frac{d}{2}-2}}\psi(\beta'x)\quad (1.84)$$

and this is of same power with  $x^2$  in denominator.

other possibility which now should give different order:

$$\frac{\gamma\alpha}{\Lambda}z^\rho\partial_\rho\bar{\psi}\lambda^a\gamma_\mu\not{x}\not{x}\not{x}\gamma^\mu\psi\quad (1.85)$$

then

$$[D : c] = \frac{ig^2}{8\pi^{\frac{3d}{2}}} C_F \int [d\alpha d\beta d\gamma] \alpha^{-d} \beta^{\frac{d}{2}-1} \gamma^{\frac{d}{2}-2} \Gamma\left(\frac{3d}{2} - 1\right) (-1)^{\frac{3d}{2}-1} \quad (1.86)$$

$$\int d^d z \int d^d y \partial_\rho \bar{\psi} \left( \frac{\alpha\beta + \gamma\alpha}{\Lambda} x \right) \lambda^a \frac{\frac{\gamma\alpha}{\Lambda} z^\rho \gamma_\mu \not{x} x^2 \gamma^\mu}{\left( y^2 - \left( \frac{\Lambda}{\alpha^2} z^2 - \frac{\beta\gamma}{\Lambda} x^2 - i\epsilon \right) \right)^{\frac{3d}{2}-1}} \psi \left( \frac{\gamma\alpha}{\Lambda} x \right) \quad (1.87)$$

$$= \frac{ig^2}{8\pi^{\frac{3d}{2}}} C_F \int [d\alpha d\beta d\gamma] \alpha^{-d} \beta^{\frac{d}{2}-1} \gamma^{\frac{d}{2}-2} (-1)^{\frac{6d}{2}-2} \left( \frac{\Lambda}{\alpha^2} \right)^{1-d} \quad (1.88)$$

$$\int d^d z i\pi^{\frac{d}{2}} \Gamma(d-1) \partial_\rho \bar{\psi} \left( \frac{\alpha\beta + \gamma\alpha}{\Lambda} x \right) \lambda^a \frac{\frac{\gamma\alpha}{\Lambda} z^\rho z^\theta \gamma_\mu \gamma_\theta x^2 \gamma^\mu}{\left( z^2 - \left( \frac{\beta\gamma}{\Lambda} x^2 + i\epsilon \right) \right)^{d-1}} \psi \left( \frac{\gamma\alpha}{\Lambda} x \right) \quad (1.89)$$

$$= \frac{ig^2}{8\pi^{\frac{d}{2}}} C_F \int [d\alpha d\beta d\gamma] \alpha^{-d} \beta^{\frac{d}{2}-1} \gamma^{\frac{d}{2}-2} (-1)^{4d-1} \left( \frac{\Lambda}{\alpha^2} \right)^{1-d} \quad (1.90)$$

$$\frac{i^2 \Gamma\left(\frac{d}{2} - 2\right)}{2} \partial_\rho \bar{\psi} \left( \frac{\alpha\beta + \gamma\alpha}{\Lambda} x \right) \lambda^a \frac{\frac{\gamma\alpha}{\Lambda} \gamma_\mu \gamma^\rho x^2 \gamma^\mu}{\left( \frac{\beta\gamma}{\Lambda} x^2 + i\epsilon \right)^{\frac{d}{2}-2}} \psi \left( \frac{\gamma\alpha}{\Lambda} x \right) \quad (1.91)$$

And this term is now of order  $x^2$  in total, which is a correction only. this term will be dropped. So i think i do not care about derivatives on quantum fields, but every x,y,z pair on top decreases the power of the denominator.

Schematically

$$\int dx^d z \int d^d y \frac{z^{2k} y^{2l} x^{2m}}{(y^2 - z^2 + x^2)^{\frac{3d}{2}-1}} \rightarrow \int dx^d z \frac{z^{2k} x^{2m}}{(z^2 + x^2)^{d-1-l}} \quad (1.92)$$

$$\rightarrow \frac{x^{2m}}{(x^2)^{\frac{d}{2}-1-l-k}} \quad (1.93)$$

$$= (x^2)^{m+l+k+1-\frac{d}{2}} \quad (1.94)$$

We are looking for  $2 = m + l + 1 + k$  and i omitt the 1 slashed  $x$ , which is always there from the middle term, where i have no choice. ( means I can have  $x^2(m=1)$ ,  $x^2 \not{x}(m=1)$  or  $\not{x}(m=0)$  only)

Applying this analysis I i only have the two terms which are the first two examples. All other terms with give higher power in  $x^2$  and are hence a correction

## II twist2

have operator (fraction):

$$O_q^1(z) = [-\infty n, zn] q(zn) \quad (2.1)$$

For one loop I have to take linear term of wilson line and inlcude first order of interaction Hamiltonian. Thus

$$A = \left[ -ig \int_{-\infty}^z d\sigma B^+(\sigma n) \right] q(zn) \left( ig \int d^d x \bar{q}(x) \not{B}(x) q(x) \right) \quad (2.2)$$

I do not write the contractions since they are clear and I am lazy. result: we shift  $z$  to 0 because it does not matter and makes it easier. ACTUALLY:: this is a dangerous move. doing it carefully and not much more difficult, one shifts integration variable  $x$  such that  $z$  is dropped in propagators. However it remains in integration in wilson line AND also in quark field shift. thus care and understand that in the end the quark field position is shifted by  $+z$ . Important mistake I made at some point: the integration limit 0 is NOT SHIFTED! it is 0 for any  $z$ ! Mid calculation shift :  $x = x + \bar{\alpha}\sigma n$  note that  $n$  is a light-like vector

$$A = -(ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x n^\mu \frac{-1}{4\pi^{\frac{d}{2}}} \frac{\Gamma(\frac{d}{2}-1)}{(-(x-\sigma n)^2 + i\epsilon)^{\frac{d}{2}-1}} \frac{-i\Gamma(\frac{d}{2})}{2\pi^{\frac{d}{2}}} \frac{\not{x}}{(-x^2 + i\epsilon)^{\frac{d}{2}}} \gamma_\mu q(x) \quad (2.3)$$

$$= -(ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x \frac{i\Gamma(\frac{d}{2}-1) \Gamma(\frac{d}{2})}{8\pi^d} \frac{\not{x}}{(-(x-\sigma n)^2 + i\epsilon)^{\frac{d}{2}-1} (-x^2 + i\epsilon)^{\frac{d}{2}}} \gamma^+ q(x) \quad (2.4)$$

$$= -(ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x \int_0^1 d\alpha \frac{i\Gamma(d-1)}{8\pi^d} \frac{\alpha^{\frac{d}{2}-1} \bar{\alpha}^{\frac{d}{2}-2} \not{x} \gamma^+}{(-\bar{\alpha}(x-\sigma n)^2 - \alpha x^2 + i\epsilon)^{d-1}} q(x) \quad (2.5)$$

$$= -(ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x \int_0^1 d\alpha \frac{i\Gamma(d-1)}{8\pi^d} \frac{\alpha^{\frac{d}{2}-1} \bar{\alpha}^{\frac{d}{2}-2} \not{x} \gamma^+}{(-x^2 + i\epsilon)^{d-1}} q(x + \bar{\alpha}\sigma n) \quad (2.6)$$

here use the formula (copied from Alexeys notes)

$$\int d^d x \frac{x^\mu x^\nu}{(-x^2 + i\epsilon)^{d-1}} = \frac{-i\pi^{\frac{d}{2}}}{2(2 - \frac{d}{2})} \frac{\eta^{\mu\nu}}{\Gamma(d-1)} \quad (2.7)$$

own adaption for four elements in numerator:

$$\int d^d x \frac{x^\mu x^\nu x^\rho x^\sigma}{(-x^2 + i\epsilon)^d} = \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} (\eta^{\mu\nu}\eta^{\rho\sigma} + \eta^{\mu\rho}\eta^{\nu\sigma} + \eta^{\mu\sigma}\eta^{\nu\rho}) \quad (2.8)$$

expand quark field and have

$$A = -(ig)^2 C_F \int_{-\infty}^z d\sigma \int_0^1 d\alpha \frac{1}{8\pi^{\frac{d}{2}}} \frac{\alpha^{\frac{d}{2}-1} \bar{\alpha}^{\frac{d}{2}-2} \gamma^\nu \gamma^+}{2(2 - \frac{d}{2})} \partial_\nu q(\bar{\alpha}\sigma n) \quad (2.9)$$

## A left quark

$$\bar{O}_q^1(z) = \bar{q}(zn) [zn, -\infty n] \quad (2.10)$$

First order corr:Mid calculation shift :  $x = x + \alpha\sigma n$

$$\bar{A} = \left( ig \int d^d x \bar{q}(x) \not{B}(x) q(x) \right) \bar{q}(zn) [zn, -\infty n] \left[ ig \int_{-\infty}^z d\sigma B^+(\sigma n) \right] \quad (2.11)$$

$$= (ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x \bar{q}(x) \gamma^+ \frac{\Gamma(\frac{d}{2})}{2\pi^{\frac{d}{2}}} \frac{i\not{x}}{(-x^2 + i\epsilon)^{\frac{d}{2}}} \frac{-1}{4\pi^{\frac{d}{2}}} \frac{\Gamma(\frac{d}{2} - 1)}{(-(x - \sigma n)^2 + i\epsilon)^{\frac{d}{2} - 1}} \quad (2.12)$$

$$= -i(ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x \int_0^1 d\alpha \bar{q}(x) \frac{\Gamma(d-1)}{8\pi^d} \gamma^+ \not{x} \frac{\alpha^{\frac{d}{2}-2} \bar{\alpha}^{\frac{d}{2}-1}}{(-\bar{\alpha}x^2 - \alpha(x - \sigma n)^2 + i\epsilon)^{d-1}} \quad (2.13)$$

$$= -i(ig)^2 C_F \int_{-\infty}^z d\sigma \int d^d x \int_0^1 d\alpha \bar{q}(x + \alpha\sigma n) \frac{\Gamma(d-1)}{8\pi^d} \gamma^+ \not{x} \frac{\alpha^{\frac{d}{2}-2} \bar{\alpha}^{\frac{d}{2}-1}}{(-x^2 + i\epsilon)^{d-1}} \quad (2.14)$$

$$= -(ig)^2 C_F \int_{-\infty}^z d\sigma \int_0^1 d\alpha \partial_\mu \bar{q}(\alpha\sigma n) \frac{1}{8\pi^{\frac{d}{2}}} \gamma^+ \gamma^\mu \frac{\alpha^{\frac{d}{2}-2} \bar{\alpha}^{\frac{d}{2}-1}}{2(2 - \frac{d}{2})} \quad (2.15)$$

$$= \frac{\alpha_s}{4\pi} C_F \int_{-\infty}^z d\sigma \int_0^1 d\alpha \frac{\alpha^{\frac{d}{2}-2} \bar{\alpha}^{\frac{d}{2}-1}}{(2 - \frac{d}{2})} \partial_\mu \bar{q}(\alpha\sigma n) \gamma^+ \gamma^\mu \quad (2.16)$$

$$= \frac{2\alpha_s}{4\pi} C_F \int_{-\infty}^z d\sigma \int_0^1 d\alpha \frac{\bar{\alpha}}{\epsilon} \partial_+ \bar{q}(\alpha\sigma n) \quad (2.17)$$

perform the last integral is not trivial for me:

$$f(x) = \int \frac{d^d p}{(2\pi)^d} e^{i(p|x)} \int d^d x' e^{-i(p|x')} f(x') \quad (2.18)$$

$$\partial_+ \bar{q}(x) = \int \frac{d^d p}{(2\pi)^d} e^{i(p|x)} \int d^d x' e^{-i(p|x')} \partial_+ \bar{q}(x') \quad (2.19)$$

$$= \int \frac{d^d p}{(2\pi)^d} \int d^d x' (-ip_+) e^{i(p|x-x')} \bar{q}(x') \quad (2.20)$$

$$= \int \frac{d^d p}{(2\pi)^d} (-ip_+) e^{i(p|x)} \bar{q}(p) \quad (2.21)$$

then

$$\int_0^1 d\alpha \int_{-\infty}^0 d\sigma \bar{\alpha} \partial_+ \bar{q}((\alpha\sigma + z)n) = \int \frac{d^d p}{(2\pi)^d} e^{ip_+ z} \bar{q}(p) \int_0^1 d\alpha \int_{-\infty}^0 d\sigma \bar{\alpha} (-ip_+) e^{i\alpha\sigma p_+} \quad (2.22)$$

to deal with oscillating integral : introduce regulator for the integral,  $\delta$

$$= \int \frac{d^d p}{(2\pi)^d} e^{ip_+ z} \bar{q}(p) \int_0^1 d\alpha \int_{-\infty}^0 d\sigma \bar{\alpha}(-ip_+) e^{i\alpha\sigma p_+} e^{\delta\sigma} \quad (2.23)$$

$$= \int \frac{d^d p}{(2\pi)^d} e^{ip_+ z} \bar{q}(p) \int_0^1 d\alpha \frac{\bar{\alpha}(-ip_+)}{i\alpha p_+ + \delta} \quad (2.24)$$

$$= \int \frac{d^d p}{(2\pi)^d} e^{ip_+ z} \bar{q}(p) \int_0^1 d\alpha \frac{\alpha - 1}{\alpha + \frac{\delta}{ip_+}} \quad (2.25)$$

$$= \int \frac{d^d p}{(2\pi)^d} e^{ip_+ z} \bar{q}(p) \int_{\frac{\delta}{ip_+}}^{1 + \frac{\delta}{ip_+}} d\alpha \frac{\alpha - 1}{\alpha} \quad (2.26)$$

$$= \int \frac{d^d p}{(2\pi)^d} e^{ip_+ z} \bar{q}(p) \left( 1 - \ln \left( \frac{ip_+}{\delta} \right) \right) \quad (2.27)$$

CARE global minus difference to Alexeys result...

have operators

$$O_g^1(z) = [-\infty n, zn] F_{\mu+}(zn) \quad (2.28)$$

$$\bar{O}_g^1(zn) = F_{\mu+}(zn) [zn, -\infty n] \quad (2.29)$$

have 3-gluon vertex term: (taken from alexeys notes, can be derived by making separation of gluon fields into classical + quantum for full QCD lagrangian, also including ghost fields.

$$-ig A_\nu^{A'} \partial_\alpha B_\beta^{B'} B_\gamma^{C'} v_{A'B'C'}^{\nu\alpha\beta\gamma} \quad (2.30)$$

$$v_{ABC}^{\nu\alpha\beta\gamma} = f_{ABC} (2\eta^{\nu\beta} \eta^{\alpha\gamma} - \eta^{\nu\alpha} \eta^{\beta\gamma} - 2\eta^{\nu\gamma} \eta^{\alpha\beta}) \quad (2.31)$$

pod:

$$\bar{A}_q^{(1)} = -ig^2 C_A \int_{-\infty}^z d\sigma d^d x \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.32)$$

$$\left\{ (\partial_{z^\mu} n^\rho - \partial_{z^+} \eta^{\mu\rho}) \Delta^{\rho\gamma}(z-x) \partial_{x^\alpha} \Delta^{+\beta}(\sigma n - x) - (\partial_{z^\mu} n^\rho - \partial_{z^+} \eta^{\mu\rho}) \partial_{x^\alpha} \Delta^{\rho\beta}(z-x) \Delta^{+\gamma}(\sigma n - x) \right\} \quad (2.33)$$

$$= \frac{(-1)^3 i C_A g^2 \Gamma^2 \left( \frac{d}{2} - 1 \right)}{4^2 \pi^d} \int_{-\infty}^z d\sigma d^d x \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.34)$$

$$\left\{ (\partial_{z^\mu} n^\rho - \partial_{z^+} \eta^{\mu\rho}) \frac{\eta^{\rho\gamma}}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \partial_{x^\alpha} \frac{\eta^{+\beta}}{-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}-1}} \right. \quad (2.35)$$

$$\left. - (\partial_{z^\mu} n^\rho - \partial_{z^+} \eta^{\mu\rho}) \partial_{x^\alpha} \frac{\eta^{\rho\beta}}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \frac{\eta^{+\gamma}}{-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}-1}} \right\} \quad (2.36)$$

Aux:

$$\partial_{x^\alpha} \frac{1}{(-(z-x)^2 + i\epsilon)^n} = \frac{-2n(x-z)_\alpha}{(-(z-x)^2 + i\epsilon)^{n+1}} \quad (2.37)$$

$$\partial_{z^\mu} \partial_{x^\alpha} \frac{1}{(-(z-x)^2 + i\epsilon)^n} = \partial_{z^\mu} \frac{-2n(x-z)_\alpha}{(-(z-x)^2 + i\epsilon)^{n+1}} \quad (2.38)$$

$$= \frac{4n(n+1)(z-x)_\mu(x-z)_\alpha}{(-(z-x)^2 + i\epsilon)^{n+2}} + \frac{2n\eta_{\mu\alpha}}{(-(z-x)^2 + i\epsilon)^{n+1}} \quad (2.39)$$

$$\bar{A}_q^{(1)} = \frac{(-1)^3 i C_A g^2 \Gamma^2 \left(\frac{d}{2} - 1\right)}{4^2 \pi^d} \int_{-\infty}^z d\sigma d^d x \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.40)$$

$$\left\{ ((z-x)_\mu n^\rho - (z-x)_+ \eta^{\mu\rho}) \frac{-2 \left(\frac{d}{2} - 1\right) \eta^{\rho\gamma}}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{-2 \left(\frac{d}{2} - 1\right) (x - \sigma n) \eta^{+\beta}}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \right. \quad (2.41)$$

$$\left. - \eta^{\rho\beta} \left( \frac{4 \left(\frac{d}{2} - 1\right) \frac{d}{2} \left( (z-x)_\mu n^\rho - (z-x)_+ \eta^{\mu\rho} \right) (x-z)_\alpha}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}+1}} + \frac{2 \left(\frac{d}{2} - 1\right) (\eta_{\mu\alpha} n^\rho - \eta_{+\alpha} \eta^{\mu\rho})}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}}} \right) \frac{\eta^{+\gamma}}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}-1}} \right\} \quad (2.42)$$

$$= \frac{(-1)^3 i C_A g^2 \Gamma^2 \left(\frac{d}{2} - 1\right)}{4^2 \pi^d} \int_{-\infty}^z d\sigma d^d x \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.43)$$

$$\left\{ ((z-x)_\mu \eta^{+\gamma} - (z-x)_+ \eta^{\mu\gamma}) \frac{4 \left(\frac{d}{2} - 1\right)^2}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{(x - \sigma n) \eta^{+\beta}}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \right. \quad (2.44)$$

$$\left. - \left( \frac{4 \left(\frac{d}{2} - 1\right) \frac{d}{2} \left( (z-x)_\mu \eta^{+\beta} - (z-x)_+ \eta^{\mu\beta} \right) (x-z)_\alpha}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}+1}} + \frac{2 \left(\frac{d}{2} - 1\right) (\eta_{\mu\alpha} \eta^{+\beta} - \eta_{+\alpha} \eta^{\mu\beta})}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}}} \right) \frac{\eta^{+\gamma}}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}-1}} \right\} \quad (2.45)$$

To keep overview, treat 3 terms individually. now invent integration variables A:

$$FIRSTTERM = \frac{(-1)^3 i C_A g^2 \Gamma^2 \left(\frac{d}{2} - 1\right)}{4^2 \pi^d} \int_{-\infty}^z d\sigma d^d x \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.46)$$

$$((z-x)_\mu \eta^{+\gamma} - (z-x)_+ \eta^{\mu\gamma}) \frac{4 \left(\frac{d}{2} - 1\right)^2}{(-(z-x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{(x - \sigma n) \eta^{+\beta}}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \quad (2.47)$$

$$= \frac{(-1)^3 i C_A g^2 \Gamma^2 \left(\frac{d}{2} - 1\right)}{4^2 \pi^d} \int_{-\infty}^z d\sigma d^d x \int_0^1 d\alpha \frac{(\alpha \bar{\alpha})^{\frac{d}{2}-1} \Gamma(d) 4 \left(\frac{d}{2} - 1\right)^2}{\Gamma^2 \left(\frac{d}{2}\right)} \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.48)$$

$$((z-x)_\mu \eta^{+\gamma} - (z-x)_+ \eta^{\mu\gamma}) \frac{(x - \sigma n) \eta^{+\beta}}{(-\alpha(z-x)^2 - \bar{\alpha}(\sigma n - x)^2 + i\epsilon)^d} \quad (2.49)$$

$$= \frac{(-1)^3 i C_A g^2 \Gamma^2 \left(\frac{d}{2} - 1\right)}{4^2 \pi^d} \int_{-\infty}^z d\sigma d^d x \int_0^1 d\alpha \frac{(\alpha \bar{\alpha})^{\frac{d}{2}-1} \Gamma(d) 4 \left(\frac{d}{2} - 1\right)^2}{\Gamma^2 \left(\frac{d}{2}\right)} \bar{q}(0) A_\nu(x) v^{\nu\alpha\beta\gamma} \quad (2.50)$$

$$((z-x)_\mu \eta^{+\gamma} - (z-x)_+ \eta^{\mu\gamma}) \frac{(x - \sigma n) \eta^{+\beta}}{(-x^2 - 2\alpha\bar{\alpha}(z - \sigma n)^2 + i\epsilon)^d} \quad (2.51)$$

shift  $x : x \rightarrow x + \bar{\alpha}\sigma n + \alpha z$ .

## B field interaction correction

there are some extra contractions, more or less one extra diagram. It comes from taking into account Field interaction term, commutator of fields, and this then yields this term:

$$F_{\mu+}^{A'}(zn) [zn, -\infty n]_c e^{i \int dt H_I} \quad (2.52)$$

$$F_{\mu\nu}^{A'} = \partial_\mu A_\nu^{A'} - \partial_\nu A_\mu^{A'} + g f^{A'B'C'} A_\mu^{B'} A_\nu^{C'} \quad (2.53)$$

$$A_\mu \rightarrow A_\mu + B_\mu \quad (2.54)$$

take only last term of field tensor and have to additional terms:

$$g f^{A'B'C'} B_\mu^{B'}(zn) B_+^{C'}(zn) 1 \left( -ig \int d^d x A_\nu^D(x) \partial_\alpha B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.55)$$

$$\rightarrow g f^{A'B'C'} B_\mu^{B'}(zn) B_+^{C'}(zn) 1 \left( -ig \int d^d x A_\nu^D(x) \partial_\alpha B_\beta^E B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.56)$$

$$+ g f^{A'B'C'} B_\mu^{B'}(zn) B_+^{C'}(zn) 1 \left( -ig \int d^d x A_\nu^D(x) \partial_\alpha B_\beta^E B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.57)$$

the first diagram is:

$$-ig^2 f^{A'B'C'} \int d^d x \partial_{x^\alpha} \Delta_{\mu\beta}(zn-x) \delta^{B'E} \Delta_{+\gamma}(zn-x) \delta^{FC'} A_\nu^D(x) v^{\nu\alpha\beta\gamma} f^{DEF} \quad (2.58)$$

$$= \frac{-ig^2(-1)^2\Gamma^2\left(\frac{d}{2}-1\right)}{4^2\pi^d} \eta_{\mu\beta}\eta_{+\gamma} v^{\nu\alpha\beta\gamma} \int d^d x \partial_{x^\alpha} \left\{ \frac{1}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \right\} \frac{1}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}-1}} A_\nu^D(x) f^{A'B'C'} f^{DB'C'} \quad (2.59)$$

$$= \frac{-ig^2(-1)^2\Gamma^2\left(\frac{d}{2}-1\right) C_A \delta^{A'D}}{4^2\pi^d} \eta_{\mu\beta}\eta_{+\gamma} v^{\nu\alpha\beta\gamma} \int d^d x \frac{2\left(\frac{d}{2}-1\right) x_\alpha}{(-x^2 + i\epsilon)^{d-1}} A_\nu^D(x+zn) \quad (2.60)$$

$$= \frac{-ig^2(-1)^2\Gamma^2\left(\frac{d}{2}-1\right) C_A \delta^{A'D}}{4^2\pi^d} \eta_{\mu\beta}\eta_{+\gamma} v^{\nu\alpha\beta\gamma} \int d^d x \frac{2\left(\frac{d}{2}-1\right) x_\alpha}{(-x^2 + i\epsilon)^{d-1}} (1 + x^\lambda \partial_\lambda + \dots) A_\nu^D(zn) \quad (2.61)$$

now one can evaluate the integral.

I however like to introduce integration variables, even though it seems there is no point to this, however it makes the Gamma functions vanish naturally. Start from first term after evaluating the derivative, but now introduce integration variables instead of just combining powers:

$$-ig^2 f^{A'B'C'} \int d^d x \partial_{x^\alpha} \Delta_{\mu\beta}(zn-x) \delta^{B'E} \Delta_{+\gamma}(zn-x) \delta^{FC'} A_\nu^D(x) v^{\nu\alpha\beta\gamma} f^{DEF} \quad (2.62)$$

$$= \frac{-ig^2(-1)^2 \Gamma\left(\frac{d}{2}-1\right) \Gamma(d-1) C_A \delta^{A'D}}{4^2 \pi^d \Gamma\left(\frac{d}{2}\right)} \eta_{\mu\beta} \eta_{+\gamma} v^{\nu\alpha\beta\gamma} \int d^d x \int du \frac{2\left(\frac{d}{2}-1\right) u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} x_\alpha}{(-x^2+i\epsilon)^{d-1}} A_\nu^D(x+zn) \quad (2.63)$$

$$= \frac{g^2 \Gamma\left(\frac{d}{2}-1\right) C_A \delta^{A'D}}{4^2 \pi^{\frac{d}{2}} \Gamma\left(\frac{d}{2}\right)} \eta_{\mu\beta} \eta_{+\gamma} \eta_\alpha^\lambda v^{\nu\alpha\beta\gamma} \frac{2\left(\frac{d}{2}-1\right)}{2\epsilon} \partial_\lambda A_\nu^D(zn) \quad (2.64)$$

$$= \frac{\alpha_s C_A \delta^{A'D}}{4\pi} \frac{1}{\epsilon} \eta_{\mu\beta} \eta_{+\gamma} \eta_\alpha^\lambda v^{\nu\alpha\beta\gamma} \partial_\lambda A_\nu^D(zn) \quad (2.65)$$

Integral over integration variables gives a factor of  $\frac{1}{2}$  here. Factors can be combined using  $\Gamma$  function identities:  $\Gamma(x+1) = \Gamma(x)x$  metric algebra:

$$\eta_{\mu\beta} \eta_{+\gamma} \eta_\alpha^\lambda v^{\nu\alpha\beta\gamma} = \eta_{\mu\beta} \eta_{+\gamma} \eta_\alpha^\lambda (2\eta^{\nu\beta} \eta^{\alpha\gamma} - \eta^{\nu\alpha} \eta^{\beta\gamma} - 2\eta^{\nu\gamma} \eta^{\alpha\beta}) \quad (2.66)$$

$$= 2\eta_\mu^\nu \eta_+^\lambda - \eta g^{\nu\lambda} \eta_{\mu+} - 2\eta^{\nu+} \eta_\mu^\lambda \quad (2.67)$$

$$\eta_{\mu\beta} \eta_{+\gamma} \eta_\alpha^\lambda v^{\nu\alpha\beta\gamma} \partial_\lambda A_\nu^D(zn) = 2(\partial_+ A_\mu^D - \partial_\mu A_+^D) - \eta_{\mu+} \partial_\nu A^{\nu D} \quad (2.68)$$

then we have the second term: from the very beginning we can just say it is the same with exchange of indices:  $(\mu+)(B'C') \leftrightarrow (+\mu)(C'B')$  since the propagators commute. The color indices switch once, this gives a total minus. Thus my result is

$$\text{2nd term} = -\text{1st term} ((\mu+) \leftrightarrow (+\mu)) \quad (2.69)$$

By looking at the lorentz structure the total result for this diagrams is

$$\frac{\alpha_s C_A}{\pi\epsilon} (\partial_+ A_\mu^{A'} - \partial_\mu A_+^{A'}) (zn) \quad (2.70)$$

Due to light-cone gauge the second term drops, we have

$$-\int_{\infty n}^{zn} dx F_{\mu+}(x) = -\int_{\infty n}^{zn} dx (\partial_\mu A_+ - \partial_+ A_\mu)(x) \quad (2.71)$$

$$= \int_{\infty n}^{zn} dx \partial_+ A_\mu(x) \quad (2.72)$$

$$= \int_{\infty}^z d\tau \partial_\tau A_\mu(\tau n) \quad (2.73)$$

$$= A_\mu(zn) \quad (2.74)$$

$$\partial_+ A_\mu = -F_{\mu+} \quad (2.75)$$

$$\frac{\alpha_s C_A}{\pi\epsilon} (\partial_+ A_\mu^{A'} - \partial_\mu A_+^{A'}) (zn) = -\frac{\alpha_s C_A}{\pi\epsilon} F_{\mu+}(zn) \quad (2.76)$$

Computatin of other term: again:

$$F_{\mu\rho} = \partial_\mu B_\rho - \partial_\rho B_\mu \quad (2.77)$$



So i can take one term for computation. In the end you can replace indices and take into account minus sign. Simplifies computation. So for computation i use these terms :

$$\mathbf{T}^{A'} \partial_\mu B_\rho^{A'}(zn) \text{ig} \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{bk}^{B'} \left( - \text{ig} \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.78)$$

$$+ \mathbf{T}_{ab}^{A'} \partial_\mu B_\rho^{A'}(zn) \text{ig} \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{bk}^{B'} \left( - \text{ig} \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.79)$$

The red color matrix is not included in the computation, but i put it there to understand what index goes where. as stated above I compute these terms now, and add interchanged  $\rho \leftrightarrow \mu$  with minus sign. (and also replace  $\rho$  by  $+$ !) for the appearing derivatives i use my formulae

Note that since the properties of the vector  $n$  and the transversality of  $\mu$  plus the role of the index  $\rho$  which is interchanged by  $\mu$  or not, and in the end is  $+$  all implies that in any case  $n_\mu = 0$  (0 if transverse, null if  $n^+$ )

$$-i^2 g^2 \int d^d x \int_{-\infty}^z d\sigma \partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn-x) \delta^{A'E} \Delta_{+\gamma}(\sigma n-x) \delta^{B'F} v_{DEF}^{\nu\alpha\beta\gamma} A_\nu^D(x) T_{bk}^{B'} \quad (2.80)$$

$$= \frac{(-1)^3 i^2 g^2 \Gamma^2\left(\frac{d}{2}-1\right)}{4^2 \pi^d} \int d^d x \int_{-\infty}^z d\sigma \partial_{z^\mu} \partial_{x^\alpha} \frac{1}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \eta_{\rho\beta} \delta^{A'E} \frac{1}{(-(\sigma n-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \eta_{+\gamma} \delta^{B'F} v_{DEF}^{\nu\alpha\beta\gamma} A_\nu^D(x) T_{bk}^{B'} \quad (2.81)$$

$$= \frac{(-1)^3 i^2 g^2 \Gamma^2\left(\frac{d}{2}-1\right)}{4^2 \pi^d} \int d^d x \int_{-\infty}^z d\sigma \left( \frac{4\left(\frac{d}{2}-1\right) \frac{d}{2} (zn-x)_\mu (x-zn)_\alpha}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}+1}} + \frac{2\left(\frac{d}{2}-1\right) \eta_{\mu\alpha}}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}}} \right) \quad (2.82)$$

$$\frac{1}{(-(\sigma n-x)^2 + i\epsilon)^{\frac{d}{2}-1}} v_{DEF}^{\nu\alpha\beta\gamma} \eta_{\rho\beta} \delta^{A'E} \eta_{+\gamma} \delta^{B'F} A_\nu^D(x) T_{bk}^{B'} \quad (2.83)$$

$$= \frac{(-1)^3 i^2 g^2 \Gamma^2\left(\frac{d}{2}-1\right)}{4^2 \pi^d} \int d^d x \int_{-\infty}^z d\sigma \int_0^1 du \left\{ \frac{4u^{\frac{d}{2}} \bar{u}^{\frac{d}{2}-2} \Gamma(d) \left(\frac{d}{2}-1\right) \frac{d}{2} (zn-x)_\mu (x-zn)_\alpha}{\Gamma\left(\frac{d}{2}-1\right) \Gamma\left(\frac{d}{2}+1\right) (-u(zn-x)^2 - \bar{u}(\sigma n-x)^2 + i\epsilon)^d} \right. \quad (2.84)$$

$$\left. + \frac{2u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d-1) \left(\frac{d}{2}-1\right) \eta_{\mu\alpha}}{\Gamma\left(\frac{d}{2}\right) \Gamma\left(\frac{d}{2}-1\right) (-u(zn-x)^2 - \bar{u}(\sigma n-x)^2 + i\epsilon)^{d-1}} \right\} \quad (2.85)$$

$$v_{DEF}^{\nu\alpha\beta\gamma} \eta_{\rho\beta} \delta^{A'E} \eta_{+\gamma} \delta^{B'F} A_\nu^D(x) T_{bk}^{B'} \quad (2.86)$$

$$= \frac{(-1)^{d+3} i^2 g^2}{4^2 \pi^d} \int d^d x \int_{-\infty}^z d\sigma \int_0^1 du \left\{ \frac{4u^{\frac{d}{2}} \bar{u}^{\frac{d}{2}-2} \Gamma(d) (zn-x)_\mu (x-zn)_\alpha}{\left(\frac{d}{2}+1\right) (u(zn-x)^2 + \bar{u}(\sigma n-x)^2 - i\epsilon)^d} \right. \quad (2.87)$$

$$\left. - \frac{2u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d-1) \eta_{\mu\alpha}}{(u(zn-x)^2 + \bar{u}(\sigma n-x)^2 - i\epsilon)^{d-1}} \right\} v_{DEF}^{\nu\alpha\beta\gamma} \eta_{\rho\beta} \delta^{A'E} \eta_{+\gamma} \delta^{B'F} A_\nu^D(x) T_{bk}^{B'} \quad (2.88)$$

$$= \frac{(-1)^{d+4} i^2 g^2}{4^2 \pi^d} \int d^d x \int_{-\infty}^z d\sigma \int_0^1 du \left\{ \frac{4u^{\frac{d}{2}} \bar{u}^{\frac{d}{2}-2} \Gamma(d) x_\mu (x-zn)_\alpha}{\left(\frac{d}{2}+1\right) (x^2 - i\epsilon)^d} \right. \quad (2.89)$$

$$\left. + \frac{2u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d-1) \eta_{\mu\alpha}}{(x^2 - i\epsilon)^{d-1}} \right\} v_{DEF}^{\nu\alpha\beta\gamma} \eta_{\rho\beta} \delta^{A'E} \eta_{+\gamma} \delta^{B'F} A_\nu^D(x + (uz + \bar{u}\sigma)n) T_{bk}^{B'} \quad (2.90)$$

Shift is  $x \rightarrow x + uz n + \bar{u}\sigma n$ . since  $n$  is light-like, there is no additional term in the denominator. To compute

color, note that

$$T_{ab}^A = -i f^{Aab} \quad (2.91)$$

Thus have the color factor:

$$f^{DEF} \delta^{A'E} \delta^{B'F} T_{bk}^{B'} = -i f^{DA'B'} f^{B'bk} \quad (2.92)$$

$$= i f^{A'DB'} f^{bkB'} \quad (2.93)$$

$$= i C_A \delta^{A'b} \quad (2.94)$$

At this point i think one has to do numerator analysis. If I am looking for  $\frac{1}{\epsilon}$  pole i need following integrals

$$\frac{x^n}{x^{d+n-2}} \quad (2.95)$$

this is why, for the second term i need two derivatives. For the first term I need 2-4 derivatives and the rest must be  $x$  from brackets

second term:

$$\int d^d x \frac{x^\tau x^\omega}{(x^2)^{d-1}} \partial_\tau \partial_\omega A_\nu^D((uz + \bar{u}\sigma)n) = \frac{i\pi^{\frac{d}{2}}}{2\epsilon\Gamma(d-1)} \partial^2 A_\nu^D((uz + \bar{u}\sigma)n) \quad (2.96)$$

first term: Terms are this one and also a term for third derivative of  $A$ :  $\partial^3 A$ . I ignore it here but it has to be examined what happens to it.

$$\int d^d x \frac{x_\mu x_\alpha x^\tau x^\omega}{(x^2)^d} \partial_\tau \partial_\omega A_\nu^D((uz + \bar{u}\sigma)n) = \frac{i\pi^{\frac{d}{2}}}{4\epsilon\Gamma(d)} \partial^\tau \partial^\omega A_\nu^D((uz + \bar{u}\sigma)n) (\eta_{\mu\alpha} \eta_{\tau\omega} + \eta_{\mu\tau} \eta_{\alpha\omega} + \eta_{\mu\omega} \eta_{\alpha\tau}) \quad (2.97)$$

$$= \frac{i\pi^{\frac{d}{2}}}{4\epsilon\Gamma(d)} (\eta_{\mu\alpha} \partial^2 A_\nu^D + 2\partial_\mu \partial_\alpha A_\nu^D) ((uz + \bar{u}\sigma)n) \quad (2.98)$$

## C Self correction

By the following idea: Wilson gauge line adds belongs to the structure. The field and the line form a gauge invariant unit, and transform a one. Hence the color indices belong to the whole object.

$$F_{\mu\nu}^A T^A \rightarrow (F_{\mu\nu}(x) [x, y])^A T^A \quad (2.99)$$

For self education i compute the following diagram

$$\overline{F_{\mu+}(zn) \left[ ig \int_{-\infty}^z d\sigma B_+(\sigma n) \right]} = \overline{F_{\mu+}^A(zn) T_{ac}^A \left[ ig \int_{-\infty}^z d\sigma B_+^B T_{cb}^B(\sigma n) \right]} \quad (2.100)$$

$$= ig \int_{-\infty}^z d\sigma \partial_\alpha \Delta_{\beta+}((z-\sigma)n) \left( \eta_\mu^\alpha \eta_+^\beta - \eta_+^\alpha \eta_\mu^\beta \right) \delta^{AB} T_{ac}^A T_{cb}^B \quad (2.101)$$

$$\delta^{AB} T_{ac}^A T_{cb}^B = -f^{Aac} f^{Bcb} \delta^{AB} \quad (2.102)$$

$$= f^{aAc} f^{bAc} \quad (2.103)$$

$$= \delta^{ab} C_2 G \quad (2.104)$$

This result is nonsense, it is not proportional to a color matrix, but to identity ( i think )  
other possibility

$$\overline{F_{\mu+}(zn) \left[ ig \int_{-\infty}^z d\sigma B_+(\sigma n) \right]} = \overline{F_{\mu+}^A(zn) \left[ ig \int_{-\infty}^z d\sigma B_+^B T_{AC}^B(\sigma n) \right]} \quad (2.105)$$

$$-i f^{BAC} \delta^{AB} = f^{AAC} \quad (2.106)$$

$$= 0 \quad (2.107)$$

if computation is correct in this way, it demonstrates that this diagram vanishes. Does this computation make sense? Field tensor can be written as sum over coefficients for color matrices. Wrong thought previously was that the color index is not on the gluon field alone, but on the gluon field coupled to the wilson line! There is the open color index. And that index survives.

Again compute the terms: now with this color combination

$$\partial_\mu B_\rho^{A'}(zn) ig \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{A'C}^{B'} \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.108)$$

$$+ \partial_\mu B_\rho^{A'}(zn) ig \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{A'C}^{B'} \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.109)$$

computation yields: (it is urgent to develop systematic procedure to compute these objects, otherwise it is time waste)

$$g^2 \int_{-\infty}^z d\sigma \int d^d x \left( \partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn-x) \Delta_{+\gamma}(\sigma n-x) \delta^{A'E} \delta^{B'F} + \partial_{z^\mu} \Delta_{\rho\gamma}(zn-x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n-x) \delta^{A'F} \delta^{B'E} \right) A_\nu^D(x) v_{DEF}^{\nu\alpha\beta\gamma} T_{A'C}^{B'} \quad (2.110)$$

computation of color yields:

$$f^{DEF} T_{A'C}^{B'} (\delta^{A'E} \delta^{B'F} + \delta^{A'F} \delta^{B'E}) = -i (f^{DEF} f^{B'A'C} \delta^{A'E} \delta^{B'F} + f^{DEF} f^{B'A'C} \delta^{A'F} \delta^{B'E}) \quad (2.111)$$

$$= -i (-f^{DEF} f^{CEF} + f^{DEF} f^{CEF}) \quad (2.112)$$

$$= i C_A \delta^{CD} (+1 - 1) \quad (2.113)$$

This means, color produces relative minus sign between diagrams.

$$ig^2 C_A \int_{-\infty}^z d\sigma \int d^d x (\partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn-x) \Delta_{+\gamma}(\sigma n-x) - \partial_{z^\mu} \Delta_{\rho\gamma}(zn-x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n-x)) A_\nu^C(x) v^{\nu\alpha\beta\gamma} \quad (2.114)$$

Now compute both propagator combinations (derivatives acting on them) Ignore common factors of  $i\Gamma\pi$  etc.

$$\partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn-x) \Delta_{+\gamma}(\sigma n-x) = \left( -\frac{4\left(\frac{d}{2}-1\right) \frac{d}{2} (zn-x)_\mu (zn-x)_\alpha}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}+1}} + \frac{2\left(\frac{d}{2}-1\right) \eta_{\mu\alpha}}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}}} \right) \frac{1}{(-(\sigma n-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \quad (2.115)$$

$$= \int_0^1 du \left( -\frac{4u^{\frac{d}{2}} \bar{u}^{\frac{d}{2}-2} \Gamma(d) \left(\frac{d}{2}-1\right) \frac{d}{2} (zn-x)_\mu (zn-x)_\alpha}{\Gamma\left(\frac{d}{2}+1\right) \Gamma\left(\frac{d}{2}-1\right) (-u(zn-x)^2 - \bar{u}(\sigma n-x) + i\epsilon)^d} \right. \quad (2.116)$$

$$\left. + \frac{2u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d-1) \left(\frac{d}{2}-1\right) \eta_{\mu\alpha}}{\Gamma\left(\frac{d}{2}\right) \Gamma\left(\frac{d}{2}-1\right) (-u(zn-x)^2 - \bar{u}(\sigma n-x) + i\epsilon)^{d-1}} \right) \quad (2.117)$$

$$= \int_0^1 du \frac{u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2}}{\Gamma^2\left(\frac{d}{2}-1\right)} \left( \frac{4u\Gamma(d) x_\mu (\bar{u}(\sigma-z)n-x)_\alpha}{(-x^2 + i\epsilon)^d} \right. \quad (2.118)$$

$$\left. + \frac{2\Gamma(d-1) \eta_{\mu\alpha}}{(-x^2 + i\epsilon)^{d-1}} \right) \quad (2.119)$$

note that we use  $n_\mu = 0$  since its a transverse component The other term yields:

$$\partial_{z^\mu} \Delta_{\rho\gamma}(zn-x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n-x) = \left( 2\left(\frac{d}{2}-1\right) \right)^2 \frac{(x-zn)_\mu}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{(x-\sigma n)_\alpha}{(-(\sigma n-x)^2 + i\epsilon)^{\frac{d}{2}}} \quad (2.120)$$

$$= 4 \int_0^1 du \frac{u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-1} \Gamma(d) x_\mu (x+u(z-\sigma)n)_\alpha}{\Gamma^2\left(\frac{d}{2}-1\right) (-x^2 + i\epsilon)^d} \quad (2.121)$$

To mention the shifts:

$$x \rightarrow x + un + \bar{u}\sigma n \quad (2.122)$$

$$x - zn \rightarrow x - \bar{u}(z-\sigma)n \quad (2.123)$$

$$x - \sigma n \rightarrow x + u(z-\sigma)n \quad (2.124)$$

$$AND \quad (2.125)$$

$$\sigma \rightarrow \sigma + z \quad (2.126)$$

Now carefully identify common factors and put them in front. Also respect different metric index and the relative minus coming from color. combination yields: (skip common previous factors); denominator Gammas drop with propagators.)

$$\int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \left( \frac{4\Gamma(d)x_\mu \{u(\bar{u}\sigma n - x)_\alpha \eta_{\rho\beta} \eta_{+\gamma} - \bar{u}(x - u\sigma n)_\alpha \eta_{\rho\gamma} \eta_{+\beta}\}}{(-x^2 + i\epsilon)^d} + \frac{2\Gamma(d-1) \eta_{\mu\alpha} \eta_{\rho\beta} \eta_{+\gamma}}{(-x^2 + i\epsilon)^{d-1}} \right) A_\nu(x + zn + \bar{u}\sigma n) \quad (2.127)$$

Now think of all terms that account to  $\frac{1}{\epsilon}$  pole. Note sign flip of second term due to  $-x^2$ ;  $d = 4$  when using integration formula.

for second term i need double derivative:

$$\int d^d x \frac{\Gamma(d-1) x^\lambda x^\tau}{(-x^2 + i\epsilon)^{d-1}} = \frac{-i\pi^{\frac{d}{2}}}{2(2 - \frac{d}{2})} \eta^{\lambda\tau} \quad (2.128)$$

For the first term I can take the bracket  $x$  or  $n$  vector, thus have two or three derivatives. In any case its four  $x$  numerator formula:

$$\int d^d x \frac{\Gamma(d) x^\mu x^\nu x^\rho x^\sigma}{(-x^2 + i\epsilon)^d} = \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})} (\eta^{\mu\nu} \eta^{\rho\sigma} + \eta^{\mu\rho} \eta^{\nu\sigma} + \eta^{\mu\sigma} \eta^{\nu\rho}) \quad (2.129)$$

Merging to total expression: (after  $d$ -dim integral), omitting factors( $\pi$ )

$$\frac{i}{\epsilon} \left( - \{u \eta_{\rho\beta} \eta_{+\gamma} + \bar{u} \eta_{\rho\gamma} \eta_{+\beta}\} (\eta_{\mu\alpha} \eta^{\tau\sigma} + \eta_\mu^\tau \eta_\alpha^\sigma + \eta_\mu^\sigma \eta_\alpha^\tau) \partial_\sigma \partial_\tau \right. \quad (2.130)$$

$$+ u \bar{u} \sigma n_\alpha \{-\eta_{\rho\beta} \eta_{+\gamma} + \eta_{\rho\gamma} \eta_{+\beta}\} (\eta_\mu^\omega \eta^{\tau\sigma} + \eta_\mu^\tau \eta^\omega{}^\sigma + \eta_\mu^\sigma \eta^\omega{}^\tau) \partial_\omega \partial_\tau \partial_\sigma \quad (2.131)$$

$$\left. - \eta_{\mu\alpha} \eta_{\rho\beta} \eta_{+\gamma} \partial^2 \right) A_\nu((z + \bar{u}\sigma)n) \quad (2.132)$$

1st line is first term taking 2 derivatives 2nd line is first term taking 3 derivatives 3rd line is second term  
Now one should contract metric (possibly?) and compute  $\sigma$  integral. To perform a to a previous one similar computation I expect some  $\partial_+ A$  term. However other terms should drop(?)

Maybe it is a good idea to contract Lorentzindices at this stage( $v^{\nu\alpha\beta\gamma}$ ). We have this expression:

$$v^{\nu\alpha\beta\gamma} = 2\eta^{\nu\beta} - \eta^{\nu\alpha} \eta^{\beta\gamma} - 2\eta^{\nu\gamma} \eta^{\alpha\beta} \quad (2.133)$$

$$v^{\nu\alpha\beta\gamma} \eta_{\mu\alpha} \eta_{\rho\beta} \eta_{+\gamma} = 0 \quad (2.134)$$

$$v^{\nu\alpha\beta\gamma} \eta_{\rho\beta} \eta_{+\gamma} = 2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\nu+} \eta^{\alpha\rho}) \quad (2.135)$$

$$v^{\nu\alpha\beta\gamma} \eta_{\rho\gamma} \eta_{+\beta} = -2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\nu+} \eta^{\alpha\rho}) \quad (2.136)$$

This is another relative minus sign!

In my notation  $\mu$  and  $\rho$  are to be interchanged, and  $\rho$  is being replaced by  $+$  in the end. Thus, in any combination:

$$\eta^{\mu+} = \eta^{\rho+} = 0 \quad (2.137)$$

$$g_{++} = n^2 = 0 \quad (2.138)$$

light like and  $\mu$  transverse. thus these terms can be immediately dropped!!!

This implies the third term is dropped completely, and the first gains common factor in metric but

relative minus sign for terms in numerator.

$$\frac{4\Gamma(d)2(\eta_\rho^\nu\eta^{\alpha+} - \eta^{\nu+}\eta_\rho^\alpha)x_\mu\{u(\bar{u}\sigma n - x)_\alpha + \bar{u}(x - u\sigma n)_\alpha\}}{(-x^2 + i\epsilon)^d}A_\nu(x + zn + \bar{u}\sigma n) \quad (2.139)$$

$$= \frac{4\Gamma(d)2(\eta_\rho^\nu\eta^{\alpha+} - \eta^{\nu+}\eta_\rho^\alpha)x_\mu(1 - 2u)x_\alpha}{(-x^2 + i\epsilon)^d}A_\nu(x + zn + \bar{u}\sigma n) \quad (2.140)$$

This result is good, no divergent term. Also i have only one contributing term further: double derivative on  $A$ .

By integration obtain factors:  $\frac{i\pi^{\frac{d}{2}}}{4(2-\frac{d}{2})\Gamma(d)}$  Further the Lorentz Structure is

$$(\eta_\rho^\nu\eta^{\alpha+} - \eta^{\nu+}\eta_\rho^\alpha)(\eta_{\mu\alpha}\eta^{\tau\sigma} + \eta_\mu^\tau\eta_\alpha^\sigma + \eta_\mu^\sigma\eta_\alpha^\tau)\partial_\tau\partial_\sigma A_\nu - (\mu \leftrightarrow \rho)[\rho = +] \quad (2.141)$$

$$= 2(\partial_\mu\partial_+A_+ - \partial_+\partial_\mu - \partial_+\partial_+A_\mu + \partial_\mu\partial_+A_+) \quad (2.142)$$

$$= 2\partial_+(\partial_\mu A_+ - \partial_+A_\mu) \quad (2.143)$$

due to light-cone gauge. Recollecting all prefactors chronologically:

$$ig^2C_A \frac{(-1)^2\Gamma^2(\frac{d}{2}-1)}{4^2\pi^d} 4 \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d)\Gamma^{-2}\left(\frac{d}{2}-1\right) 2(1-2u) \frac{i\pi^{\frac{d}{2}}}{4(2-\frac{d}{2})\Gamma(d)} \quad (2.144)$$

$$= \frac{-\alpha_s C_A}{2\pi} du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (1-2u) \quad (2.145)$$

Thus

$$\frac{-\alpha_s C_A}{\pi} \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (1-2u) \int_{-\infty}^0 d\sigma \partial_+ (\partial_\mu A_+ - \partial_+ A_\mu) (zn + \bar{u}\sigma n) \quad (2.146)$$

check for errors, second go through

$$\partial_\mu B_\rho^{A'}(zn) ig \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{A'C}^{B'} \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.147)$$

$$+ \partial_\mu B_\rho^{A'}(zn) ig \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{A'C}^{B'} \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.148)$$

computation yields: (it is urgent to develop systematic procedure to compute these objects, otherwise it is time waste)

$$g^2 \int_{-\infty}^z d\sigma \int d^d x \left( \partial_{z\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn-x) \Delta_{+\gamma}(\sigma n-x) \delta^{A'E} \delta^{B'F} + \partial_{z\mu} \Delta_{\rho\gamma}(zn-x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n-x) \delta^{A'F} \delta^{B'E} \right) A_\nu^D(x) v_{DEF}^{\nu\alpha\beta\gamma} T_{A'C}^{B'} \quad (2.149)$$

computation of color yields:

$$f^{DEF} T_{A'C}^{B'} (\delta^{A'E} \delta^{B'F} + \delta^{A'F} \delta^{B'E}) = -i(f^{DEF} f^{B'A'C} \delta^{A'E} \delta^{B'F} + f^{DEF} f^{B'A'C} \delta^{A'F} \delta^{B'E}) \quad (2.150)$$

$$= -i(-f^{DEF} f^{CEF} + f^{DEF} f^{CEF}) \quad (2.151)$$

$$= iC_A \delta^{CD} (+1 - 1) \quad (2.152)$$

This means, color produces relative minus sign between diagrams.

$$ig^2 C_A \int_{-\infty}^z d\sigma \int d^d x (\partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn - x) \Delta_{+\gamma}(\sigma n - x) - \partial_{z^\mu} \Delta_{\rho\gamma}(zn - x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n - x)) A_\nu^C(x) v^{\nu\alpha\beta\gamma} \quad (2.153)$$

Now compute both propagator combinations (derivatives acting on them) Ignore common factors of  $i\Gamma\pi$  etc.

$$\partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn - x) \Delta_{+\gamma}(\sigma n - x) = \left( -\frac{4 \left(\frac{d}{2} - 1\right) \frac{d}{2} (zn - x)_\mu (zn - x)_\alpha}{(-(zn - x)^2 + i\epsilon)^{\frac{d}{2}+1}} - \frac{2 \left(\frac{d}{2} - 1\right) \eta_{\mu\alpha}}{(-(zn - x)^2 + i\epsilon)^{\frac{d}{2}}} \right) \frac{1}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}-1}} \quad (2.154)$$

$$= \int_0^1 du \left( -\frac{4u^{\frac{d}{2}} \bar{u}^{\frac{d}{2}-2} \Gamma(d) \left(\frac{d}{2} - 1\right) \frac{d}{2} (zn - x)_\mu (zn - x)_\alpha}{\Gamma\left(\frac{d}{2} + 1\right) \Gamma\left(\frac{d}{2} - 1\right) (-u(zn - x)^2 - \bar{u}(\sigma n - x) + i\epsilon)^d} \right. \quad (2.155)$$

$$\left. - \frac{2u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d-1) \left(\frac{d}{2} - 1\right) \eta_{\mu\alpha}}{\Gamma\left(\frac{d}{2}\right) \Gamma\left(\frac{d}{2} - 1\right) (-u(zn - x)^2 - \bar{u}(\sigma n - x) + i\epsilon)^{d-1}} \right) \quad (2.156)$$

$$= \int_0^1 du \frac{u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2}}{\Gamma^2\left(\frac{d}{2} - 1\right)} \left( \frac{4u\Gamma(d) x_\mu (\bar{u}\sigma n - x)_\alpha}{(-x^2 + i\epsilon)^d} \right. \quad (2.157)$$

$$\left. - \frac{2\Gamma(d-1) \eta_{\mu\alpha}}{(-x^2 + i\epsilon)^{d-1}} \right) \quad (2.158)$$

note that we use  $n_\mu = 0$  since its a transverse component The other term yields:

$$\partial_{z^\mu} \Delta_{\rho\gamma}(zn - x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n - x) = \left( 2 \left(\frac{d}{2} - 1\right) \right)^2 \frac{-(x - zn)_\mu}{(-(zn - x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{(x - \sigma n)_\alpha}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \quad (2.159)$$

$$= -4 \int_0^1 du \frac{u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-1} \Gamma(d)}{\Gamma^2\left(\frac{d}{2} - 1\right)} \frac{x_\mu (x - u\sigma n)_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.160)$$

To mention the shifts:

$$x \rightarrow x + un + \bar{u}\sigma n \quad (2.161)$$

$$x - zn \rightarrow x - \bar{u}(z - \sigma)n \quad (2.162)$$

$$x - \sigma n \rightarrow x + u(z - \sigma)n \quad (2.163)$$

$$AND \quad (2.164)$$

$$\sigma \rightarrow \sigma + z \quad (2.165)$$

Now carefully identify common factors and put them in front. Also respect different metric index and the relative minus coming from color. combination yields: (skip common previous factors); denominator Gammas drop with propagators.)

$$\int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \left( \frac{4\Gamma(d)x_\mu \{u(\bar{u}\sigma n - x)_\alpha \eta_{\rho\beta} \eta_{+\gamma} - \bar{u}(x - u\sigma n)_\alpha \eta_{\rho\gamma} \eta_{+\beta}\}}{(-x^2 + i\epsilon)^d} + \frac{2\Gamma(d-1) \eta_{\mu\alpha} \eta_{\rho\beta} \eta_{+\gamma}}{(-x^2 + i\epsilon)^{d-1}} \right) A_\nu(x + zn + \bar{u}\sigma n) \quad (2.166)$$

Now think of all terms that account to  $\frac{1}{\epsilon}$  pole. Note sign flip of second term due to  $-x^2$ ;  $d = 4$  when using integration formula.

for second term i need double derivative:

$$\int d^d x \frac{\Gamma(d-1) x^\lambda x^\tau}{(-x^2 + i\epsilon)^{d-1}} = \frac{-i\pi^{\frac{d}{2}}}{2(2 - \frac{d}{2})} \eta^{\lambda\tau} \quad (2.167)$$

For the first term I can take the bracket  $x$  or  $n$  vector, thus have two or three derivatives. In any case its four  $x$  numerator formula:

$$\int d^d x \frac{\Gamma(d) x^\mu x^\nu x^\rho x^\sigma}{(-x^2 + i\epsilon)^d} = \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})} (\eta^{\mu\nu} \eta^{\rho\sigma} + \eta^{\mu\rho} \eta^{\nu\sigma} + \eta^{\mu\sigma} \eta^{\nu\rho}) \quad (2.168)$$

Merging to total expression: (after  $d$ -dim integral), omitting factors( $\pi$ )

$$\frac{i}{\epsilon} \left( - \{u\eta_{\rho\beta} \eta_{+\gamma} + \bar{u}\eta_{\rho\gamma} \eta_{+\beta}\} (\eta_{\mu\alpha} \eta^{\tau\sigma} + \eta_\mu^\tau \eta_\alpha^\sigma + \eta_\mu^\sigma \eta_\alpha^\tau) \partial_\sigma \partial_\tau \right. \quad (2.169)$$

$$+ u\bar{u}\sigma n_\alpha \{-\eta_{\rho\beta} \eta_{+\gamma} + \eta_{\rho\gamma} \eta_{+\beta}\} (\eta_\mu^\omega \eta^{\tau\sigma} + \eta_\mu^\tau \eta^\omega{}^\sigma + \eta_\mu^\sigma \eta^\omega{}^\tau) \partial_\omega \partial_\tau \partial_\sigma \quad (2.170)$$

$$\left. - \eta_{\mu\alpha} \eta_{\rho\beta} \eta_{+\gamma} \partial^2 \right) A_\nu((z + \bar{u}\sigma)n) \quad (2.171)$$

1st line is first term taking 2 derivatives 2nd line is first term taking 3 derivatives 3rd line is second term Now one should contract metric (possibly?) and compute  $\sigma$  integral. To perform a to a previous one similar computation I expect some  $\partial_+ A$  term. However other terms should drop(?)

Maybe it is a good idea to contract Lorentzindices at this stage( $v^{\nu\alpha\beta\gamma}$ ) . We have this expression:

$$v^{\nu\alpha\beta\gamma} = 2\eta^{\nu\beta} \eta^{\alpha\gamma} - \eta^{\nu\alpha} \eta^{\beta\gamma} - 2\eta^{\nu\gamma} \eta^{\alpha\beta} \quad (2.172)$$

$$v^{\nu\alpha\beta\gamma} \eta_{\mu\alpha} \eta_{\rho\beta} \eta_{+\gamma} = 0 \quad (2.173)$$

$$v^{\nu\alpha\beta\gamma} \eta_{\rho\beta} \eta_{+\gamma} = 2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\nu+} \eta^{\alpha\rho}) \quad (2.174)$$

$$v^{\nu\alpha\beta\gamma} \eta_{\rho\gamma} \eta_{+\beta} = -2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\nu+} \eta^{\alpha\rho}) \quad (2.175)$$

This is another relative minus sign!

In my notation  $\mu$  and  $\rho$  are to be interchanged, and  $\rho$  is being replaced by  $+$  in the end. Thus, in any



combination:

$$\eta^{\mu+} = \eta^{\rho+} = 0 \quad (2.176)$$

$$g_{++} = n^2 = 0 \quad (2.177)$$

light like and  $\mu$  transverse. thus these terms can be immediately dropped!!!

This implies the third term is dropped completely, and the first gains common factor in metric but relative minus sign for terms in numerator.

$$\frac{4\Gamma(d)2(\eta_\rho^\nu \eta^{\alpha+} - \eta^{\nu+} \eta_\rho^\alpha) x_\mu \{u(\bar{u}\sigma n - x)_\alpha + \bar{u}(x - u\sigma n)_\alpha\}}{(-x^2 + i\epsilon)^d} A_\nu(x + zn + \bar{u}\sigma n) \quad (2.178)$$

$$= \frac{4\Gamma(d)2(\eta_\rho^\nu \eta^{\alpha+} - \eta^{\nu+} \eta_\rho^\alpha) x_\mu (1 - 2u) x_\alpha}{(-x^2 + i\epsilon)^d} A_\nu(x + zn + \bar{u}\sigma n) \quad (2.179)$$

This result is good, no divergent term. Also i have only one contributing term further: double derivative on  $A$ .

By integration obtain factors:  $\frac{i\pi^{\frac{d}{2}}}{4(2-\frac{d}{2})\Gamma(d)}$  Further the Lorentz Structure is

$$(\eta_\rho^\nu \eta^{\alpha+} - \eta^{\nu+} \eta_\rho^\alpha) (\eta_{\mu\alpha} \eta^{\tau\sigma} + \eta_\mu^\tau \eta_\alpha^\sigma + \eta_\mu^\sigma \eta_\alpha^\tau) \partial_\tau \partial_\sigma A_\nu - (\mu \leftrightarrow \rho) [\rho = +] \quad (2.180)$$

$$= 2(\partial_\mu \partial_+ A_+ - \partial_+ \partial_\mu - \partial_+ \partial_+ A_\mu + \partial_\mu \partial_+ A_+) \quad (2.181)$$

$$= 2\partial_+ (\partial_\mu A_+ - \partial_+ A_\mu) \quad (2.182)$$

due to light-cone gauge. Recollecting all prefactors chronologically:

$$ig^2 C_A \frac{(-1)^2 \Gamma^2(\frac{d}{2} - 1)}{4^2 \pi^d} 4 \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \Gamma(d) \Gamma^{-2} \left( \frac{d}{2} - 1 \right) 2(1 - 2u) \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} \quad (2.183)$$

$$= \frac{-\alpha_s C_A}{2\pi} du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (1 - 2u) \quad (2.184)$$

Thus

$$\frac{-\alpha_s C_A}{\pi} \int_0^1 du u (1 - 2u) \int_{-\infty}^0 d\sigma \partial_+ (\partial_\mu A_+ - \partial_+ A_\mu) (zn + \bar{u}\sigma n) \quad (2.185)$$

$$\int_0^1 du \int_{-\infty}^0 d\sigma u(1-2u) \partial_+ f(zn + \bar{u}\sigma n) \quad (2.186)$$

$$= \int \frac{d^d p}{(2\pi)^d} \int_0^1 du \int_{-\infty}^0 d\sigma u(1-2u) (-ip_+) e^{ip_+(z+\bar{u}\sigma)+\delta\sigma} f(p) \quad (2.187)$$

$$= \int \frac{d^d p}{(2\pi)^d} \int_0^1 du u(1-2u) \frac{-ip_+}{ip_+ \bar{u} + \delta} e^{ip_+ z} f(p) \quad (2.188)$$

$$= \int \frac{d^d p}{(2\pi)^d} \int_0^1 d\bar{u} \frac{1}{\bar{u} + \frac{\delta}{ip_+}} (1-3\bar{u}+2\bar{u}^2) e^{ip_+ z} f(p) \quad (2.189)$$

$$= \int \frac{d^d p}{(2\pi)^d} \left( \ln \left( \frac{ip_+}{\delta} \right) - 2 \right) e^{ip_+ z} f(p) \quad (2.190)$$

$$= \left( \ln \left( \frac{i\hat{p}_+}{\delta} \right) - 2 \right) f(zn) \quad (2.191)$$

$$\frac{-\alpha_s C_A}{\pi} \left( \ln \left( \frac{i\hat{p}_+}{\delta} \right) - 2 \right) F_{\mu+}(zn) \quad (2.192)$$

Then there are diagrams with inverted structure: Fields at  $-zn$  and wilson line is now from  $-\infty$  to  $-zn$ . To check how the result changes, carefully compare the computation.

$$\left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} \overline{B_\beta^E(x)} \overline{B_\gamma^F(x)} \right) v_{DEF}^{\nu\alpha\beta\gamma} ig \int_{-z}^{-\infty} d\sigma B_+^{B'}(\sigma n) T_{CA'}^{B'} \partial_\mu B_\rho^{A'}(-zn) \quad (2.193)$$

$$+ \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} \overline{B_\beta^E(x)} \overline{B_\gamma^F(x)} \right) v_{DEF}^{\nu\alpha\beta\gamma} ig \int_{-z}^{-\infty} d\sigma B_+^{B'}(\sigma n) T_{CA'}^{B'} \partial_\mu B_\rho^{A'}(-zn) \quad (2.194)$$

next step is identical;  $zn - x$  becomes  $zn + x$  in propagators. But other propagator does not change, only integration limits. Note the change of order of fields also implies other color structure. It will result in total minus sign, so i expect.

$$-ig^2 \int_{-zn}^{-\infty} d\sigma \int d^d x \left( \partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn+x) \Delta_{+\gamma}(\sigma n-x) \delta^{A'E} \delta^{B'F} + \partial_{z^\mu} \Delta_{\rho\gamma}(zn+x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n-x) \delta^{A'F} \delta^{B'E} \right) A_\nu^D(x) v_{DEF}^{\nu\alpha\beta\gamma} T_{CA'}^{B'} \quad (2.195)$$

For the color computation we have the same structure, but a total minus sign due to the interchange in  $T_{A'C}^{B'} \rightarrow T_{CA'}^{B'}$ .

The shifts in this case are

$$x \rightarrow x - uz n + \bar{u} \sigma n \quad (2.196)$$

$$AND \quad (2.197)$$

$$\sigma \rightarrow \sigma - z \quad (2.198)$$

$$Result \text{ in} \quad (2.199)$$

$$x \rightarrow x + \bar{u} \sigma n - z n \quad (2.200)$$

$$x + z n \rightarrow x + \bar{u} \sigma n \quad (2.201)$$

$$x - \sigma n \rightarrow x - u \sigma n \quad (2.202)$$

I go on and copy the expression eq. (2.158) Comment: the derivative on field is in the end evaluated for  $z = -zn$ . This is how i get this result ( so i rewrite  $-zn = z$ , compute the derivative and then reinsert)

$$\partial_{z^\mu} \partial_{x^\alpha} \Delta_{\rho\beta}(zn+x) \Delta_{+\gamma}(\sigma n-x) = \left( + \frac{4 \left(\frac{d}{2}-1\right) \frac{d}{2} (zn+x)_\mu (zn+x)_\alpha}{(- (zn+x)^2 + i\epsilon)^{\frac{d}{2}+1}} - \frac{2 \left(\frac{d}{2}-1\right) \eta_{\mu\alpha}}{(- (zn+x)^2 + i\epsilon)^{\frac{d}{2}}} \right) \frac{1}{(- (\sigma n-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \quad (2.203)$$

$$= \int_0^1 du \frac{u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2}}{\Gamma^2 \left(\frac{d}{2}-1\right)} \left( \frac{4u\Gamma(d) x_\mu (\bar{u}\sigma n+x)_\alpha}{(-x^2+i\epsilon)^d} \right. \quad (2.204)$$

$$\left. - \frac{2\Gamma(d-1) \eta_{\mu\alpha}}{(-x^2+i\epsilon)^{d-1}} \right) \quad (2.205)$$

this is obtained by thinking, but to be 100% sure one should really check it fully explicitly. But im sure it is correct. Other term (copyied from eq. (2.160)): (and here for the derivative same comment as above!)

$$\partial_{z^\mu} \Delta_{\rho\gamma}(zn+x) \partial_{x^\alpha} \Delta_{+\beta}(\sigma n-x) = +4 \int_0^1 du \frac{u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-1} \Gamma(d) x_\mu (x-u\sigma n)_\alpha}{\Gamma^2 \left(\frac{d}{2}-1\right) (-x^2+i\epsilon)^d} \quad (2.206)$$

Then I should contract the metric for all the terms. For double derivative on propagator i have

$$\eta_{\rho\beta} \eta_{\gamma+} v^{\nu\alpha\beta\gamma} = 2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\rho\alpha} \eta^{\nu+}) \quad (2.207)$$

where the second term in eq. (2.205) drops since it implies multiplication with  $\eta^{\alpha\mu}$  and thus gives zero. Other contraction is

$$\eta_{+\beta} \eta_{\gamma\rho} v^{\nu\alpha\beta\gamma} = -2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\rho\alpha} \eta^{\nu+}) \quad (2.208)$$

In other words it is the same calulation, obviously. Nothing else to expect. So, from color i have relative minus (and total minus ), from metric i have relative minus (relative here between first and second term) that is why nominator structure of added term is (no color factors and Gammas)

$$+8 \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{x_\mu x_\alpha}{(-x^2+i\epsilon)^d} (\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\rho\alpha} \eta^{\nu+}) \quad (2.209)$$

Field  $A(x \rightarrow x + \bar{u}\sigma n - zn)$  double expanded, then integration over  $x$ .

$$\int d^d x \frac{x_\mu x_\alpha x_\tau x_\sigma}{(-x^2 + i\epsilon)^d} \partial^\tau \partial^\sigma A(zn + \bar{u}n) = \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} (\text{"sym.metric"}) \quad (2.210)$$

contracting produced metric with existing and subtract interchanged term ( $\rho \leftrightarrow \mu$   $\rho = +$ ) (i also treat  $\sigma$  and  $\tau$  as identical indices since they can be interchanged in the derivative.)

$$2(\eta_{\mu\tau}\eta_{\sigma+}\eta_{\nu+} - \eta_{\mu\nu}\eta_{\tau+}\eta_{\sigma+}) \quad (2.211)$$

Combine all this:

$$\frac{-16i^2 C_A \delta^{CD} g^2}{4^2 4\epsilon\pi^2} \int_0^{-\infty} d\sigma \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (\partial_\mu \partial_+ A_+^D - \partial_+ \partial_+ A_\mu^D) \quad (2.212)$$

$$= \frac{4\alpha_s C_A}{\epsilon} \int_0^{-\infty} d\sigma \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (\partial_\mu \partial_+ A_+^C - \partial_+ \partial_+ A_\mu^C) (-zn + \bar{u}\sigma n) \quad (2.213)$$

$$\int_0^{-\infty} d\sigma \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (\partial_\mu \partial_+ A_+^C - \partial_+ \partial_+ A_\mu^C) (-zn + \bar{u}\sigma n) \quad (2.214)$$

$$= \int_0^{-\infty} d\sigma \int_0^1 du u \partial_+ f(y + \bar{u}\sigma n) \quad (2.215)$$

$$= \int \frac{d^d p}{(2\pi)^4} \int_0^{-\infty} d\sigma \int_0^1 du u (-ip_+) e^{i(y + \bar{u}\sigma n|p) + \delta\sigma} f(p) \quad (2.216)$$

$$= - \int \frac{d^d p}{(2\pi)^4} \int_0^1 du u \frac{-ip_+}{i\bar{u}p_+ + \delta} e^{i(y|p)} f(p) \quad (2.217)$$

$$= \int \frac{d^d p}{(2\pi)^4} \int_0^1 d\bar{u} \frac{1 - \bar{u}}{\bar{u} + \frac{\delta}{ip_+}} e^{i(y|p)} f(p) \quad (2.218)$$

$$= \int \frac{d^d p}{(2\pi)^4} \int_{\frac{\delta}{ip_+}}^{1 + \frac{\delta}{ip_+}} d\bar{u} \frac{1 - \bar{u} - 0}{\bar{u}} e^{i(y|p)} f(p) \quad (2.219)$$

$$= \int \frac{d^d p}{(2\pi)^4} \left( 1 + \ln \left( \frac{ip_+}{\delta} \right) \right) e^{i(y|p)} f(p) \quad (2.220)$$

$$= \left( 1 + \ln \left( \frac{ip_+}{\delta} \right) \right) f(y) \quad (2.221)$$

COMBINED RESULT FOR GLUON TWIST1:

## D gluons

A better working method is maybe to parallel write serious expressions and then drop from them from time to time parts into "prefactor" expression

Let me do parallel computing.

I take color red for right field, color blue for left field. Common factors i put out in front will be black,

no color. Thus here we have the following diagrams.

$$\left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} ig \int_{-z}^{-\infty} d\sigma B_+^{B'}(\sigma n) T_{CA'}^{B'} \partial_\mu B_\rho^{A'}(-zn) \quad (2.222)$$

$$+ \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} ig \int_{-z}^{-\infty} d\sigma B_+^{B'}(\sigma n) T_{CA'}^{B'} \partial_\mu B_\rho^{A'}(-zn) \quad (2.223)$$

$$\partial_\mu B_\rho^{A'}(zn) ig \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{A'C}^{B'} \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.224)$$

$$+ \partial_\mu B_\rho^{A'}(zn) ig \int_{-\infty}^z d\sigma B_+^{B'}(\sigma n) T_{A'C}^{B'} \left( -ig \int d^d x A_\nu^D(x) \partial_{x^\alpha} B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.225)$$

For the colors:

$$f^{DEF} T_{A'C}^{B'} (\delta^{A'E} \delta^{B'F} + \delta^{A'F} \delta^{B'E}) = -i(f^{DEF} f^{B'A'C} \delta^{A'E} \delta^{B'F} + f^{DEF} f^{B'A'C} \delta^{A'F} \delta^{B'E}) \quad (2.226)$$

$$= -i(-f^{DEF} f^{CEF} + f^{DEF} f^{CEF}) \quad (2.227)$$

$$= iC_A \delta^{CD} (+1 - 1) \quad (2.228)$$

+1 - 1 means, the first line has +, the second line has -1.

$$f^{DEF} T_{CA'}^{B'} (\delta^{A'E} \delta^{B'F} + \delta^{A'F} \delta^{B'E}) = -i(f^{DEF} f^{B'CA'} \delta^{A'E} \delta^{B'F} + f^{DEF} f^{B'CA'} \delta^{A'F} \delta^{B'E}) \quad (2.229)$$

$$= -i(f^{DEF} f^{FCE} + f^{DEF} f^{ECF}) \quad (2.230)$$

$$= -i f^{DEF} f^{FCE} (+1 - 1) \quad (2.231)$$

$$= -iC_A \delta^{CD} (+1 - 1) \quad (2.232)$$

$$= -iC_A \delta^{CD} (+1 - 1) \quad (2.233)$$

so from color the two diagrams differ by a total sign First common factors:

$$+i^3 C_A g^2 \int d^d v v^{\nu\alpha\beta\gamma} \delta^{CD} \quad (2.234)$$

$$\int_{-\infty}^z \partial_{zn^\mu} \partial_{x^\alpha} \Delta(zn - x)_{\rho\beta} \Delta(x - \sigma n)_{\gamma+} - \partial_{zn^\mu} \Delta(zn - x)_{\rho\gamma} \partial_{x^\alpha} \Delta(x - \sigma n)_{\beta+} \quad (2.235)$$

$$- \int_{-z}^{-\infty} \partial_{zn^\mu} \partial_{x^\alpha} \Delta(zn + x)_{\rho\beta} \Delta(x - \sigma n)_{\gamma+} - \partial_{zn^\mu} \Delta(zn + x)_{\rho\gamma} \partial_{x^\alpha} \Delta(x - \sigma n)_{\beta+} \quad (2.236)$$

Then we have the to compute as usual, derivatives on propagators. Since i know that the metric computation will kill the additional term from double derivative on one propagator (additional term  $\tilde{\eta}_{\mu\alpha}$  I IGNORE IT HERE.)

$$+i^3 C_A g^2 \int d^d \delta^{CD} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^d} \quad (2.237)$$

$$\int_{-\infty}^z d\sigma v^{\nu\alpha\beta\gamma} \quad (2.238)$$

$$\frac{4(\frac{d}{2}-1)\frac{d}{2}(zn-x)_\mu(x-zn)_\alpha}{(-(x-zn)^2+i\epsilon)^{\frac{d}{2}+1}} \frac{1}{(-(x-\sigma n)^2+i\epsilon)^{\frac{d}{2}-1}} \eta_{\rho\beta}\eta_{\gamma+} \quad (2.239)$$

$$- \frac{-2(\frac{d}{2}-1)(zn-x)_\mu - 2(\frac{d}{2}-1)(x-\sigma n)_\alpha}{(-(x-zn)^2+i\epsilon)^{\frac{d}{2}}} \frac{1}{(-(x-\sigma n)^2+i\epsilon)^{\frac{d}{2}}} \eta_{\rho\gamma}\eta_{\beta+} \quad (2.240)$$

$$- \int_{-z}^{-\infty} d\sigma v^{\nu\alpha\beta\gamma} \quad (2.241)$$

$$\frac{4(\frac{d}{2}-1)\frac{d}{2}(-zn-x)_\mu(x+zn)_\alpha}{(-(x+zn)^2+i\epsilon)^{\frac{d}{2}+1}} \frac{1}{(-(x-\sigma n)^2+i\epsilon)^{\frac{d}{2}-1}} \eta_{\rho\beta}\eta_{\gamma+} \quad (2.242)$$

$$- \frac{-2(\frac{d}{2}-1)(-zn-x)_\mu - 2(\frac{d}{2}-1)(x-\sigma n)_\alpha}{(-(x+zn)^2+i\epsilon)^{\frac{d}{2}}} \frac{1}{(-(x-\sigma n)^2+i\epsilon)^{\frac{d}{2}}} \eta_{\rho\gamma}\eta_{\beta+} \quad (2.243)$$

shifts:

$$x \rightarrow x + uz n + \bar{u}\sigma n \quad (2.244)$$

$$\sigma \rightarrow \sigma + z \quad (2.245)$$

$$HENCE \quad (2.246)$$

$$x \rightarrow x + zn + \bar{u}\sigma n \quad (2.247)$$

$$x - zn \rightarrow x + \bar{u}\sigma n \quad (2.248)$$

$$x - \sigma n \rightarrow x - u\sigma n \quad (2.249)$$

$$x \rightarrow x - uz n + \bar{u}\sigma n \quad (2.250)$$

$$\sigma \rightarrow \sigma - z \quad (2.251)$$

$$HENCE \quad (2.252)$$

$$x \rightarrow x - zn + \bar{u}\sigma n \quad (2.253)$$

$$x + zn \rightarrow x + \bar{u}\sigma n \quad (2.254)$$

$$x - \sigma n \rightarrow x - u\sigma n \quad (2.255)$$

Lorentz index computation is the same for both, red and blue. it is

$$v^{\nu\alpha\beta\gamma} = 2\eta^{\nu\beta}\eta^{\alpha\gamma} - \eta^{\nu\alpha}\eta^{\beta\gamma} - 2\eta^{\nu\gamma}\eta^{\alpha\beta} \quad (2.256)$$

$$v^{\nu\alpha\beta\gamma}\eta_{\mu\alpha}\eta_{\rho\beta}\eta_{+\gamma} = 0 \text{ (that is why there is no additional term from the double derivative on prop.)} \quad (2.257)$$

$$v^{\nu\alpha\beta\gamma}\eta_{\rho\beta}\eta_{+\gamma} = 2(\eta^{\nu\rho}\eta^{\alpha+} - \eta^{\nu+}\eta^{\alpha\rho}) \quad (2.258)$$

$$v^{\nu\alpha\beta\gamma}\eta_{\rho\gamma}\eta_{+\beta} = -2(\eta^{\nu\rho}\eta^{\alpha+} - \eta^{\nu+}\eta^{\alpha\rho}) \quad (2.259)$$

and results in relative minus sign between the respective lines plus a common factor:

$$+ i^3 C_A g^2 \int d^d \delta^{CD} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^d} 2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\nu+} \eta^{\alpha\rho}) \quad (2.260)$$

The minus sign from this computation annihilates the relative minus from color (the minus between the diagrams! this has nothing to do with the minus between red and blue!)

$$\int_{-\infty}^0 \int_0^1 d \frac{\Gamma(d)}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (-4x_\mu) \frac{u(x + \bar{u}\sigma n)_\alpha + \bar{u}(x - u\sigma n)_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.261)$$

$$= -4 \int_{-\infty}^0 \int_0^1 d \frac{\Gamma(d)}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{x_\mu x_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.262)$$

$$- \int_0^{-\infty} \int_0^1 d \frac{\Gamma(d)}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} (-4x_\mu) \frac{u(x + \bar{u}\sigma n)_\alpha + \bar{u}(x - u\sigma n)_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.263)$$

$$= +4 \int_0^{-\infty} \int_0^1 d \frac{\Gamma(d)}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{x_\mu x_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.264)$$

$$= -4 \int_{-\infty}^0 \int_0^1 d \frac{\Gamma(d)}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{x_\mu x_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.265)$$

at this point it looks like both computations are identical, but there is still the free field  $A_\nu(x)$ , which after shift is

$$A(x + zn + \bar{u}\sigma n) \quad A(x - zn + \bar{u}\sigma n) \quad (2.266)$$

This has to be kept in mind. Further the integration can be done in for both terms identically. I write  $A(x + yn + \bar{u}\sigma n)$ , and  $y = \pm z$ , blue or red case. It follows the following computation:

$$+ i^3 C_A g^2 \delta^{CD} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^d} 2(\eta^{\nu\rho} \eta^{\alpha+} - \eta^{\nu+} \eta^{\alpha\rho}) (-4) \int_{-\infty}^0 \int_0^1 d \frac{\Gamma(d)}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \quad (2.267)$$

$$\int d^d \frac{x_\mu x_\alpha x_\tau x_\sigma}{(-x^2 + i\epsilon)^d} \partial^\tau \partial^\sigma A(yn + \bar{u}\sigma n) = \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} (\eta_{\mu\alpha} \eta_{\tau\sigma} + \eta_{\mu\tau} \eta_{\alpha\sigma} + \eta_{\mu\sigma} \eta_{\alpha\tau}) A_\nu(yn + \bar{u}\sigma n) \quad (2.268)$$

Together metric and fields give

$$(\eta_{\mu\alpha} \eta_{\tau\sigma} + \eta_{\mu\tau} \eta_{\alpha\sigma} + \eta_{\mu\sigma} \eta_{\alpha\tau}) A_\nu(yn + \bar{u}\sigma n) - (\mu \leftrightarrow \rho) [\rho = +] \quad (2.269)$$

$$= 2\partial_+ (\partial_\mu A_+ - \partial_+ A_\mu) (yn + \bar{u}\sigma n) \quad (2.270)$$

Together with prefactors and so on

$$\frac{(-4)2^2 i^4 C_A g^2}{4^3 \pi^{\frac{d}{2}} \epsilon} \int_{-\infty}^0 d\sigma \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial_+ (\partial_\mu A_+^C - \partial_+ A_\mu^C) (yn + \bar{u}\sigma n) \quad (2.271)$$

now integration:

$$\int_0^{-\infty} d\sigma \int_0^1 du u \partial_+ f(yn + \bar{u}\sigma n) \quad (2.272)$$

$$= \int \frac{d^d p}{(2\pi)^4} \int_0^{-\infty} d\sigma \int_0^1 du u (-ip_+) e^{i(y + \bar{u}\sigma n|p) + \delta\sigma} f(p) \quad (2.273)$$

$$= - \int \frac{d^d p}{(2\pi)^4} \int_0^1 du u \frac{-ip_+}{i\bar{u}p_+ + \delta} e^{i(y|p)} f(p) \quad (2.274)$$

$$= \int \frac{d^d p}{(2\pi)^4} \int_0^1 d\bar{u} \frac{1 - \bar{u}}{\bar{u} + \frac{\delta}{ip_+}} e^{i(y|p)} f(p) \quad (2.275)$$

$$= \int \frac{d^d p}{(2\pi)^4} \int_{\frac{\delta}{ip_+}}^{1 + \frac{\delta}{ip_+}} d\bar{u} \frac{1 - \bar{u} - 0}{\bar{u}} e^{i(y|p)} f(p) \quad (2.276)$$

$$= \int \frac{d^d p}{(2\pi)^4} \left( -1 + \ln \left( \frac{ip_+}{\delta} \right) \right) e^{i(y|p)} f(p) \quad (2.277)$$

note that i have in my previous computation a different integration order, this minus sign. From the two cases  $y = \pm zn$  there are two results

$$\int_{-\infty}^0 d\sigma \int_0^1 du u \partial_+ f(\pm zn + \bar{u}\sigma n) = - \left( -1 + \ln \left( \frac{\pm i\hat{p}_+}{\delta} \right) \right) f(\pm zn) \quad (2.278)$$

then the final result is

$$\frac{g^2 C_A}{4\pi^2 \epsilon} \left( -1 + \ln \left( \frac{\pm i\hat{p}_+}{\delta} \right) \right) F_{\mu+}^C(\pm zn) \quad (2.279)$$

Other diagram.

$$gf^{A'B'C'} B_\mu^{B'}(zn) B_+^{C'}(zn) 1 \left( -ig \int d^d x A_\nu^D(x) \partial_\alpha B_\beta^E(x) B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.280)$$

$$\rightarrow gf^{A'B'C'} B_\mu^{B'}(zn) B_+^{C'}(zn) 1 \left( -ig \int d^d x A_\nu^D(x) \partial_\alpha B_\beta^E B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.281)$$

$$+ gf^{A'B'C'} B_\mu^{B'}(zn) B_+^{C'}(zn) 1 \left( -ig \int d^d x A_\nu^D(x) \partial_\alpha B_\beta^E B_\gamma^F(x) \right) v_{DEF}^{\nu\alpha\beta\gamma} \quad (2.282)$$

here i compute both terms/diagrams, but again split common factor. color computation:

$$f^{A'B'C'} f^{DEF} (\delta^{B'E} \delta^{C'F} + \delta^{B'F} \delta^{C'E}) \quad (2.283)$$

$$= f^{A'EF} f^{DEF} (+1 + (-1)) \quad (2.284)$$

$$= C_A \delta^{A'D} (+1 + (-1)) \quad (2.285)$$

$$-ig^2 \int d^d x A_\nu(x) C_A \delta^{A'D} v^{\nu\alpha\beta\gamma} \quad (2.286)$$

$$\partial_{x^\alpha} \Delta_{\mu\beta}(zn - x) \Delta_{\gamma+}(zn - x) - \Delta_{\mu\gamma}(zn - x) \partial_{x^\alpha} \Delta_{\beta+}(zn - x) \quad (2.287)$$



$$\frac{-ig^2\Gamma^2\left(\frac{d}{2}-1\right)}{4^2\pi^d}\int d^d x A_\nu^{A'}(x)C_A v^{\nu\alpha\beta\gamma} \quad (2.288)$$

$$\frac{-2\left(\frac{d}{2}-1\right)(x-zn)_\alpha}{(- (x-zn)^2 + i\epsilon)^{\frac{d}{2}}}\frac{1}{(- (x-zn)^2 + i\epsilon)^{\frac{d}{2}}-1}(\eta_{\gamma+}\eta_{\mu\beta}-\eta_{\beta+}\eta_{\mu\gamma}) \quad (2.289)$$

$$= \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{\Gamma(d-1)}{\Gamma\left(\frac{d}{2}-1\right)\Gamma\left(\frac{d}{2}\right)} \frac{-2\left(\frac{d}{2}-1\right)(x-zn)_\alpha}{(- (x-zn)^2 + i\epsilon)^{d-1}}(\eta_{\gamma+}\eta_{\mu\beta}-\eta_{\beta+}\eta_{\mu\gamma}) \quad (2.290)$$

$$= \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{\Gamma(d-1)}{\Gamma^2\left(\frac{d}{2}-1\right)} \frac{-2(x-zn)_\alpha}{(- (x-zn)^2 + i\epsilon)^{d-1}}(\eta_{\gamma+}\eta_{\mu\beta}-\eta_{\beta+}\eta_{\mu\gamma}) \quad (2.291)$$

The shift is  $x \rightarrow x + zn$ . The metric can be contracted:

$$(\eta_{\gamma+}\eta_{\mu\beta}-\eta_{\beta+}\eta_{\mu\gamma})v^{\nu\alpha\beta\gamma} = 4(\eta_\mu^\nu\eta^{\alpha+}-\eta_\mu^\alpha\eta_{\nu+}) \quad (2.292)$$

$$\frac{-ig^2C_A\Gamma(d-1)}{4^2\pi^d}\int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} 4(\eta_\mu^\nu\eta^{\alpha+}-\eta_\mu^\alpha\eta_{\nu+}) \quad (2.293)$$

$$\int d^d x \frac{-2(x-zn)_\alpha x_\tau}{(- (x)^2 + i\epsilon)^{d-1}} \partial^\tau A_\nu^{A'}(zn) \quad (2.294)$$

$$= \frac{+2i\pi^{\frac{d}{2}}}{2\epsilon} \frac{1}{\Gamma(d-1)} \partial_\alpha A_\nu^{A'}(zn) \quad (2.295)$$

Have in total

$$\frac{g^2C_A}{4\pi^2\epsilon}(\partial_+A_\mu-\partial_\mu A_+)^{A'}(zn) \quad (2.296)$$

$$= -\frac{g^2C_A}{4\pi^2\epsilon}F_{\mu+}^{A'}(zn) \quad (2.297)$$

Looks ok but not sure about factor. So this has to be checked again. But it does not contribute to the pole with regulator  $\delta$ . CHECKED IT, CORRECTED IT AND I THINK FACTOR IS OK NOW.

For other diagram, meaning other side of wilson line there is NO difference in color and also nothing else that makes me think there would be a difference. So it is the same result (of course at other space-time point.)

Next diagram:

$$gf^{A'B'C'}B_\mu^{B'}(zn)B_+^{C'}(zn)\int_{-\infty}^z d\sigma igB_+(\sigma n) \quad (2.298)$$

To the outgoing field has to be classical, so I have to choose which one it should be. However Once needs to




Table 1: diagrams for gluon TMD operator on twist 1

understand that already here the suspicion that the diagram is null can be made. Two contractions:

$$gf^{A'B'C'} \overline{B_\mu^{B'}(zn)A_+^{C'}(zn)} \int_{-\infty}^z d\sigma igB_+^D(\sigma n)T_{A'C}^D \quad (2.299)$$

$$+ gf^{A'B'C'} A_\mu^{B'}(zn) \overline{B_+^{C'}(zn)} \int_{-\infty}^z d\sigma igB_+^D(\sigma n)T_{A'C}^D \quad (2.300)$$

color is of course a relative minus. But the second diagram is null since  $n^2$  is 0. First diagram should be null in light cone gauge, and because  $g_{\mu+}$  is 0. So anyway it is 0.

Now the results: The diagrams are in table 1:

The results are

$$A = \frac{g^2 C_A}{4\pi^2 \epsilon} \left( -1 + \ln \left( \frac{i\hat{p}_+}{\delta} \right) \right) F_{\mu+}^C(zn) \quad (2.301)$$

$$B = -\frac{g^2 C_A}{4\pi^2 \epsilon} F_{\mu+}^{A'}(zn) \quad (2.302)$$

$$C = 0 \quad (2.303)$$

1-loop diagrams for gluon pdf: twist2:

operators with this twist should look have the form

$$\int d\sigma F_{\mu+}[zn, \sigma n] F_{\nu+}[\sigma, -\infty n] \quad (2.304)$$

Probably it is ok to ignore the integration which should be kept until the end anyway? Like I have learned before the color of the whole object is connected, and just one index is open here. so I need to know what to do for:

no disturbance disturbance in first path disturbance in second path

$$F_{\mu+}^A F_{\nu+}^B T_{AC}^B \quad (2.305)$$

$$ig \int_{\sigma}^z d\tau F_{\mu+}^A B_+^B(\tau n) T_{AC}^B F_{\nu+}^D T_{CE}^D \quad (2.306)$$

$$ig \int_{-\infty}^{\sigma} d\tau F_{\mu+}^A F_{\nu+}^B T_{AC}^B B_+^D(\tau n) T_{CE}^D \quad (2.307)$$

Note: I will compute for antisymmetric part of tensor: (i know it is fully as)  $partial_\mu A_M$ , since  $M$  is capital  $\mu$ . at some convenient point in computation i will subtract interchanged term and replace  $M$  by  $+$ . Same for  $\nu \leftrightarrow N$ . with this:

Result table:

A:

$$F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.308)$$

$$\rightarrow \quad (2.309)$$

$$\overbrace{F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma}} \quad (2.310)$$

$$+ \overbrace{F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma}} \quad (2.311)$$

Color factor

$$T_{AC}^B f^{A'B'C'} T_{CE}^D (\delta^{BC'} \delta^{AB'} - \delta^{BB'} \delta^{AC'}) = +i C_A \delta^{A'C} (1-1) T_{CE}^D \quad (2.312)$$

The remaining term is

$$-i^2 g^2 F_{\nu+}^D \int d^d x A_l^{A'}(x) \int_\sigma^z d\tau \left( \partial_\mu \partial_\alpha \Delta_{M\beta}(zn-x) \Delta_{+\gamma}(\tau n-x) - \partial_\mu \Delta_{M\gamma}(zn-x) \partial_\alpha \Delta_{+\beta}(\tau n-x) \right) v^{\iota\alpha\beta\gamma} \quad (2.313)$$

$$\int_\sigma^z d\tau \int d^d x A_l^{A'}(x) \left( \partial_\mu \partial_\alpha \Delta_{M\beta}(zn-x) \Delta_{+\gamma}(\tau n-x) - \partial_\mu \Delta_{M\gamma}(zn-x) \partial_\alpha \Delta_{+\beta}(\tau n-x) \right) v^{\iota\alpha\beta\gamma} \quad (2.314)$$

$$= \int_\sigma^z d\tau \int d^d x A_l^{A'}(x) \frac{\Gamma^2(\frac{d}{2}-1)}{4^2 \pi^d} \left( \frac{4(\frac{d}{2}-1) \frac{d}{2} (x-zn)_\alpha (zn-x)_\mu}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{1}{(-(\tau n-x)^2 + i\epsilon)^{\frac{d}{2}-1}} \eta_{M\beta} \eta_{+\gamma} \right) \quad (2.315)$$

$$- \frac{2(\frac{d}{2}-1)(zn-x)_\mu - 2(\frac{d}{2}-1)(x-\tau n)_\alpha}{(-(zn-x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{1}{(-(\tau n-x)^2 + i\epsilon)^{\frac{d}{2}}} \eta_{M\gamma} \eta_{+\beta} \right) v^{\iota\alpha\beta\gamma} \quad (2.316)$$

$$= \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \int d^d x A_l^{A'}(x) \frac{4\Gamma(d)}{4^2 \pi^d} \left( \frac{u(x+\bar{u}\tau n)_\alpha - x_\mu}{(-x^2 + i\epsilon)^d} \eta_{M\beta} \eta_{+\gamma} - \frac{-\bar{u}x_\mu(x-u\tau n)_\alpha}{(-x^2 + i\epsilon)^d} \eta_{M\gamma} \eta_{+\beta} \right) v^{\iota\alpha\beta\gamma} \quad (2.317)$$

$$= \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \int d^d x A_l^{A'}(x+zn+\bar{u}\tau n) \frac{4\Gamma(d)}{4^2 \pi^d} \left( \frac{u(x+\bar{u}\tau n)_\alpha - x_\mu}{(-x^2 + i\epsilon)^d} + \frac{-\bar{u}x_\mu(x-u\tau n)_\alpha}{(-x^2 + i\epsilon)^d} \right) 2(\eta^{\iota M} \eta^{\alpha+} - \eta^{\iota+} \eta^{M\alpha}) \quad (2.318)$$

$$= \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \int d^d x A_l^{A'}(x+zn+\bar{u}\tau n) \frac{4\Gamma(d)}{4^2 \pi^d} \left( \frac{-x_\mu x_\alpha}{(-x^2 + i\epsilon)^d} \right) 2(\eta^{\iota M} \eta^{\alpha+} - \eta^{\iota+} \eta^{M\alpha}) \quad (2.319)$$

$$= \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \int d^d x \frac{\partial^\zeta \partial^\eta}{2} A_l^{A'}(zn+\bar{u}\tau n) \frac{4\Gamma(d)}{4^2 \pi^d} \left( \frac{-x_\mu x_\alpha x_\eta x_\zeta}{(-x^2 + i\epsilon)^d} \right) 2(\eta^{\iota M} \eta^{\alpha+} - \eta^{\iota+} \eta^{M\alpha}) \quad (2.320)$$

$$= \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{\partial^\zeta \partial^\eta}{2} A_l^{A'}(zn+\bar{u}\tau n) \frac{4\Gamma(d)}{4^2 \pi^d} \frac{i\pi^{\frac{d}{2}}}{4(2-\frac{d}{2})\Gamma(d)} (\eta_{\mu\alpha} \eta_{\eta\zeta} + \eta_{\mu\eta} \eta_{\alpha\zeta} + \eta_{\mu\zeta} \eta_{\alpha\eta}) 2(\eta^{\iota M} \eta^{\alpha+} - \eta^{\iota+} \eta^{M\alpha}) \quad (2.321)$$

$$= \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \frac{4\Gamma(d)}{4^2 \pi^d} \frac{i\pi^{\frac{d}{2}}}{4(2-\frac{d}{2})\Gamma(d)} \frac{2}{2} \cdot 2\partial_+ (\partial_\mu A_+ - \partial_+ A_\mu)(zn+\bar{u}\tau n) \quad (2.322)$$

$$= \int_{\sigma-z}^0 d\tau \frac{i}{8\pi^{\frac{d}{2}} \epsilon} \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial_+ \left( \partial_\mu A_+^{A'} - \partial_+ A_\mu^{A'} \right) (zn+\bar{u}\tau n) \quad (2.323)$$

This is an important intermediate result. With it i can compute the diagrams quicker. However i made the mistake to not include the  $\tau$  integral, thus i need to check for the shifts!

ok so we have for  $A$ :

$$-i^2 g^2 F_{\nu+}^D \frac{i}{4\pi^{\frac{d}{2}}\epsilon} \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial_+ \left( \partial_\mu A_+^{A'} - \partial_+ A_\mu^{A'} \right) (zn + \bar{u}\tau n) \quad (2.324)$$

$$= \frac{-g^2 C_A}{4\pi^{\frac{d}{2}}\epsilon} F_{\nu+}^D (\sigma n) T_{CE}^D \int_{\sigma-z}^0 d\tau \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial_+ F_{\mu+}^C (zn + \bar{u}\tau n) \quad (2.325)$$

for the integrals i have a formula in appendix, so i just insert and have:

$$\frac{+g^2 C_A}{4\pi^{\frac{d}{2}}\epsilon} F_{\nu+}^D (\sigma n) T_{CE}^D \int_0^1 du \frac{u}{\bar{u}} \left( F_{\mu+}^C (zn + \bar{u}(\sigma - z)n) - F_{\mu+}^C (zn) \right) \quad (2.326)$$

Result for diagram  $A$ !

Diagram  $A'$ :

$$F_{\mu M}^A F_{\nu N}^D T_{AC}^D i g B_+^B T_{CE}^B \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.327)$$

$$\rightarrow g^2 F_{\nu N}^D \int_{-\infty}^{\sigma} d\tau \int d^d x A_l^{A'}(x) \left( \partial_\mu \partial_\alpha \Delta_{M\beta}(zn - x) \Delta_{+\gamma}(\tau n - x) - \partial_\mu \Delta_{M\gamma}(zn - x) \partial_\alpha \Delta_{\beta+}(\tau n - x) \right) v^{\iota\alpha\beta\gamma} i \frac{C_A}{2} T_{A'E}^D \quad (2.328)$$

$$= g^2 F_{\nu N}^D \int_{-\infty}^{\sigma-z} d\tau \frac{i}{4\pi^{\frac{d}{2}}\epsilon} \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial_+ \left( \partial_\mu A_+^{A'} - \partial_+ A_\mu^{A'} \right) (zn + \bar{u}\tau n) i \frac{C_A}{2} T_{A'E}^D \quad (2.329)$$

$$= g^2 F_{\nu N}^D \left( \int_{-\infty}^0 + \int_0^{\sigma-z} \right) d\tau \frac{i}{4\pi^{\frac{d}{2}}\epsilon} \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial_+ \left( \partial_\mu A_+^{A'} - \partial_+ A_\mu^{A'} \right) (zn + \bar{u}\tau n) i \frac{C_A}{2} T_{A'E}^D \quad (2.330)$$

$$= \frac{-g^2 C_A}{8\pi^{\frac{d}{2}}\epsilon} \int_{-\infty}^0 d\tau \int_0^1 du u \partial_+ F_{\mu+}^{A'}(zn + \bar{u}\tau n) F_{\nu+}^D T_{A'E}^D \quad (2.331)$$

$$+ \frac{-g^2 C_A}{8\pi^{\frac{d}{2}}\epsilon} \int_0^{\sigma-z} d\tau \int_0^1 du u \partial_+ F_{\mu+}^{A'}(zn + \bar{u}\tau n) F_{\nu+}^D T_{A'E}^D \quad (2.332)$$

$$= \frac{-g^2 C_A}{8\pi^{\frac{d}{2}}\epsilon} \left( 1 - \ln \left( \frac{ip_+}{\delta} \right) \right) F_{\mu+}^{A'}(zn) F_{\nu+}^D (\sigma n) T_{A'E}^D \quad (2.333)$$

$$+ \frac{-g^2 C_A}{8\pi^{\frac{d}{2}}\epsilon} \int_0^1 du \frac{u}{\bar{u}} \left( F_{\mu+}^{A'}(\sigma n + u(z - \sigma)n) - F_{\mu+}^{A'}(zn) \right) F_{\nu+}^D T_{A'E}^D \quad (2.334)$$

Diagram  $B$ :

$$\int_{\sigma}^z F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.335)$$

$$\rightarrow \quad (2.336)$$

$$\int_{\sigma}^z (-1)^2 (i)^3 \frac{C_A}{2} g^2 T_{AE}^{A'} F_{\mu M}^A \int d^d x A_l^{A'}(x) \left( \partial_\mu \partial_\alpha \Delta_{N\beta}(\sigma n - x) \Delta_{+\gamma}(\tau n - x) - \partial_\mu \Delta_{N\gamma}(\sigma n - x) \partial_\alpha \Delta_{+\beta}(\tau n - x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.337)$$

$$= (-1)^2 (i)^3 \frac{C_A}{2} g^2 T_{AE}^{A'} F_{\mu M}^A \frac{i}{4\pi^{\frac{d}{2}} \epsilon} \int_0^{z-\sigma} \int_0^1 du u \partial_+ F_{\nu+}^{A'}(\sigma n + \bar{u}\tau n) \quad (2.338)$$

$$= \frac{g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} F_{\mu M}^A \int_0^1 du \frac{u}{u} \left( F_{\nu+}^{A'}(\sigma n + \bar{u}(z - \sigma)n) - F_{\nu+}^{A'}(\sigma n) \right) T_{AE}^{A'} \quad (2.339)$$

Diagram  $B'$

$$\int_{-\infty}^{\sigma} F_{\mu M}^A F_{\nu N}^D T_{AC}^D i g B_+^B T_{CE}^B \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.340)$$

$$\rightarrow \quad (2.341)$$

$$\int_{-\infty}^{\sigma} \frac{-i^3 g^2 C_A}{2} T_{AE}^{A'} F_{\mu M}^A \int d^d x A_l^{A'}(x) \left( \partial_\mu \partial_\alpha \Delta_{N\beta}(\sigma n - x) \Delta_{+\gamma}(\tau n - x) - \partial_\mu \Delta_{N\gamma}(\sigma n - x) \partial_\alpha \Delta_{+\beta}(\tau n - x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.342)$$

$$= \int_{-\infty}^0 \frac{-i^3 g^2 C_A}{2} T_{AE}^{A'} F_{\mu M}^A \frac{i}{4\pi^{\frac{d}{2}} \epsilon} \int_0^1 du u \partial_+ F_{\nu+}^{A'}(\sigma n + \bar{u}\tau n) \quad (2.343)$$

$$= T_{AE}^{A'} F_{\mu M}^A \frac{-g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \int_{-\infty}^0 \int_0^1 du u \partial_+ F_{\nu+}^{A'}(\sigma n + \bar{u}\tau n) \quad (2.344)$$

$$= T_{AE}^{A'} F_{\mu M}^A \frac{-g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \left( 1 - \ln \left( \frac{i p_+}{\delta} \right) \right) F_{\nu+}^{A'}(\sigma n) \quad (2.345)$$

Diagram  $A1$

$$g f^{ABC} B_\mu^B B_+^C(zn) F_{\nu+}^D T_{AE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.346)$$

$$\rightarrow g f^{ABC} \overbrace{B_\mu^B B_+^C(zn) F_{\nu+}^D T_{AE}^D} \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.347)$$

$$+ g f^{ABC} \overbrace{B_\mu^B B_+^C(zn) F_{\nu+}^D T_{AE}^D} \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.348)$$

$$= -i g^2 C_A F_{\nu+}^D T_{AE}^D \int d^d x A_l^A(x) \left( \partial_\alpha \Delta_{\mu\beta}(zn - x) \Delta_{\gamma+}(zn - x) - \Delta_{\mu\gamma}(zn - x) \partial_\alpha \Delta_{+\beta}(zn - x) \right) v^{\iota\alpha\beta\gamma} \quad (2.349)$$

as this steps returns:

$$\int d^d x A_\ell^A(x) \left( \partial_\alpha \Delta_{\mu\beta}(zn-x) \Delta_{\gamma+}(zn-x) - \Delta_{\mu\gamma}(zn-x) \partial_\alpha \Delta_{+\beta}(zn-x) \right) v^{\iota\alpha\beta\gamma} \quad (2.350)$$

$$= \frac{\Gamma^2\left(\frac{d}{2}-1\right)}{\left(4\pi^{\frac{d}{2}}\right)^2} \int d^d x A_\ell^A(x) \left( \frac{-2\left(\frac{d}{2}-1\right)(x-zn)_\alpha}{(-(zn-x)^2+i\epsilon)^{\frac{d}{2}}} \frac{\eta_{\mu\beta}\eta_{+\gamma}}{(-(zn-x)^2+i\epsilon)^{\frac{d}{2}-1}} - \frac{\eta_{\mu\gamma}\eta_{+\beta}}{(-(zn-x)^2+i\epsilon)^{\frac{d}{2}-1}} \frac{-2\left(\frac{d}{2}-1\right)(x-zn)_\alpha}{(-(zn-x)^2+i\epsilon)^{\frac{d}{2}}} \right) v^{\iota\alpha\beta\gamma} \quad (2.351)$$

$$= \frac{\Gamma(d-1)}{\left(4\pi^{\frac{d}{2}}\right)^2} \int d^d x \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} A_\ell^A(x+zn) \left( \frac{-2x_\alpha}{(-x^2+i\epsilon)^{d-1}} (\eta_{\mu\beta}\eta_{+\gamma} - \eta_{\mu\gamma}\eta_{+\beta}) \right) v^{\iota\alpha\beta\gamma} \quad (2.352)$$

$$= \frac{\Gamma(d-1)}{\left(4\pi^{\frac{d}{2}}\right)^2} \int d^d x \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial^\lambda A_\ell^A(zn) \frac{-2x_\alpha x_\lambda}{(-x^2+i\epsilon)^{d-1}} 4 (\eta_\mu^\iota \eta^{+\alpha} - \eta_\mu^\alpha \eta^{+\iota}) \quad (2.353)$$

$$= \frac{\Gamma(d-1)}{\left(4\pi^{\frac{d}{2}}\right)^2} \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-2} \partial^\lambda A_\ell^A(zn) \frac{2i\pi^{\frac{d}{2}}}{2(2-\frac{d}{2})\Gamma(d-1)} \frac{\eta_{\alpha\lambda}}{4} (\eta_\mu^\iota \eta^{+\alpha} - \eta_\mu^\alpha \eta^{+\iota}) \quad (2.354)$$

$$= \frac{i}{4\pi^{\frac{d}{2}}\epsilon} \frac{1}{2} (\partial_+ A_\mu^A - \partial_\mu A_+^A)(zn) \quad (2.355)$$

$$= \frac{-i}{8\pi^{\frac{d}{2}}\epsilon} F_{\mu+}^A(zn) \quad (2.356)$$

also a nice intermediate result, which i can use for other diagrams too. together for diagram A1

$$-ig^2 C_A F_{\nu+}^D T_{AE}^D \frac{-i}{8\pi^{\frac{d}{2}}\epsilon} F_{\mu+}^A(zn) \quad (2.357)$$

$$= \frac{-g^2 C_A}{8\pi^{\frac{d}{2}}\epsilon} F_{\mu+}^A(zn) F_{\nu+}^D T_{AE}^D \quad (2.358)$$

diagram B2: it is just the same with field at different position. but same factor! (it has to be)

Diagram C:

$$\partial_\mu B_M^A \partial_\nu B_N^D T_{AE}^D \left( -ig \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \left( -ig \int d^d y A_\theta^{D'}(y) \partial_{y^\omega} B_\kappa^{E'}(y) B_\lambda^{F'}(y) \right) v_{D'E'F'}^{\theta\omega\kappa\lambda} \quad (2.359)$$

$$\rightarrow 2 \quad (2.360)$$

$$\partial_\mu B_M^A \partial_\nu B_N^D T_{AE}^D \left( -ig \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \left( -ig \int d^d y A_\theta^{D'}(y) \partial_{y^\omega} B_\kappa^{E'}(y) B_\lambda^{F'}(y) \right) v_{D'E'F'}^{\theta\omega\kappa\lambda} \quad (2.361)$$

$$+ \partial_\mu B_M^A \partial_\nu B_N^D T_{AE}^D \left( -ig \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \left( -ig \int d^d y A_\theta^{D'}(y) \partial_{y^\omega} B_\kappa^{E'}(y) B_\lambda^{F'}(y) \right) v_{D'E'F'}^{\theta\omega\kappa\lambda} \quad (2.362)$$

$$+ \partial_\mu B_M^A \partial_\nu B_N^D T_{AE}^D \left( -ig \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \left( -ig \int d^d y A_\theta^{D'}(y) \partial_{y^\omega} B_\kappa^{E'}(y) B_\lambda^{F'}(y) \right) v_{D'E'F'}^{\theta\omega\kappa\lambda} \quad (2.363)$$

$$+ \partial_\mu B_M^A \partial_\nu B_N^D T_{AE}^D \left( -ig \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \left( -ig \int d^d y A_\theta^{D'}(y) \partial_{y^\omega} B_\kappa^{E'}(y) B_\lambda^{F'}(y) \right) v_{D'E'F'}^{\theta\omega\kappa\lambda} \quad (2.364)$$

$$= -i^2 \frac{C_A}{2} g^2 \int d^d x \int d^d y A_\ell^{A'}(x) A_\theta^{D'}(y) v^{\iota\alpha\beta\gamma} v^{\theta\omega\kappa\lambda} \quad (2.365)$$

$$\left( \partial_\mu \partial_\omega \Delta_{M\kappa}(zn - y) \partial_\nu \partial_\alpha \Delta_{N\kappa}(\sigma n - x) \Delta_{\gamma\lambda}(x - y) \right) \quad (2.366)$$

$$- \partial_\mu \Delta_{\lambda M}(zn - y) \partial_\nu \partial_\alpha \Delta_{N\beta}(\sigma n - x) \partial_\omega \Delta_{\gamma\kappa}(x - y) \quad (2.367)$$

$$- \partial_\mu \partial_\omega \Delta_{M\kappa}(zn - y) \partial_\nu \Delta_{N\gamma}(\sigma n - x) \partial_\alpha \Delta_{\beta\lambda}(x - y) \quad (2.368)$$

$$+ \partial_\mu \Delta_{M\lambda}(zn - y) \partial_\nu \Delta_{N\gamma}(\sigma n - x) \partial_\alpha \partial_\omega \Delta_{\beta\kappa}(x - y) \quad (2.369)$$

nightmare continues. I have to CHECK whether the double derivative on propagator contributes with the

additional term. I think not, that is why i ignore it. Check it(imperative)

$$v^{\iota\alpha\beta\gamma}v^{\theta\omega\kappa\lambda} \quad (2.370)$$

$$\left( \partial_\mu \partial_\omega \Delta_{M\kappa}(zn - y) \partial_\nu \partial_\alpha \Delta_{N\beta}(\sigma n - x) \Delta_{\gamma\lambda}(x - y) \right. \quad (2.371)$$

$$- \partial_\mu \Delta_{\lambda M}(zn - y) \partial_\nu \partial_\alpha \Delta_{N\beta}(\sigma n - x) \partial_\omega \Delta_{\gamma\kappa}(x - y) \quad (2.372)$$

$$- \partial_\mu \partial_\omega \Delta_{M\kappa}(zn - y) \partial_\nu \Delta_{N\gamma}(\sigma n - x) \partial_\alpha \Delta_{\beta\lambda}(x - y) \quad (2.373)$$

$$\left. + \partial_\mu \Delta_{M\lambda}(zn - y) \partial_\nu \Delta_{N\gamma}(\sigma n - x) \partial_\alpha \partial_\omega \Delta_{\beta\kappa}(x - y) \right) \quad (2.374)$$

$$= v^{\iota\alpha\beta\gamma}v^{\theta\omega\kappa\lambda} \frac{(-1)^3 \Gamma^3\left(\frac{d}{2} - 1\right)}{(4\pi^{\frac{d}{2}})^3} \quad (2.375)$$

$$\left( \frac{4(zn - y)_\mu (y - zn)_\omega}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{4(\sigma n - x)_\nu (x - \sigma n)_\alpha}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{\left(\frac{d}{2} - 1\right)^2}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}-1}} \eta_{M\kappa} \eta_{N\beta} \eta_{\gamma\lambda} \right. \quad (2.376)$$

$$- \frac{-2(zn - y)_\mu}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{4(\sigma n - x)_\nu (x - \sigma n)_\alpha}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{-2(y - x)_\omega}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{d}{2} \left(\frac{d}{2} - 1\right)^3 \eta_{M\lambda} \eta_{N\beta} \eta_{\gamma\kappa} \quad (2.377)$$

$$- \frac{4(zn - y)_\mu (y - zn)_\omega}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{-2(\sigma n - x)_\nu}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{-2(x - y)_\alpha}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{d}{2} \left(\frac{d}{2} - 1\right)^3 \eta_{M\kappa} \eta_{N\gamma} \eta_{\beta\lambda} \quad (2.378)$$

$$+ \frac{-2(zn - y)_\mu}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{-2(\sigma n - x)_\nu}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{4(y - x)_\omega (x - y)_\alpha}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{d}{2} \left(\frac{d}{2} - 1\right)^3 \eta_{M\lambda} \eta_{N\gamma} \eta_{\beta\kappa} \quad (2.379)$$

now 3 variables for integration  $u_1, u_2, u_3$  maybe. As it was done already for the exercise. introduce integration variables and the notations

$$\Lambda = u_1 u_2 + u_2 u_3 + u_3 u_1 \quad (2.380)$$

$$\bar{u}_1 = \frac{u_2 u_3}{\Lambda} \text{first shift:} \quad (2.381)$$

$$x \rightarrow x + on + \bar{w}y \quad (2.382)$$

The values for the shifts should be looked up in Mathematica. FOLLOWING FORMULA NOT CORRECT

$$v^{\iota\alpha\beta\gamma}v^{\theta\omega\kappa\lambda} \frac{(-1)^3 \Gamma\left(\frac{3d}{2} + 1\right)}{(4\pi^{\frac{d}{2}})^3} \int [du_1 du_2 du_3] \frac{1}{(-Kx^2 - f(u_1 - 3, n, y) + i\epsilon)^{d+1}} 16 \quad (2.383)$$

$$\left( -y_\mu (y - zn)_\omega - x_\nu (x - \sigma n)_\alpha \eta_{M\kappa} \eta_{N\beta} \eta_{\gamma\lambda} \right. \quad (2.384)$$

$$- \frac{-2(zn - y)_\mu}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{4(\sigma n - x)_\nu (x - \sigma n)_\alpha}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{-2(y - x)_\omega}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{d}{2} \left(\frac{d}{2} - 1\right)^3 \eta_{M\lambda} \eta_{N\beta} \eta_{\gamma\kappa} \quad (2.385)$$

$$- \frac{4(zn - y)_\mu (y - zn)_\omega}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{-2(\sigma n - x)_\nu}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{-2(x - y)_\alpha}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{d}{2} \left(\frac{d}{2} - 1\right)^3 \eta_{M\kappa} \eta_{N\gamma} \eta_{\beta\lambda} \quad (2.386)$$

$$+ \frac{-2(zn - y)_\mu}{(-(zn - y)^2 + i\epsilon)^{\frac{d}{2}}} \frac{-2(\sigma n - x)_\nu}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{4(y - x)_\omega (x - y)_\alpha}{(-(x - y)^2 + i\epsilon)^{\frac{d}{2}+1}} \frac{d}{2} \left(\frac{d}{2} - 1\right)^3 \eta_{M\lambda} \eta_{N\gamma} \eta_{\beta\kappa} \quad (2.387)$$

I denote the operator for twist 2 as the following object:



diagram	correction
$A$	$\frac{g^2 C_A}{4\pi^{\frac{d}{2}} \epsilon} \int_0^1 du \frac{u}{u} (\mathcal{X}_{\mu,\nu}^A(z_{21}^u, z_2) - \mathcal{X}_{\mu,\nu}^A(z_1, z_2))$
$A'$	$\frac{-g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \left[ \left( 1 - \ln \left( \frac{ip_+^{z_1}}{\delta} \right) \right) \mathcal{X}_{\mu,\nu}^A(z_1, z_2) + \int_0^1 du \frac{u}{u} (\mathcal{X}_{\mu,\nu}^A(z_{21}^u, z_2) - \mathcal{X}_{\mu,\nu}^A(z_1, z_2)) \right]$
$B$	$\frac{g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \int_0^1 du \frac{u}{u} (\mathcal{X}_{\mu,\nu}^A(z_1, z_{12}^u) - \mathcal{X}_{\mu,\nu}^A(z_1, z_2))$
$B'$	$\frac{-g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \left( 1 - \ln \left( \frac{ip_+^{z_2}}{\delta} \right) \right) \mathcal{X}_{\mu,\nu}^A(z_1, z_2)$
$A2$	$-\frac{g^2 C_A}{8\pi^2 \epsilon} \mathcal{X}_{\mu,\nu}^A(z_1, z_2)$
$B2$	$-\frac{g^2 C_A}{8\pi^2 \epsilon} \mathcal{X}_{\mu,\nu}^A(z_1, z_2)$

Table 2: gluon twist 2

The correction is in one-loop. with  $z_1 = z$ ,  $z_2 = \sigma$ ,  $z_{ij}^w = z_i + w(z_j - z_i)$  (from first index to second index running along  $w$ ) and  $p_+^x$  denoting on which argument the momentum operator is acting.

$$\mathcal{X}_{\mu,\nu}^A(z_1, z_2) = F_{\mu+}^B(z_1 n) [z_1 n, z_2 n]^{BC} F_{\nu+}^D(z_2 n) T_{CE}^D [z_2 n, -\infty n]^{EA} \quad (2.388)$$

$$= F_{\mu+}^B(z_1 n) F_{\nu+}^D(z_2 n) T_{BA}^D \quad (2.389)$$

in lc gauge and then i re-represent the results

Sum together should

Diagram A:

$$F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.390)$$

$$\rightarrow \quad (2.391)$$

$$F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.392)$$

$$+ F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_\ell^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.393)$$

I claim that the computation is - after computing color - 100% identical to the computation for twist 1. The intermediate field is simply ignored. (one of course has to add the diagram  $A'$ ). color is

$$(\delta^{AB'} \delta^{BC'} - \delta^{AC'} \delta^{BB'}) f^{A'B'C'} T_{AC}^B T_{CE}^D = -i f^{A'AB} f^{BAC} T_{CE}^D (1 - 1) \quad (2.394)$$

$$= i C_A T_{A'E}^D \quad (2.395)$$

so if i trust my intuition i end with the expression

$$\pm \int_{-\infty}^{\pm z} d\sigma \frac{g^2 C_A}{4\pi^2 \epsilon} \left( -1 + \ln \left( \frac{\pm i \hat{p}_+}{\delta} \right) \right) F_{\mu+}^{A'}(\pm z n) F_{\nu+}^D(\sigma n) T_{A'E}^D \quad (2.396)$$

all +-es are for field at  $zn$ , and all --es are for other case. Minuses i am not so sure, should check. Most likely there is a sign flip in this expression. I need to check it.

Diagram B: here i show full computation. we start with the same expression as for diagram A.

$$F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.397)$$

$$\rightarrow \quad (2.398)$$

$$= F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.399)$$

$$+ F_{\mu M}^A i g B_+^B T_{AC}^B F_{\nu N}^D T_{CE}^D \left( -i g \int d^d x A_l^{A'}(x) \partial_{x^\alpha} B_\beta^{B'}(x) B_\gamma^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.400)$$

unfortunately i forgot to put arguments, on the fields of the operator, but they are clear.

the color factor requires the identity for 3 structure function product. it can be easily derived from Jacobi-identity (for structure functions) plus identity for product of two structure functions. the formula reads

$$f^{DEA} f^{EFB} f^{FDC} = \frac{1}{2} C_A f^{ABC} \quad (2.401)$$

The color factor computation for this diagrams is

$$T_{AC}^B T_{CE}^D f^{A'B'C'} (\delta^{BC} \delta^{DB'} + \delta^{BB'} \delta^{DC'}) = -(-i)^2 f^{CBA} f^{BDA'} f^{DCE} (1-1) \quad (2.402)$$

$$= \frac{-C_A}{2} (-i)^2 f^{AA'E} (1-1) \quad (2.403)$$

again the minus is between different possibilities of contracting the fields. We can remove the remaining structure function back to the color matrix to restore the initial expression.

$$T_{AC}^B T_{CE}^D f^{A'B'C'} (\delta^{BC} \delta^{DB'} + \delta^{BB'} \delta^{DC'}) = \frac{-i C_A}{2} T_{AE}^{A'} (1-1) \quad (2.404)$$

The terms yield

$$-i^2 g^2 F_{\mu M}^A \int d^d x A_l^{A'}(x) \frac{-i C_A}{2} T_{AE}^{A'} v^{\iota\alpha\beta\gamma} \quad (2.405)$$

$$\Delta_{+\gamma}(\tau n - x) \partial_\nu \partial_\alpha \Delta_{N\beta}(\sigma n - x) - \partial_\alpha \Delta_{+\beta}(\tau n - x) \partial_\nu \Delta_{N\gamma}(\sigma n - x) \quad (2.406)$$

always keep in mind to which argument the derivatives are related. I do omit the extra term arising from double derivative on the propagator, since its contribution is null (Lorentz contracted is 0)

$$-i^2 g^2 F_{\mu M}^A \int d^d A_l^{A'}(x) \frac{-i C_A}{2} T_{AE}^{A'} v^{\iota \alpha \beta \gamma} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^{\frac{d}{2}}} \quad (2.407)$$

$$\frac{1}{(-(\tau n - x)^2 + i\epsilon)^{\frac{d}{2}-1}} \frac{4 \left( \frac{d}{2} - 1 \right) \frac{d}{2} (\sigma n - x)_\nu (x - \sigma n)_\alpha}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}+1}} \eta_{\gamma+} \eta_{N\beta} \quad (2.408)$$

$$- \frac{-2 \left( \frac{d}{2} - 1 \right) (x - \tau n)_\alpha}{(-(\tau n - x)^2 + i\epsilon)^{\frac{d}{2}}} \frac{-2 \left( \frac{d}{2} - 1 \right) (\sigma n - x)_\nu}{(-(\sigma n - x)^2 + i\epsilon)^{\frac{d}{2}}} \Gamma(d) \eta_{\beta+} \eta_{N\gamma} \quad (2.409)$$

$$-i^2 g^2 F_{\mu M}^A \int d^d A_l^{A'}(x) \frac{-i C_A}{2} T_{AE}^{A'} v^{\iota \alpha \beta \gamma} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^{\frac{d}{2}}} 4 \frac{1}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} \quad (2.410)$$

$$\int_0^1 du u^{\frac{d}{2}-2} \bar{u}^{\frac{d}{2}} \frac{-x_\nu (x - \sigma n)_\alpha}{(-u(\tau n - x)^2 - \bar{u}(\sigma n - x)^2 + i\epsilon)^d} \eta_{\gamma+} \eta_{N\beta} \quad (2.411)$$

$$+ \int_0^1 du u^{\frac{d}{2}-1} \bar{u}^{\frac{d}{2}-1} \frac{x_\nu (x - \tau n)_\alpha}{(-u(\tau n - x)^2 - \bar{u}(\sigma n - x)^2 + i\epsilon)^d} \eta_{\beta+} \eta_{N\gamma} \quad (2.412)$$

The shifts in this setup are:

$$\tau \rightarrow \tau + \sigma \text{ first shift} \quad (2.413)$$

$$x \rightarrow x + u\tau n + \sigma n \quad (2.414)$$

$$x - \sigma n \rightarrow x + u\tau n \quad (2.415)$$

$$x - \tau n \rightarrow x - \bar{u}\tau n \quad (2.416)$$

compute metric

$$\eta_{\gamma+} \eta_{N\beta} v_{\iota \alpha \beta \gamma} = 2(\eta_{\iota N} \eta_{\alpha+} - \eta_{\iota+} \eta_{\alpha N}) \quad (2.417)$$

$$= -\eta_{\beta+} \eta_{N\gamma} v_{\iota \alpha \beta \gamma} \quad (2.418)$$

Hence another relative sign between term and identical metric term Have:

$$-i^2 g^2 F_{\mu M}^A \int d^d A_l^{A'}(x + u\tau n + \sigma n) \frac{-i C_A}{2} T_{AE}^{A'} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^{\frac{d}{2}}} 4 \frac{1}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} \int_0^1 du u^{\frac{d}{2}-2} \bar{u}^{\frac{d}{2}-1} 2(\eta_{\iota N} \eta_{\alpha+} - \eta_{\iota+} \eta_{\alpha N}) \quad (2.419)$$

$$\frac{-\bar{u}x_\nu (x + u\tau n)_\alpha - ux_\nu (x - \bar{u}\tau n)_\alpha}{(-x^2 + i\epsilon)^d} = \frac{-x_\nu x_\alpha}{(-x^2 + i\epsilon)^d} \quad (2.420)$$

Then from integration we have double derivative

$$\int d^d x \frac{-x_\nu x_\alpha x_\rho x_\kappa}{(-x^2 + i\epsilon)^d} \partial_\rho \partial_\kappa A^{A'}(u\tau n + \sigma n) = \frac{-i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} (\eta_{\nu\alpha} \eta_{\rho\kappa} + \eta_{\nu\rho} \eta_{\alpha\kappa} + \eta_{\nu\kappa} \eta_{\alpha\rho}) \partial^\kappa \partial^\rho A_l^{A'}(\sigma n + u\tau n) \quad (2.421)$$

now again contract metrics:

$$(\eta_{\nu\alpha}\eta_{\rho\kappa} + \eta_{\nu\rho}\eta_{\alpha\kappa} + \eta_{\nu\kappa}\eta_{\alpha\rho}) \partial^\kappa \partial^\rho A_\nu^{A'} (\sigma n + u\tau n) (\eta_{\iota N}\eta_{\alpha+} - \eta_{\iota+}\eta_{\alpha N}) = 2\partial_+ (\partial_\nu A_+ - \partial_+ A_\nu)^{A'} (\sigma n + u\tau n) \quad (2.422)$$

Total , combination of last two steps

$$-i^2 g^2 F_{\mu M}^A \frac{-iC_A}{2} T_{AE}^{A'} \Gamma^2 \left( \frac{d}{2} - 1 \right) \frac{1}{4^2 \pi^{\frac{d}{2}}} 4 \frac{1}{\Gamma^2 \left( \frac{d}{2} - 1 \right)} \quad (2.423)$$

$$\int_0^1 du u^{\frac{d}{2}-2} \bar{u}^{\frac{d}{2}-1} 2 \partial_+ (\partial_\nu A_+ - \partial_+ A_\nu)^{A'} (\sigma n + u\tau n) \frac{-i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} \quad (2.424)$$

$$= \int_0^1 du \bar{u} \frac{(-1)^3 i^4 g^2 C_A^2}{4^3 2^1 \pi^2 \epsilon} F_{\mu M}^A(zn) T_{AE}^{A'} \partial_+ F_{\nu+}^{A'} (\sigma n + u\tau n) \quad (2.425)$$

of course, there are the two integrals, over  $\tau$  and  $\sigma$ . first is necessary for the computation, the other one belongs to the operator itself. i already shifted the  $\tau$  integral by  $+\sigma$ . Now i want to subtract  $\sigma$  again but add  $z$ . I also show the integral over  $\sigma$  now, which is from  $-\infty$  to  $z$ , and i also shift this one. I end up

$$\int_{-\infty}^0 d\tau \int_0^1 du \bar{u} \partial_+ f(y + u\tau n) = - \int \frac{d^d p}{(2\pi)^d} \left( \ln \left( \frac{ip_+}{\delta} \right) - 1 \right) e^{i(p|y)} f(p) \quad (2.426)$$

$$= \left( 1 - \ln \left( \frac{ip_+}{\delta} \right) \right) f(\sigma n) \quad (2.427)$$

so in my case here:

$$\frac{-g^2 C_A}{2(4\pi)^2 \epsilon} F_{\mu M}^A(zn) \int_{-\infty}^z d\sigma \left( 1 - \ln \left( \frac{ip_+(\sigma)}{\delta} \right) \right) F_{\nu+}^{A'}(\sigma n) T_{AE}^{A'} \quad (2.428)$$

the argument of  $p_+$  indicates on which field its acting.

The next diagram i want to compute is diagram AB. I think its the last out three diagrams for twist 2 which are not absolutely the same. Probably this is the most interesting. the first two are the same computation as for twist 1, essentially.

the therm is

$$\overline{F_{\mu+}^A F_{\nu+}^D T_{AE}^D} \quad (2.429)$$

$$\rightarrow g f^{ABC} \overline{B_\mu^B B_+^C} g f^{DFH} \overline{B_\nu^F B_+^H} T_{AE}^D \quad (2.430)$$

I only have this contraction, since all other 3 possibilities yield zero Color computation yields:

$$+ig^2 f^{DAE} f^{ABC} f^{BDH} \Delta_{\mu\nu}(zn - \sigma n) \quad (2.431)$$

$$= + \frac{g^2 C_A}{2} f^{ECH} \Delta_{\mu\nu}(zn - \sigma n) \quad (2.432)$$

$$= \frac{-g^2 C_A}{2} T_{HE}^C \Delta_{\mu\nu}(zn - \sigma n) A_+^C A_+^H \quad (2.433)$$

Maybe one can argue that these drop since the outgoing fields are classical so they drop in lc-gauge?

0 diagrams from metric: the following diagrams are 0 for the reason  $\eta_{++} = \eta_{\mu+} = \eta_{\nu+} = 0$ :  
 $A1, A1', B1, B1', E1, E2, E3$  The remaining computation is for the diagrams A2 and B2. It should be very similar to what i have seen for twist 1:

$$gf^{ABC} B_{\mu}^B B_{+}^C(zn) F_{\nu+}^D T_{AE}^D \left( -ig \int d^d x A_{\mu}^{A'}(x) \partial_{x^{\alpha}} B_{\beta}^{B'}(x) B_{\gamma}^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.434)$$

$$\rightarrow gf^{ABC} \overbrace{B_{\mu}^B B_{+}^C(zn) F_{\nu+}^D T_{AE}^D} \left( -ig \int d^d x A_{\mu}^{A'}(x) \partial_{x^{\alpha}} B_{\beta}^{B'}(x) B_{\gamma}^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.435)$$

$$+ gf^{ABC} \overbrace{B_{\mu}^B B_{+}^C(zn) F_{\nu+}^D T_{AE}^D} \left( -ig \int d^d x A_{\mu}^{A'}(x) \partial_{x^{\alpha}} B_{\beta}^{B'}(x) B_{\gamma}^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.436)$$

apart from color it is identical to twist one, so i will not do perform the calculation in all details. color gives

$$f^{ABC} T_{AE}^D f^{A'B'C'} (\delta^{BB'} \delta^{CC'} + \delta^{BC'} \delta^{CB'}) = -i f^{ABC} f^{DAE} f^{A'BC} (1 - 1) \quad (2.437)$$

$$= -i C_A f^{DA'E} \quad (2.438)$$

$$= C_A T_{A'E}^D \quad (2.439)$$

So color is identical ( gives factor  $C_A$  and connects color indices) Apart from that i see no difference, either.  
 So the result is the same:

$$- \frac{g^2 C_A}{4\pi^2 \epsilon} F_{\mu+}^{A'}(zn) \int_{-\infty}^z d\sigma F_{\nu+}^D(\sigma n) T_{A'E}^D \quad (2.440)$$

last diagram: B2

$$F_{\mu+}^A(zn) gf^{DBC} B_{\nu}^B B_{+}^C(\sigma n) T_{AE}^D \left( -ig \int d^d x A_{\mu}^{A'}(x) \partial_{x^{\alpha}} B_{\beta}^{B'}(x) B_{\gamma}^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.441)$$

$$\rightarrow F_{\mu+}^A(zn) gf^{DBC} \overbrace{B_{\nu}^B B_{+}^C(\sigma n) T_{AE}^D} \left( -ig \int d^d x A_{\mu}^{A'}(x) \partial_{x^{\alpha}} B_{\beta}^{B'}(x) B_{\gamma}^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.442)$$

$$+ F_{\mu+}^A(zn) gf^{DBC} \overbrace{B_{\nu}^B B_{+}^C(\sigma n) T_{AE}^D} \left( -ig \int d^d x A_{\mu}^{A'}(x) \partial_{x^{\alpha}} B_{\beta}^{B'}(x) B_{\gamma}^{C'}(x) \right) v_{A'B'C'}^{\iota\alpha\beta\gamma} \quad (2.443)$$

COLOR:


$$f^{DBC} T_{AE}^D f^{A'B'C'} (\delta^{BB'} \delta^{CC'} + \delta^{BC'} \delta^{CB'}) = \delta^{A'D} T_{AE}^D C_A (1 - 1) \quad (2.444)$$


again, the rest of the computation must be exactly the same. here the point for the fixed field is  $\sigma n$  and not  $zn$  but it is not integrated over (here) so it must be identical. the color gives the same result (just connects and  $C_A$ ) FINAL:

$$- \frac{g^2 C_A}{4\pi^2 \epsilon} F_{\mu+}^{A'}(zn) \int_{-\infty}^z d\sigma F_{\nu+}^D(\sigma n) T_{A'E}^D \quad (2.445)$$

diagram	correction
$A$	$\frac{g^2 C_A}{4\pi^{\frac{d}{2}} \epsilon} F_{\nu+}^D(\sigma n) T_{CE}^D \int_0^1 du \frac{u}{\bar{u}} (F_{\mu+}^C(zn + \bar{u}(\sigma - z)n) - F_{\mu+}^C(zn))$
$A'$	$\frac{-g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \left(1 - \ln\left(\frac{ip_+}{\delta}\right)\right) F_{\mu+}^{A'}(zn) F_{\nu+}^D(\sigma n) T_{A'E}^D$
$B$	$\frac{g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} F_{\mu M}^A \int_0^1 du \frac{u}{\bar{u}} (F_{\nu+}^{A'}(\sigma n + \bar{u}(z - \sigma)n) - F_{\nu+}^{A'}(\sigma n)) T_{AE}^{A'}$
$B'$	$F_{\mu M}^A \frac{-g^2 C_A}{8\pi^{\frac{d}{2}} \epsilon} \left(1 - \ln\left(\frac{ip_+}{\delta}\right)\right) F_{\nu+}^{A'}(\sigma n) T_{AE}^{A'}$
$A2$	$-\frac{g^2 C_A}{8\pi^2 \epsilon}$
$B2$	$-\frac{g^2 C_A}{8\pi^2 \epsilon}$

Table 3: gluon twist 2; the "correction" is the factor to the "default" expression. Together it is the result for the one loop diagram with the corresponding Name. For example, diagram  $B + B' = \frac{-g^2 C_A}{2(4\pi)^2 \epsilon} F_{\mu M}^A(zn) \int_{-\infty}^z d\sigma \left(1 - \ln\left(\frac{ip_+(\sigma)}{\delta}\right)\right) F_{\nu+}^{A'}(\sigma n) T_{AE}^{A'}$ . In blue the "default" expression, for clarification.





### III Tipps and Tricks

- $\int_0^1 du f(u, \bar{u}) = \int_0^1 d\bar{u} f(\bar{u}, u)$
- Gamma function properties
- index  $\mu$  is transverse!!!! has more implications than one might think!

Auxiliaries:

$$\partial_{y^\alpha} \frac{1}{(-(y-w)^2 + i\epsilon)^n} = \frac{-2n(y-w)_\alpha}{(-(y-w)^2 + i\epsilon)^{n+1}} \quad (3.1)$$

$$\partial_{w^\beta} \partial_{y^\alpha} \frac{1}{(-(y-w)^2 + i\epsilon)^n} = \frac{4n(n+1)(y-w)_\alpha(w-y)_\beta}{(-(y-w)^2 + i\epsilon)^{n+2}} + \frac{2n\eta_{\alpha\beta}}{(-(y-w)^2 + i\epsilon)^{n+1}} \quad (3.2)$$

$$\int_{-\infty}^0 d\sigma \int_{-\infty}^\sigma d\tau f(\sigma, \tau) = \int_{-\infty}^0 d\sigma \int_{-\infty}^0 d\tau f(\sigma + \tau, \tau) \quad (3.3)$$

$$\int d^d x \frac{x^\mu x^\nu}{(-x^2 + i\epsilon)^{d-1}} = \frac{-i\pi^{\frac{d}{2}}}{2(2 - \frac{d}{2})} \frac{\eta^{\mu\nu}}{\Gamma(d-1)} \quad (3.4)$$

own adaption for four elements in numerator:

$$\int d^d x \frac{x^\mu x^\nu x^\rho x^\sigma}{(-x^2 + i\epsilon)^d} = \frac{i\pi^{\frac{d}{2}}}{4(2 - \frac{d}{2})\Gamma(d)} (\eta^{\mu\nu}\eta^{\rho\sigma} + \eta^{\mu\rho}\eta^{\nu\sigma} + \eta^{\mu\sigma}\eta^{\nu\rho}) \quad (3.5)$$

$$\int_{-\infty}^0 d\tau \int_0^1 du \bar{u} \partial_+ f(y + u\tau n) = - \int \frac{d^d p}{(2\pi)^d} \left( \ln \left( \frac{ip_+}{\delta} \right) - 1 \right) e^{i(p|y)} f(p) \quad (3.6)$$

$$= \left( 1 - \ln \left( \frac{ip_+}{\delta} \right) \right) f(\sigma n) \quad (3.7)$$

$$\int_0^x d\tau \int_0^1 du u \partial_+ f(y + \bar{u}\tau n) = \int_0^1 du \frac{u}{\bar{u}} [f(y + \bar{u}x) - f(y)] \quad (3.8)$$

## IV Diagrams.

