

Assignment No.4

Valli Devi Bolla

Download all python codes from

<https://github.com/Vallidevibolla/Assignment-4/blob/main/code.py>

and latex-tikz codes from

<https://github.com/Vallidevibolla/Assignment-4/blob/main/main.tex>

Question taken from

https://github.com/gadepall/ncert/blob/main/linalg/linear_forms/gvv_ncert_linear_forms.pdf-Q. no.2.1

1 QUESTION No.2.1

Check which of the following are solutions of the equation

$$(1 \quad -2) \mathbf{x} = 4 \quad (1.0.1)$$

$$(a) \begin{pmatrix} 0 \\ 2 \end{pmatrix} (b) \begin{pmatrix} 4 \\ 0 \end{pmatrix} (c) \begin{pmatrix} 2 \\ 0 \end{pmatrix} (d) \begin{pmatrix} \sqrt{2} \\ 4\sqrt{2} \end{pmatrix} (e) \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (1.0.2)$$

2 SOLUTION

Given

$$(1 \quad -2) \mathbf{x} = 4 \quad (2.0.1)$$

$$\text{Let } \mathbf{A} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{B} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \quad (2.0.3)$$

$$\mathbf{C} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad (2.0.4)$$

$$\mathbf{D} = \begin{pmatrix} \sqrt{2} \\ 4\sqrt{2} \end{pmatrix} \quad (2.0.5)$$

$$\mathbf{E} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (2.0.6)$$

Let 'y' be the solution then equation be

$$\mathbf{y} = (1 \quad -2) \mathbf{x} \quad (2.0.7)$$

Substitute(2.0.2)in (2.0.7)

$$\mathbf{x} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \text{in}(2.0.7) \quad (2.0.8)$$

$$\mathbf{y} = (1 \quad -2) \begin{pmatrix} 0 \\ 2 \end{pmatrix} \quad (2.0.9)$$

$$\boxed{\mathbf{y} = -4} \quad (2.0.10)$$

Substitute (2.0.3)in (2.0.7)

$$\mathbf{x} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \text{in}(2.0.7) \quad (2.0.11)$$

$$\mathbf{y} = (1 \quad -2) \begin{pmatrix} 4 \\ 0 \end{pmatrix} \quad (2.0.12)$$

$$\boxed{\mathbf{y} = 4} \quad (2.0.13)$$

Substitute(2.0.4)in (2.0.7)

$$\mathbf{x} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \text{in}(2.0.7) \quad (2.0.14)$$

$$\mathbf{y} = (1 \quad -2) \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad (2.0.15)$$

$$\boxed{\mathbf{y} = 2} \quad (2.0.16)$$

Substitute(2.0.5)in (2.0.7)

$$\mathbf{x} = \begin{pmatrix} \sqrt{2} \\ 4\sqrt{2} \end{pmatrix} \text{in}(2.0.7) \quad (2.0.17)$$

$$\mathbf{y} = (1 \quad -2) \begin{pmatrix} \sqrt{2} \\ 4\sqrt{2} \end{pmatrix} \quad (2.0.18)$$

$$\boxed{\mathbf{y} = -7\sqrt{2}} \quad (2.0.19)$$

Substitute (2.0.6)in (2.0.7)

$$\mathbf{x} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{in}(2.0.7) \quad (2.0.20)$$

$$\mathbf{y} = (1 \quad -2) \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (2.0.21)$$

$$\boxed{\mathbf{y} = -1} \quad (2.0.22)$$

$\therefore x = \begin{pmatrix} 4 \\ 0 \end{pmatrix}$ i.e., Point B is the solution of the equation
 $(1 \ -2)x = 4$




Fig. 2.1: Solution