1

Assignment 6

K.A. Raja Babu

Download all python codes from

https://github.com/ka-raja-babu/Matrix-Theory/tree/main/Assignment6/Codes

and latex-tikz codes from

https://github.com/ka-raja-babu/Matrix-Theory/ tree/main/Assignment6

1 Question No. 2.29

Find the coordinates of the focus, axis, the equation of the directrix and latus rectum of the parabola $y^2 = 8x$.

2 Solution

Para-	Sym-	Value	General
meter	bol	value	Formula
Vertex	С	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} \mathbf{u}^T + \eta \mathbf{p_1}^T \\ \mathbf{V} \end{pmatrix} \mathbf{c}$ $= \begin{pmatrix} -f \\ \eta \mathbf{p_1} - \mathbf{u} \end{pmatrix}$
Focal	β	2	$\left \begin{array}{c c} \frac{1}{4} & \frac{2\eta}{\lambda_2} \end{array} \right $
Length			$\overline{4} \overline{\lambda_2} $
Focus	F	$\begin{pmatrix} 2 \\ 0 \end{pmatrix}$	$\mathbf{F} = \mathbf{c} + \mathbf{a}^T$
Axis		$\begin{pmatrix} 0 & 1 \end{pmatrix} \mathbf{x} = 0$	$\mathbf{x} = \mathbf{F} + k\mathbf{m}$
Direct-		$(1 0)\mathbf{x} = -2$	$\mathbf{v_1}\mathbf{x} = -\beta$
rix		$\begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = -2$	$\mathbf{v_1x} = -\rho$
Latus		$(1 0)\mathbf{x} = 2$	$\mathbf{v_2}\mathbf{x} = \boldsymbol{\beta}$
rectum		$(1 0) \mathbf{x} = 2$	$\mathbf{v}_2\mathbf{x} = \boldsymbol{\rho}$
End			
points	M, N	(2) (2)	$ \beta \rangle$
of latus	''-, ' '	$\begin{pmatrix} 2 \\ 4 \end{pmatrix}, \begin{pmatrix} 2 \\ -4 \end{pmatrix}$	$\begin{pmatrix} \beta \\ \pm y(\beta) \end{pmatrix}$
rectum		· / · /	(/
Length			
of latus	<i>l</i>	8	$ \mathbf{M} - \mathbf{N} $
rectum			

TABLE 2.1: Parameters of parabola $y^2 = 8x$

All parameters of parabola $y^2 = 8x$ can be summarised in table 2.1

Note : Given general formula is valid only when parabola is in standard form i.e. |V| = 0 and $\lambda_1 = 0$

Given parabola is

$$y^2 = 8x (2.0.1)$$

$$\implies y^2 - 8x = 0 \tag{2.0.2}$$

Vector form of given parabola is

$$\mathbf{x}^{T} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} -4 & 0 \end{pmatrix} \mathbf{x} + 0 = 0 \tag{2.0.3}$$

:.

$$\mathbf{V} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \mathbf{u} = \begin{pmatrix} -4 \\ 0 \end{pmatrix}, f = 0 \tag{2.0.4}$$

|V| = 0 and $\lambda_1 = 0$ i.e. it is in standard form

$$\mathbf{P} = \mathbf{I} \implies \mathbf{p_1} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \tag{2.0.5}$$

$$\eta = \mathbf{u}^T \mathbf{p_1} = -4 \tag{2.0.6}$$

The vertex \mathbf{c} is given by

$$\begin{pmatrix} -8 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{c} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \tag{2.0.7}$$

$$\implies \mathbf{c} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \tag{2.0.8}$$

The focal length β is given by

$$\beta = \frac{1}{4} \left| \frac{2\eta}{\lambda_2} \right| = \frac{1}{4} \left| \frac{-8}{1} \right| = 2 \tag{2.0.9}$$

The focus **F** is given by

$$\mathbf{a} = \frac{-2\eta \begin{pmatrix} 1 & 0 \end{pmatrix}}{4} = \begin{pmatrix} 2 & 0 \end{pmatrix} \tag{2.0.10}$$

$$\mathbf{F} = \mathbf{c} + \mathbf{a}^T = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \tag{2.0.11}$$

 \therefore Axis of parabola passes through both vertex and focus .

:. Axis of parabola is given by

$$\mathbf{m} = \mathbf{F} - \mathbf{c} = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \tag{2.0.12}$$

$$\mathbf{x} = \mathbf{F} + k\mathbf{m} = \begin{pmatrix} 2 + 2k \\ 0 \end{pmatrix} \tag{2.0.13}$$

$$\implies \begin{pmatrix} 0 & 1 \end{pmatrix} \mathbf{x} = 0 \tag{2.0.14}$$

- : Vertex of parabola is at equal distance from focus and directrix and is perpendicular to axis.
- .. Directrix of parabola is given by

$$\begin{pmatrix} 0 & 1 \end{pmatrix} \mathbf{v_1}^T = 0 \tag{2.0.15}$$

$$\implies \mathbf{v_1} = \begin{pmatrix} 1 & 0 \end{pmatrix} \tag{2.0.16}$$

So,

$$\mathbf{v_1}\mathbf{x} = -\beta \tag{2.0.17}$$

$$\implies \begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = -2 \tag{2.0.18}$$

- : Latus rectum of parabola passes through focus and is perpendicular to axis.
- :. Latus rectum of parabola is given by

$$\begin{pmatrix} 0 & 1 \end{pmatrix} \mathbf{v_2}^T = 0 \tag{2.0.19}$$

$$\implies \mathbf{v_2} = \begin{pmatrix} 1 & 0 \end{pmatrix} \tag{2.0.20}$$

So,

$$\mathbf{v_2}\mathbf{x} = \beta \tag{2.0.21}$$

$$\implies \begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{x} = 2 \tag{2.0.22}$$

End points of latus rectum are

$$\mathbf{M} = \begin{pmatrix} 2\\4 \end{pmatrix} \tag{2.0.23}$$

$$\mathbf{N} = \begin{pmatrix} 2 \\ -4 \end{pmatrix} \tag{2.0.24}$$

So, the length of latus rectum l is

$$l = ||\mathbf{M} - \mathbf{N}|| = 8 \tag{2.0.25}$$

Plot of given parabola




Fig. 2.1: Parabola $y^2 = 8x$