

Quiz 3

Matthew Farah
June 28, 2026

Contents

Problem 1	3
(a)	3
(b)	3
(c)	4
(d)	4
Problem 2	6
Problem 3	7
Problem 4	8
Problem 5	9
Problem 6	11
(a)	11
(b)	11
(c)	11
Problem 7	13
Problem 8	14
(a)	14
(b)	19
Problem 9	20
Problem 10	21
(a)	21
(b)	21
(c)	21
(d)	21
(e)	23
(f)	23
Problem 11	25
Problem 12	26
(a)	26
(b)	26
(c)	26
(d)	26
(e)	27
Problem 13	28

Please note the quiz is based on the following topics:

- Conjugation.
- Normal Subgroups.

Problem 1:

In the group D_4 , let $r = (1234)$ be a clockwise rotation, and let $s = (12)(34)$ be a reflection. Which of the following elements are equal to sr ?

(a) rs .


Solution:

rs does not equal sr .

For this and the following questions, we will use the fact that:

$$\begin{aligned} sr &= (12)(34)(1234) \\ &= (24) \end{aligned}$$


Which graphically looks like the following adjacency-preserving permutation of the vertices of a square:



to determine whether or not the given permutations are equivalent to sr . In this case, since:

$$\begin{aligned} rs &= (1234)(12)(34) \\ &= (13) \end{aligned}$$

Which can be graphically represented by the transformation:



It is clear that rs and sr are different permutations, and thus, are unequal.


(b) r^2s .

Solution:

r^2s doesn't equal sr as well. To see why, we can leverage the fact that $rs = (13)$ (which we determined in part (a)) to conclude that:

$$\begin{aligned}
 r^2 s &= r(rs) \\
 &= (1234)(13) \\
 &= (14)(23)
 \end{aligned}$$

Which visually looks like this:



And since this is clearly a different permutation than sr , $r^2 s \neq sr$.

(c) $r^3 s$.


Solution:

$r^3 s$ actually does equal sr !

Proceeding as we did before, since:

$$\begin{aligned}
 r^3 s &= r(r^2 s) \\
 &= (1234)(14)(23) \\
 &= (24)
 \end{aligned}$$

Which looks like:



We can see that $r^3 s$ is indeed the same permutation as sr , and so $r^3 s = sr$.

(d) $r^4 s$.


Solution:

$r^4 s$ doesn't equal sr .

This is because:

$$\begin{aligned} r^4 s &= r(r^3 s) \\ &= (1234)(24) \\ &= (12)(34) \end{aligned}$$

Or, graphically:



Thus, $r^4 s$ and sr are clearly different permutations and so $r^4 s \neq sr$.

Problem 2:

Let G be a group, and let $a \in G$. What notations represent the conjugacy class of a ?

Solution:

a^G and $[a]$ are both ways of denoting the conjugacy class of a . Note that a^G is perhaps a little more descriptive than $[a]$, since it explicitly states which group it is a conjugacy class of.

Problem 3:

Let G be a group and let $a, b \in G$. What is the definition of a^b ?

Solution:

a^b is defined by:

$$a^b = b^{-1}ab$$

Problem 4:

If $\sigma = (153)(24)$ and $g = (5314)$, then what does σ^g equal?

Solution:

By the definition of σ^g given as the solution to the problem just above, we know that:

$$\begin{aligned}\sigma^g &= g^{-1}\sigma g && \text{(By definition of conjugation)} \\ &= (5314)^{-1}(153)(24)(5314) && \text{(Substituting)}\end{aligned}$$

So, to compute σ^g , we'll need to find g^{-1} ! This is thankfully not very difficult to do. Although I won't prove it (:P), it is generally true that the inverse of any permutation of the form $(abc\dots z)$ is just $(zxy\dots a)$, and so $g^{-1} = (4135)$.

So carrying out our earlier computation, we have that:

$$\begin{aligned}\sigma^g &= (5314)^{-1}(153)(24)(5314) && \text{(From earlier equation)} \\ &= (4135)(153)(24)(5314) && \text{(Since } (5314)^{-1} = (4135)\text{)} \\ &= (4135)(12435) && \text{(Simplifying)} \\ &= (12)(345) && \text{(Simplifying)}\end{aligned}$$

Thus, $\sigma^g = (12)(345)$.

Problem 5:

Find all elements $g \in S_3$ such that $g^{-1}(123)g = (132)$.

Solution:

We know that $S_3 = \{(), (12), (23), (13), (123), (132)\}$ so, together with the quick and easy method for finding the inverse of a permutation I mentioned in the solution to the problem above, we can brute force all possible $g \in S_3$ to find that:

(I) When $g = ()$:

$$\begin{aligned} g^{-1}(123)g &= ()(123)() \\ &= ()(123) \\ &= (123) \end{aligned}$$

(II) When $g = (12)$:

$$\begin{aligned} g^{-1}(123)g &= (21)(123)(12) \\ &= (21)(13) \\ &= \boxed{(132)} \end{aligned}$$

(III) When $g = (23)$:

$$\begin{aligned} g^{-1}(123)g &= (32)(123)(23) \\ &= (32)(12) \\ &= \boxed{(132)} \end{aligned}$$

(IV) When $g = (13)$:

$$\begin{aligned} g^{-1}(123)g &= (31)(123)(13) \\ &= (31)(23) \\ &= \boxed{(132)} \end{aligned}$$

(V) When $g = (123)$:

$$\begin{aligned}g^{-1}(123)g &= (321)(123)(123) \\&= (321)(132) \\&= (123)\end{aligned}$$

(VI) When $g = (132)$:

$$\begin{aligned}g^{-1}(123)g &= (231)(123)(132) \\&= (231)() \\&= (231) \\&= (123)\end{aligned}$$

Thus, the set of all g in S_3 satisfying $g^{-1}(123)g = (132)$ is $\{(12), (23), (13)\}$.

Problem 6:

How many elements do the following conjugacy classes contain?

(a) $[(123)]$ in S_4 .

Solution:

$[(123)]$ in S^4 has 8 elements.

By Theorem 5.3, we know that (123) is conjugate to another permutation, $\tau \in S_4$, if and only if (123) and τ have the same cycle type.

The cycle type of (123) in S_4 is $(3, 1)$ and so all permutations in S_4 conjugate to (123) must likewise have a cycle type of $(3, 1)$. Therefore, all permutations in S_4 conjugate to (123) are expressible as (abc) , where $a, b, c \in \{1, 2, 3, 4\}$ and a, b , and c are all pairwise distinct.

Clearly then, there are $4! = 24$ possible such permutations in S_4 , since:

- There are 4 possible choices for a .
- 3 choices for b so as to not conflict with a .
- 2 for c to avoid conflicting with both a and b .
- Only one choice as to which element the permutation leave fixed, after a, b , and c have all been chosen.

However, recall that the permutations (abc) , (bca) , and (cab) are all equivalent, meaning 24 over counts the number of distinct such permutations in S_4 by a factor of 3. Thus, there are actually $\frac{24}{3} = 8$ permutations with a cycle type of $(3, 1)$ in S_4 .

Thus, since there are exactly 8 permutations in S_4 with the same cycle type as (123) , there are also exactly 8 elements in the conjugacy class of (123) .

(b) $[(123)]$ in A_4 .

Solution:

(c) $[5]$ in $\mathbb{Z}_6^\times$.

Solution:

$[5]$ in $\mathbb{Z}_6^\times$ has 1 element.

$\mathbb{Z}_6^\times$ is a relatively small group, so we will find the size of $[5]$ in $\mathbb{Z}_6^\times$ by brute force. By Proposition 1.7, we know that because:

$$\begin{array}{l} \gcd(0, 6) = 6 \\ \boxed{\gcd(1, 6) = 1} \\ \gcd(2, 6) = 2 \end{array}$$

$$\begin{array}{l} \gcd(3, 6) = 3 \\ \gcd(4, 6) = 2 \\ \boxed{\gcd(5, 6) = 1} \end{array}$$

1 and 5 are the only units modulo 6. Thus, $\mathbb{Z}_6^\times = \{1, 5\}$. Now, since:

$$\begin{aligned}5^1 &\equiv (1)^{-1}(5)(1) \pmod{6} \\&\equiv (1)(5)(1) \pmod{6} \\&\equiv (1)(5) \pmod{6} \\&\equiv 5 \pmod{6} \\&\equiv 5\end{aligned}$$

and:

$$\begin{aligned}5^5 &\equiv (5)^{-1}(5)(5) \pmod{6} \\&\equiv (5)(5)(5) \pmod{6} \\&\equiv (5)(25) \pmod{6} \\&\equiv 125 \pmod{6} \\&\equiv 5\end{aligned}$$

We therefore have that the conjugacy class of 5 in $\mathbb{Z}_6^\times$ is just the set containing 5. Thus, since $[5] = \{5\}$, $[5]$ contains only one element.

Problem 7:

How many conjugacy classes are there in S_5 .

Solution:

There are 7 conjugacy classes in S_5 .

By Theorem 5.3 and the fact that conjugacy classes form equivalence partitions over their domain, we know there must be as many conjugacy classes in S_5 as there are distinct cycle types in S_5 .

Since there are 7 distinct cycle types in S_5 , given by the set $\{ (1, 1, 1, 1, 1), (2, 1, 1, 1), (2, 2, 1), (3, 1, 1), (3, 2), (4, 1), (5) \}$, (derived from the integer partitions of 5), there must therefore be 7 conjugacy classes in S_5 , as argued earlier.

Problem 8:

Which of the following subgroups are normal subgroups of $G = GL_2(\mathbb{Z}_2)$?

$$(a) H = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \right\}$$

Solution:

H is a normal subgroup of G .

To see why, recall that for H to be a normal subgroup of G , it must be closed under conjugation, meaning that for all $g \in G$ and $h \in H$, $h^g \in H$. Since I couldn't really think of a better way, our approach will be to compute h^g for all possible h and g , but cleverly take shortcuts where possible to cut down on the number of required calculations.

For one, since $h^g = g^{-1}hg$, it'd be convenient to have an easy formula for g^{-1} , given g , that we can use to plug and chug with. It is for that reason that I will quickly sidetrack to prove that for any $g \in G$ of the form:

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$g^{-1} \in G$ is given by:

$$g^{-1} = \begin{pmatrix} d & b \\ c & a \end{pmatrix}$$

To understand how this is the case, all that's required is to understand two things:

1. For any $x \in \mathbb{Z}_2$, $2x = 0$.
 - This is because if $x = 0$, then $2x = (2)(0) \equiv 0$, and if $x = 1$, then $(2)(1) = 2 = 0 \pmod{2}$.
2. $ad + bc = 1$.
 - This is because exactly one of ad or bc can be 1, and the other must be 0.
 - Since $\det(g) = ad - bc \neq 0$, by definition of g belonging to $GL_2(\mathbb{Z}_2)$.

And so, we have that:

$$\begin{aligned}
\boxed{\begin{pmatrix} d & b \\ c & a \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}} &= \begin{pmatrix} ad + cb & db + bd \\ ca + ac & cb + ad \end{pmatrix} && \text{(Simplifying)} \\
&= \begin{pmatrix} ad + bc & 2bd \\ 2ac & ad + bc \end{pmatrix} && \text{(Simplifying)} \\
&= \begin{pmatrix} ad + bc & 0 \\ 0 & ad + bc \end{pmatrix} && \text{(Since } 2x = 0 \text{ for all } x \in \mathbb{Z}_2) \\
&= \boxed{\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}} && \text{(Since } ad + bc = 1) \\
&= \begin{pmatrix} bc + ad & 0 \\ 0 & bc + ad \end{pmatrix} && \text{(Since } bc + ad = 1) \\
&= \begin{pmatrix} bc + ad & 2ab \\ 2cd & bc + ad \end{pmatrix} && \text{(Since } 2x = 0 \text{ for all } x \in \mathbb{Z}_2) \\
&= \boxed{\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} d & b \\ c & a \end{pmatrix}}
\end{aligned}$$

And thus, $g^{-1} = \begin{pmatrix} d & b \\ c & a \end{pmatrix}$. So now that we know how to compute $g^{-1}hg$, we need only consider 3 possible cases, recalling that we need only check that h is closed under conjugation by $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, and $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, since all other matrices in $GL_2(\mathbb{Z}_2)$ belong to H , which is given to be a group.

(I) $h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

If $h = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, then we wouldn't even need to consider the different choices of g to see that:

$$\begin{aligned}
h^g &= g^{-1} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} g && \text{(By definition of conjugation)} \\
&= g^{-1}g && \text{(By definition of the identity)} \\
&= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} && \text{(By definition of inverses)}
\end{aligned}$$

And because $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \in H, h^g \in H$ in this case. (Note that we also could've determined this from the fact the conjugacy class of the identity element is always the set containing just itself).

$$\text{(II)} \quad h = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

Here, we'll need to consider each possible g individually.

$$\text{(a)} \quad g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

If $g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, then $g^{-1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, and so we can compute:

$$h^g = g^{-1}hg \quad \text{(By definition of conjugation)}$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{(Substituting)}$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \quad \text{(Simplifying)}$$

$$= \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{(Simplifying)}$$

And since $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \in H, h^g \in H$ in this case.

$$\text{(b)} \quad g = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

If $g = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, then $g^{-1} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, and so we can compute:

$$h^g = g^{-1}hg \quad \text{(By definition of conjugation)}$$

$$= \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \quad \text{(Substituting)}$$

$$= \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{(Simplifying)}$$

$$= \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{(Simplifying)}$$

And since $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \in H, h^g \in H$ in this case as well.

$$\text{(c)} \quad g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

If $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, then $g^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, and so we can compute:

$$h^g = g^{-1}hg \quad \text{(By definition of conjugation)}$$

$$= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{(Substituting)}$$

$$= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{(Simplifying)}$$

$$= \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{(Simplifying)}$$

And since $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \in H, h^g \in H$ once again.

Thus, since $h^g \in H$ in cases **(a)**, **(b)**, and **(c)**, we have that $h = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ in H is closed under conjugation.

$$\text{(III)} \quad h = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}.$$

We'll once again have to consider each case of g separately.

$$\text{(a)} \quad g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

When $g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, then $g^{-1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, and so:

$$h^g = g^{-1}hg \quad (\text{By definition of conjugation})$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (\text{Substituting})$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad (\text{Simplifying})$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \quad (\text{Simplifying})$$

And because $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \in H$, then $h^g \in H$ in this case.

(b) $g = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$.

Assuming $g = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, then $g^{-1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and:

$$h^g = g^{-1}hg \quad (\text{By definition of conjugation})$$

$$= \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \quad (\text{Substituting})$$

$$= \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (\text{Simplifying})$$

$$= \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad (\text{Simplifying})$$

And since $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \in H$, h^g belongs to H in this case as well.

(c) $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Finally, if $g = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, then $g^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and thus:

$$h^g = g^{-1}hg \quad (\text{By definition of conjugation})$$

$$= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad (\text{Substituting})$$

$$= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \quad (\text{Simplifying})$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \quad (\text{Simplifying})$$

And since $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \in H$, h^g belongs to H in this case as well.

Therefore, since h^g belongs to H in cases **(a)**, **(b)**, and **(c)**, we can therefore conclude that $h = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ in H is closed under conjugation.

Thus, since cases **(I)**, **(II)**, and **(III)** all show that the elements of H are closed under conjugation, H is closed under conjugation as well and is thus a normal subgroup of G . Phew!

$$\text{(b)} \quad K = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\}.$$

Solution:

K is not a normal subgroup of G . To see why, all we need is to observe that when $k = \begin{pmatrix} 0 & 1 & 1 & 0 \end{pmatrix}$ and $g = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$:

$$k^g = g^{-1}kg \quad (\text{By definition of conjugation})$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \quad (\text{Substituting})$$

$$= \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \quad (\text{Simplifying})$$

$$= \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

which doesn't belong to K , despite the fact that $k \in K$ and $g \in G$! Thus, K cannot be closed under conjugacy and thus doesn't constitute a normal subgroup of G .

Problem 9:

Consider the following proposition:

Proposition: Conjugation satisfies the following:

- (a) Let $a, g, h \in G$. Then $a^{gh} = (a^g)^h$.
- (b) Let $a, b, g \in G$. Then $(ab)^g = a^g b^g$.

Now, suppose that the definition of a^g was changed to $a^g = gag^{-1}$. How would claim (a) and (b) made by the proposition change, if at all?

Solution:

We'll consider the impact that this new definition of conjugacy would have on claims (a) and (b) separately.

(I) How would claim (a) change?

Claim (a) would change to become $a^{gh} = (a^h)^g$.

Using this new definition of conjugacy, we can see that for arbitrary $a, g, h \in G$:

$$\begin{aligned}
 a^{gh} &= (gh)a(gh)^{-1} && \text{(By our new definition of conjugation)} \\
 &= (gh)a(h^{-1}g^{-1}) && \text{(By exponent laws)} \\
 &= g(hah^{-1})g^{-1} && \text{(By associativity)} \\
 &= g(a^h)g^{-1} && \text{(By our new definition of conjugation)} \\
 &= (a^h)^g && \text{(By our new definition of conjugation)}
 \end{aligned}$$

And thus, $a^{gh} = (a^h)^g$ when conjugacy is defined this way.

(II) How would claim (b) change?

Claim (b) wouldn't change at all!

Using this new definition of conjugacy, we can see that for arbitrary $a, b, g \in G$:

$$\begin{aligned}
 (ab)^g &= g(ab)g^{-1} && \text{(By our new definition of conjugation)} \\
 &= g(a(1)b)g^{-1} && \text{(By definition of identity)} \\
 &= g(a(g^{-1}g)b)g^{-1} && \text{(By definition of inverse)} \\
 &= (gag^{-1})(gbg^{-1}) && \text{(By associativity)} \\
 &= a^g b^g && \text{(By our new definition of conjugation)}
 \end{aligned}$$

Thus, even with the definition of conjugacy changes, $(ab)^g$ still equals $a^g b^g$.

Problem 10:

Which of the following statements are true?

- (a) In an abelian group, every subgroup is normal.

Solution:

This statement is true.

Let G be any abelian group. To prove the statement true, we'll need to show that every subgroup of G , $H \leq G$, is a normal subgroup. So, let $H \leq G$ be an arbitrary subgroup of G , and let $h \in H$ and $g \in G$ be arbitrary elements of H and G respectively.

By definition of normalcy, H is a normal subgroup of G if $h^g = g^{-1}hg \in H$. Recall, however, that G is abelian, and so for arbitrary elements x and y , $xy = yx$. Thus, we know that:

$$\begin{aligned}
 h^g &= g^{-1}hg && \text{(By definition of conjugation)} \\
 &= g^{-1}gh && \text{(By commutativity, since } G \text{ is abelian)} \\
 &= (1)h && \text{(By definition of inverses)} \\
 &= h && \text{(By definition of identity)}
 \end{aligned}$$

And since $h \in H$ by assumption, we have that $h^g \in H$. Thus H constitutes a normal subgroup of G . Since H was arbitrary, we can thereby conclude that every subgroup of an abelian group G must be normal.

- (b) Every group has at least two normal subgroups.

Solution:

- (c) The intersection of two normal subgroups is also a normal subgroup.

Solution:

This statement, once again, is true!

Let G be an arbitrary group, and let $H \trianglelefteq G$ and $K \trianglelefteq G$ be any two normal subgroups of G . To prove the statement correct, we need to show that $H \cap K$ is itself always a normal subgroup of G . To do that, we need to show that $x^g \in (H \cap K)$ for all $x \in (H \cap K)$ and $g \in G$.

So, let $x \in (H \cap K)$ and $g \in G$ be arbitrary elements of their respective groups. By definition of set intersection, we know that $x \in H$ and $x \in K$. Since H and K were assumed to be normal subgroups of G , we therefore know that $x^g \in H$ and that $x^g \in K$, since both H and K must both be closed under conjugation. However, since $x^g \in H$ and $x^g \in K$, we have that $x^g \in (H \cap K)$.

Thus, since $x \in (H \cap K)$ and $g \in G$ implies $x^g \in (H \cap K)$, $H \cap K$ is closed under conjugation, making $H \cap K$ a normal subgroup of G . Because H , K , and G were all arbitrary, this altogether proves the statement.

- (d) If A is a normal subgroup of B , and B is a normal subgroup of C , then A is a normal subgroup of C .

Solution:

This statement is false.

We will prove this statement false by use of a neat little counter example. Consider the groups S_3 , A_3 , and $(123)^{A_3}$. To prove that the statement doesn't hold, we'll show that although $(123)^{A_3} \trianglelefteq A_3$ and $A_3 \trianglelefteq S_3$, $(123)^{A_3}$ is not a normal subgroup of S_3 .

Our first order of business will be to show that $(123)^{A_3} \trianglelefteq A_3$ and $A_3 \trianglelefteq S_3$.

(I) Prove $(123)^{A_3} \trianglelefteq A_3$.

This statement holds definitionally. Since $(123)^{A_3}$ is the set of all permutations in A_3 that (123) is conjugate to, and since conjugacy constitutes an equivalence relation over its domain, $(123)^{A_3}$ is necessarily closed under conjugation, making $(123)^{A_3}$ a normal subgroup of A_3 .

(II) Prove $A_3 \trianglelefteq S_3$.

Here, we'll try a clever approach to show that A_3 is closed under conjugation by relying on the very definition of A_3 itself.

Let $\sigma \in A_3$ and $\tau \in S_3$ be arbitrary permutations in their respective groups, and let P_σ and P_τ be their associated permutation matrices. By definition of A_3 , we know that $\det(P_\sigma) = 1$ since A_3 is the set of all even permutations in S_3 .

Because $\det(AB) = \det(A)\det(B)$ for arbitrary matrices in $M_n(\mathbb{R})$, and **Proposition 3.13 (a)** implies that $\det(P_{\sigma^{-1}}) = \frac{1}{\det(P_\sigma)}$ since:

$$\begin{aligned}
 \det(P_{\sigma^{-1}})\det(P_\sigma) &= \det(P_{\sigma^{-1}}P_\sigma) && \text{(By aforementioned property of the determinant)} \\
 &= \det(P_{\sigma\sigma^{-1}}) && \text{(By Proposition 3.13 (a))} \\
 &= \det(P_{\text{id}}) && \text{(By definition of inverse)} \\
 &= \det(I_3) && \text{(By definition of permutation matrices)} \\
 &= 1 && \text{(By definition of the identity)}
 \end{aligned}$$

We can thus conclude that:

$$\begin{aligned}
 \det(P_\sigma^\tau) &= \det(P_{\tau^{-1}\sigma\tau}) && \text{(By definition of conjugation)} \\
 &= \det(P_{\tau^{-1}}P_\sigma P_\tau) && \text{(By Proposition 3.13 (a))} \\
 &= \det(P_{\tau^{-1}})\det(P_\sigma)\det(P_\tau) && \text{(By aforementioned property of determinants)} \\
 &= \frac{1}{\det(P_\tau)}\det(P_\sigma)\det(P_\tau) && \text{(Since } P_{\sigma^{-1}} = \frac{1}{P_\sigma}\text{)} \\
 &= \det(P_\sigma) && \text{(Simplifying)} \\
 &= 1 && \text{(By assumption } \sigma \in A_3\text{)}
 \end{aligned}$$

And so, since σ^τ clearly belongs to S_3 , and its associated permutation matrix P_{σ^τ} has a determinant of 1, we know that σ^τ is an even permutation and so belongs to A_3 , by definition.

Because $\sigma \in A_3$ and $\tau \in S_3$ were arbitrary, we can therefore conclude that A_3 is closed under conjugation and is thus a normal subgroup of S_3 , as required.

Thus, considerations **(I)** and **(II)** together tell us that $(123)^{A_3} \trianglelefteq A_3$ and $A_3 \trianglelefteq S_3$. But is $(123)^{A_3}$ a normal subgroup of S_3 ? Well, since A_3 is an abelian group, we can appeal to part **(a)** of this problem to conclude that $(123)^{A_3} = \{(123)\}$. However, we also know that $(12) \in S_3$ satisfies:

$$\begin{aligned}(123)^{(12)} &= (21)(123)(12) \\ &= (21)(13) \\ &= (123)\end{aligned}$$

And yet, (132) doesn't belong to $(123)^{A_3}$! Thus, $(123)^{A_3}$ isn't closed under conjugation and so cannot be a normal subgroup of S_3 .

Therefore, since $(123)^{A_3} \trianglelefteq A_3$ and $A_3 \trianglelefteq S_3$, but $(123)^{A_3} \not\trianglelefteq S_3$, we found a counter example to the claim that normalcy is transitive.

(e) G is always a normal subgroup of itself.

Solution:

This statement is true, and is fairly easy to prove!

To show G is always its own subgroup, we need to show that $a^b \in G$ for all $a, b \in G$. Since G is a group, we know it must be closed under its group operation. Thus, there must be some $c \in G$ such that $ab = c$. Furthermore, G being a group also guarantees to us that G is closed under inversion as well. Thus, b^{-1} likewise belongs to G , implying there must exist some $d \in G$ to satisfy $d = b^{-1}c$.

Therefore, because:

$$\begin{aligned}a^b &= b^{-1}ab && \text{(By definition of conjugation)} \\ &= b^{-1}c && \text{(By definition of } c\text{)} \\ &= d && \text{(By definition of } d\text{)}\end{aligned}$$

and $d \in G$ by assumption, then a^b likewise belongs to G ! Thus, G is closed under conjugation and is thereby a normal subgroup of itself.

(f) Every group is partitioned into its conjugacy classes.

Solution:

This statement is true!

Recall that a group G is said to be partitioned by a collection of sets $\{P_1, \dots, P_n\}$ if the following two criteria are satisfied:

1. For all $g \in G$, there exists some P_i such that $x \in P_i$.
2. $P_i \cap P_j = \emptyset$ for all $i \neq j$.

So, to prove that a group G is partitioned by its conjugacy classes, we need to show that:

(I) Every element of G belongs to some conjugacy class.

This immediately follows as a corollary to **Exercise 5.1**, which asserts that conjugation constitutes an equivalence relation on any group G .

Since conjugation is an equivalence relation, it must necessarily be reflexive, meaning for arbitrary g in G , g is its own conjugate.

Thus, $g \in [g]$ at the very least, and since $[g]$ is evidently a conjugacy class of G , because $g \in G$ was arbitrary, we therefore have that every element in a group must belong to some conjugacy class.

(II) Every pair of distinct conjugacy classes are pairwise disjoint.

Let P and Q be any two conjugacy classes of G such that $P \neq Q$. To prove that they're pairwise disjoint, we need to show that this implies $P \cap Q = \emptyset$. This thankfully follows quite readily, yet again, from the fact conjugation is an equivalence relation on G .

Let's assume, for the purposes of deriving a contradiction, that $P \cap Q$ is nonempty. There must therefore exist some element $x \in (P \cap Q)$. By definition of set intersection, we thus have that $x \in P$ and $x \in Q$.

Let $p \in P$ and $q \in Q$ be two arbitrary elements of their respective conjugacy classes. Since x and p both belong to P , by definition of conjugacy classes, we know that $p\tilde{x}$. Similarly, since q and x both belong to Q , we also know $x\tilde{q}$.

Since conjugation is an equivalence relation, it must therefore be transitive. Thus, by definition of transitivity, since $p\tilde{x}$ and $x\tilde{q}$, we have that $p\tilde{q}$. But p and q were arbitrary! This means every element in P is conjugate to every element in Q and vice versa (since conjugation is also symmetric, owing to the fact it is an equivalence relation). Hence, by definition of conjugacy classes $P = Q$!

This contradicts our prior assumption that P and Q must be non-equal conjugacy classes of G . Thus, $P \cap Q$ must be empty, meaning any two distinct conjugacy classes satisfy this criterion as well! Thus, since conjugacy classes of any arbitrary group G always satisfy the criteria needed to partition G (as per the findings of considerations **(I)** and **(II)**), we can therefore conclude that every group must be partitioned by its conjugacy classes.

Problem 11:

In **Chapter 6**, the quaternion group Q_8 serves as a counter example to which conjecture?

Solution:

The conjecture was:

If every subgroup of G is normal, then G must be abelian.

Problem 12:

Complete the following computations in the quaternion group, Q_8 .

(a) ij .

Solution:

$ij = k$, since:

$$\begin{aligned}
 ij &= ij(1) && \text{(By definition of the identity)} \\
 &= ij(-1)^2 && \text{(Expanding)} \\
 &= ij(-1)(-1) && \text{(Expanding)} \\
 &= ijk^2(-1) && \text{(By fundamental identity of quaternions)} \\
 &= ijk k(-1) && \text{(Expanding)} \\
 &= (-1)k(-1) && \text{(By fundamental identity of quaternions)} \\
 &= k && \text{(Simplifying)}
 \end{aligned}$$

(b) ik .

Solution:

$ik = -j$ because:

$$\begin{aligned}
 ik &= i(1)k && \text{(By definition of the identity)} \\
 &= i(-1)(-1)k && \text{(Expanding)} \\
 &= i(ik)(-1)k && \text{(By fundamental identity of quaternions)} \\
 &= -i^2jk^2 && \text{(Simplifying)} \\
 &= jk^2 && \text{(By fundamental identity of quaternions)} \\
 &= -j && \text{(By fundamental identity of quaternions)}
 \end{aligned}$$

(c) ji .

Solution:

$ji = -k$ because:

$$\begin{aligned}
 ji &= ji(1) && \text{(By definition of the identity)} \\
 &= ji(-1)(-1) && \text{(Expanding)} \\
 &= ji(ik)(-1) && \text{(By fundamental identity of quaternions)} \\
 &= -ji^2jk && \text{(Simplifying)} \\
 &= j^2k && \text{(By fundamental identity of quaternions)} \\
 &= -k && \text{(By fundamental identity of quaternions)}
 \end{aligned}$$

(d) i^{-1} .

Solution:

$i^{-1} = -i$, since:

$$\begin{aligned}
 \boxed{(i)(-i)} &= -i^2 && \text{(Simplifying)} \\
 &= -(-1) && \text{(By fundamental identity of quaternions)} \\
 &= \boxed{1} && \text{(Simplifying)} \\
 &= -(-1) && \text{(Expanding)} \\
 &= -i^2 && \text{(By fundamental identity of quaternions)} \\
 &= \boxed{(-i)(i)} && \text{(Expanding)}
 \end{aligned}$$

(e) $(-j)^{-1}$.

Solution:

$(-j)^{-1} = j$ because:

$$\begin{aligned}
 \boxed{(-j)(j)} &= -j^2 && \text{(Simplifying)} \\
 &= -(-1) && \text{(By fundamental identity of quaternions)} \\
 &= \boxed{1} && \text{(Simplifying)} \\
 &= (-1)(-1) && \text{(Expanding)} \\
 &= (j^2)(-1) && \text{(By fundamental identity of quaternions)} \\
 &= (j)(j)(-1) && \text{(Expanding)} \\
 &= \boxed{(j)(-j)} && \text{(Simplifying)}
 \end{aligned}$$

Problem 13:

How many conjugacy classes does Q_8 have?

Solution: