


SYSTEMATIC REVIEW OPEN ACCESS

Evaluation of Complete Blood Count Parameters in Patients With Diabetes Mellitus: A Systematic Review

Mojtaba Aghaei^{1,2,3}  | Seyed Sobhan Bahreiny⁴  | Zeynab Deris Zayeri⁵  | Nader Davari²  |
Mohammad Mehdi Abolhasani³  | Najmaldin Saki^{1,2} 

¹Department of Medical Laboratory Science, School of Allied Medicine, Ahvaz Jundishapur University of Medical Science, Ahvaz, Iran | ²Health Research Institute, Thalassemia & Hemoglobinopathy Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran | ³Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran | ⁴Department of Physiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran | ⁵Golestan Hospital Clinical Research Development Unit, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Correspondence: Najmaldin Saki (Najmaldinsaki@gmail.com)

Received: 6 September 2024 | **Revised:** 10 January 2025 | **Accepted:** 21 January 2025

Keywords: blood cell count | diabetes mellitus | inflammation | leukocytes | mean platelet volume

ABSTRACT

Background and Aims: Several studies were performed to evaluate the relationship between CBC and patients with diabetes mellitus (DM). In this review, we discussed the prognostic value of CBC parameters in DM patients.

Methods: English literature was searched and retrieved from the Google Scholar search engine and PubMed database (1980–2024). “Diabetes mellitus,” “Blood cell count,” “Mean platelet volume,” “Leukocytes,” and “Inflammation” were used as keywords.

Results: DM increases vascular inflammation and oxidative stress, while vascular inflammation affects erythropoiesis and red blood cell deformation, thus increasing red cell distribution width (RDW). Mean platelet volume (MPV) is another useful prognostic biomarker for DM patients. Additionally, elevated neutrophil-lymphocyte ratio (NLR) levels are associated with poor glycemic control in T2DM patients, so it can be used as a screening tool in diabetic follow-up.

Conclusion: RDW can be used as a valuable independent biomarker to assess the prognosis of patients with DM. MPV can also be used as a noninvasive, widely available, and low-cost marker as a key factor as well as a Prognostic/diagnostic biomarker that could be used for DM patients. Total white blood cell count, NLR, Mean platelet volume lymphocyte ratio (MPVLR), and monocyte to high-density lipoprotein ratio (MHR) are valuable biomarkers in predicting DM.

1 | Introduction

Diabetes mellitus (DM) is a group of metabolic disorders characterized by chronic hyperglycemia, which results from defects in insulin secretion, insulin function, or insulin resistance (IR) [1–3]. The condition is most commonly classified into two main types: type 1 diabetes (T1DM) and type 2 diabetes (T2DM). T1DM, which is an autoimmune disorder, is characterized by the destruction of insulin-producing beta cells in the pancreas, resulting in an absolute deficiency of insulin [4–6]. In contrast, T2DM is primarily associated with insulin resistance

and relative insulin deficiency, where the body's cells become less responsive to insulin, leading to increased blood glucose levels. T2DM is the most prevalent form of diabetes and is strongly associated with obesity, physical inactivity, and an unhealthy diet, and its incidence has been increasing globally, contributing to a significant public health burden [7–9]. Both types of diabetes, however, share a common complication—hyperglycemia—which is a primary contributor to the development of various diabetic complications [10]. The complications of DM can be broadly categorized into microvascular and macrovascular disorders. Microvascular complications include

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2025 The Author(s). *Health Science Reports* published by Wiley Periodicals LLC.

diabetic retinopathy, nephropathy, and neuropathy, which can significantly impair the quality of life and lead to disability [11, 12]. Macrovascular complications, such as cardiovascular disease (CVD), stroke, and peripheral artery disease (PAD), are major contributors to the increased morbidity and mortality in diabetic patients. These complications are not only associated with prolonged hyperglycemia but also with a chronic inflammatory state that exacerbates disease progression [13–15]. In particular, Type 2 diabetes is known to be closely linked with systemic low-grade inflammation, which is thought to play a central role in the development of both microvascular and macrovascular complications. This inflammatory burden is often reflected by elevated levels of circulating inflammatory markers, such as C-reactive protein (CRP), interleukins, and tumor necrosis factor-alpha (TNF- α) and is considered a key mechanism in the pathophysiology of DM [16].

In addition to traditional biomarkers, several hemogram-derived markers have emerged as potential indicators of inflammation and disease progression in diabetes. These include mean platelet volume (MPV), red cell distribution width (RDW), and neutrophil-to-lymphocyte ratio (NLR). MPV is a marker of platelet activation and function, with increased levels often reflecting enhanced platelet reactivity and inflammation [17]. Elevated MPV levels have been observed in T2DM patients, as well as in those with diabetic kidney disease, and have been associated with an increased risk of cardiovascular events [18–21]. RDW, which reflects the variation in size of red blood cells, has been increasingly recognized as a marker of inflammation and oxidative stress. Elevated RDW levels have been reported in a wide range of inflammatory conditions, including autoimmune diseases, rheumatoid arthritis, malignancy, and CVD, as well as in DM patients, particularly those with poor blood sugar control and complications such as nephropathy [22–24]. RDW has also been identified as a predictor of adverse outcomes in various diseases, including mortality in diabetic patients [25–28]. Similarly, NLR, which represents the ratio of neutrophils to lymphocytes in the blood, is a robust marker of systemic inflammation. Increased NLR levels have been observed in conditions characterized by chronic inflammation, such as T2DM, thyroid disorders, CVD, and infections. In the context of DM, elevated NLR has been linked to insulin resistance and the development of diabetic complications, making it a potential prognostic marker for patients with T2DM [27, 29–32].

The growing body of evidence linking these hemogram-derived markers to inflammatory processes in diabetes has prompted further exploration into their role in monitoring disease progression and predicting outcomes. These markers, which are routinely measured as part of the complete blood count (CBC), offer a noninvasive, cost-effective, and accessible means of assessing inflammation and disease status in diabetic patients. Given the chronic inflammatory nature of T2DM and its complications, incorporating CBC parameters into clinical practice could provide valuable insights into disease mechanisms, allowing for better risk stratification, early detection of complications, and personalized management strategies [33].

This review aims to provide a comprehensive examination of the relationship between CBC parameters and diabetes

mellitus, focusing on markers such as MPV, RDW, and NLR. By synthesizing the available literature, we aim to highlight the potential of these biomarkers in understanding the pathophysiological mechanisms underlying diabetes and its complications, as well as their role in monitoring disease progression and prognosis. Ultimately, we hope to provide evidence supporting the integration of these markers into routine clinical practice as tools for improving the management and outcomes of diabetic patients.

2 | Methods

2.1 | Literature Search

This Systematic review was conducted following the PRISMA guidelines to systematically evaluate the relationship between CBC parameters and DM. We aimed to identify relevant studies that examined the prognostic value of various CBC indices in diabetic patients. The search terms included “Diabetes mellitus,” “Complete Blood Count,” “mean platelet volume,” “Red Cell Distribution Width,” “Leukocytes,” and “Inflammation.” We conducted searches across multiple databases, including PubMed, Scopus, and Google Scholar, from their inception up to October 2024. Additionally, we reviewed the reference lists of included studies to discover further relevant articles. This comprehensive search strategy yielded a total of 141 studies for initial review.

2.2 | Eligibility Criteria

Inclusion criteria for this study required that participants be diagnosed with diabetes mellitus, specifically type 1 or type 2, as defined by the American Diabetes Association. Studies included must have assessed CBC parameters such as RDW, MPV, and white blood cell (WBC) counts, focusing on their relationship with glycemic control and diabetes-related complications. Additionally, only studies published in English between 1980 and 2024 were considered, ensuring the accessibility and relevance of the literature. Exclusion criteria encompassed research that did not specifically examine CBC parameters in diabetic patients, studies focused on nondiabetic populations, and those lacking original data, such as reviews, commentaries, and conference abstracts. The exit qualities of the study aimed to ensure that the final selection of articles contributed valuable insights into the prognostic implications of CBC parameters in diabetes management, thereby enhancing clinical understanding and guiding future research directions.

2.3 | Data Extraction

To maintain rigor in our review process, two independent reviewers initially screened the titles and abstracts of the identified studies for relevance according to our inclusion and exclusion criteria. The selected articles underwent a full-text review by the same reviewers to confirm their suitability for inclusion. In cases of discrepancies or differing opinions between the reviewers, a third reviewer was involved to mediate

and reach a consensus. This systematic approach helped minimize selection bias, ensuring that only relevant and high-quality studies were included in our review.

3 | RDW as an Independent Prognostic Biomarker in DMs

RDW is a parameter of the change in size (anisocytosis) of circulating erythrocytes, which is usually obtained by a standard complete red blood cell count [34]. High RDW indicates anisocytosis and it is associated with impaired erythropoiesis and erythrocyte degradation, which reflects chronic inflammation and increased oxidative stress, which are obvious symptoms of T2DM and may significantly lead to atherosclerotic disease [35, 36]. Studies have shown an association between high RDW and adverse health outcomes such as increased mortality [37], increased incidence of atrial fibrillation and heart failure [38, 39] and poor prognosis in patients with heart failure or coronary artery disease [40, 41]. Increasing the rate of RDW changes is associated with increasing the incidence of DM [42]. In several studies on DM patients, higher mortality has been reported in primary coronary interventions (PCI) in individuals with high RDW [43]. Recent studies suggested that RDW may be helpful in assessing the risk of patients at risk for

T2DM. Veeranna et al. reported RDW has gradually increased among HBA1Cs group in 15,343 nondiabetic adults free of CVD based on using the National Health and Nutrition Examination Survey (NHANES) database [44]. Similarly, in a large population-based cohort study, 26,709 nondiabetic Swedes patients reported that RDW is positively associated with HBA1C so that every 1 SD increase in RDW results in a 0.10% increase in HBA1C [45]. DM increases vascular inflammation and oxidative stress, while vascular inflammation affects erythropoiesis and erythrocyte deformation, thus increase RDW [37, 40, 41, 46] (Figure 1). RDW is strongly associated with chronic inflammation and is a strong indicator of mortality risk in CVD, DM as well as in the general population. Inflammatory conditions, such as DM, cause red blood cells to deform due to multiple pathophysiological mechanisms [47]. It has recently been reported that hyperglycemia affects human RBCs by altering the content and dispersion of the three isotypes of tubulin, thereby reducing RBC deformation and osmotic resistance. Although tubulin is not the major protein in RBC base on volume, it plays an important structural role and regulates enzymatic activity such as Na⁺/K⁺ ATPase and PMCA (Plasma membrane ca²⁺ ATPase) [48]. Several studies and their almost identical results confirm the association between RDW and DM compared to other CBC parameters (Table 1). As a result, RDW can be considered as a valuable independent biomarker for DM prognosis.




FIGURE 1 | Diabetes mellitus and its complications. Diabetes mellitus can be classified into two classes, type 1 (absence of insulin secretion due to the destruction of beta cells) and type 2 (insulin resistance). Inadequate blood sugar control in diabetic people is associated with complications such as vascular inflammation, hypertriglyceridemia, hyperglycemia, and increased oxidative stress so that vascular inflammation affects the process of erythropoiesis in the bone marrow and changes the shape of RBCs, and ultimately causes anisocytosis. Hyperglycemia also leads to a decrease in deformation and osmotic resistance of RBCs through changes in the content and distribution of three isotypes of tubulin, which is a protein that regulates the activity of plasma membrane Ca²⁺ ATPase and Na⁺/K⁺ ATPase. On the other hand, hyperglycemia, insulin resistance, or destruction of beta cells through increasing thromboxane biosynthesis and calcium mobilization lead to an increase in MPV and subsequently to an increase in the reactivity of platelets that these cells have a greater tendency to thrombotic events and lead to complications such as atherothrombosis and micro and macrovascular complications. B.M, Bone marrow; PBS, Peripheral blood smear; PMCA, Plasma membrane ca²⁺ ATPase.

TABLE 1 | Summary of CBC parameters in the patients with diabetes mellitus.

Parameters	Index	M/F (%)	Median/ mean age	Number of participants	Clinical findings	Level of CBC parametr	p value	Ref.
WBC	Neu	64/56	30–90	DM:120 Control:120	Neutrophil count was statistically positively correlated with FBG, BMI and SBP	3.96 (2.27)	0.226	[49]
	Baso	64/56	30–90	DM:120 Control:120	Basophil count was statistically positively correlated with FBG, BMI and SBP	0.04 (0.03)	0.000	[49]
	Total WBC	48/52	66 [59–72]	DM:301 Control:359	Total WBC count was positively associated with HbA1c and higher in T2DM	7.29 (6.11–8.99)	< 0.001	[50]
	Mono	60/40	60.71 ± 9.07	DM:494 Control:1848	Increased monocyte count is associated with T2DM patients	0.47 ± 0.14	0.254	[51]
	Eos	46/54	57.7 ± 9.8	DM:37 Control:34	There was no statistically significant between eosinophil count and DM	0.25 (0.2)	0.51	[52]
RBC	Lymph	45/55	61 ± 9.8	DM:689 Control:NA	Low lymphocytes count were predictors of macrovascular complication and mortality in individuals with T2DM	1.8 ± 0.5	NA	[53]
	MCV	46/54	50 [40–60]	DM:105 Control:NA	Red blood cell indices such as MCV, MCH and MCHC were decreased in diabetic patients	87.52 ± 6 27.27 ± 2.71	0.016 0.054	[54]
	MCH							
	MCHC							
	Hgb	58/42	NA	DM:164 Control:82	The mean values of Hgb among DM patients were significantly lower than those of nondiabetic	31.09 ± 1.68 13.8 ± 1.8	0.829 < 0.001	[55]
PLT	HCT	48/52	56.34	DM:100 Control:100	There was no significant association	41.376	0.736	[56]
	RDW	63/37	73 (11.2)	DM:71 Control:147	RDW was associated with higher event rate both in HF patients with and without DM	15.2 (1.5)	0.744	[57]
	Plt	70/30	69.3 (±9.8)	DM:1272 Control:2210	Mean MPV values were significantly higher in diabetics than in non-diabetics	215.6 (±66.4) 11 (±0.99)	0.72 < 0.001	[58]
	MPV							

Abbreviations: BMI, body mass index; DM, diabetes mellitus; FBG, fasting blood glucose; HF, Heart failure; IR, insulin resistance; M/F, Male to Female ratio; MCH, Mean corpuscular hemoglobin concentration; MCV, Mean corpuscular volume; MPV, mean platelet volume; RDW, Red cell distribution width; SBP, systolic blood pressure.




FIGURE 2 | Inflammatory markers derived from CBC can predict DM events. WBC is an inflammatory marker in diabetes mellitus, which can be used as a risk factor for atherosclerosis, and its increase is associated with vascular complications in T2DM patients, metabolic syndrome, CVDs, CHDs, and cerebrovascular diseases, and it can be the possible cause of increased stiffness in T2DM patients. The increase of MHR is also a prognostic marker in CVDs and T2DM patients and it can be attributed to the reduction of HDL as an inflammatory lipid and the subsequent increase in CD11b expression. NLR is an important predictor of cardiac complications in diabetic patients, and its increase is related to poor blood sugar control in these patients. Factors such as apoptosis, oxidative DNA damage, and insufficient proliferation of lymphocytes lead to an increase in NLR. MPVLR is also known as an independent predictor of short-term and long-term prognosis in diabetic patients with myocardial infarction (MI), whose increase can be attributed to increased platelet activity and decreased lymphocytes. CHD, coronary heart disease; CVD, cardiovascular disease; HDL, High-density lipoprotein; MHR, monocyte to high-density lipoprotein ratio; MI, myocardial infarction; MPVLR, mean platelet volume lymphocyte ratio; NLR, neutrophil-lymphocyte ratio; T2DM, Type 2 diabetes mellitus; WBC, white blood cell.

4 | MPV as a Key Factor for DMs

Platelets are abundant cells in the bloodstream, which play a classic role in thrombosis and homeostasis [59]. Increased platelet activity plays an important role in the development of vascular complications in DM [60]. Although there are several measurements for platelet activity such as (platelet aggregometry, platelet surface p-selectin, platelet-activated glycoprotein GPIIb/IIIa, serum thromboxane B2, and urinary 11 dehydro thromboxane B2), almost all of these BIOMARKERS are time-consuming, expensive or require special training [61, 62]. Mean platelet volume (MPV) is an alternative marker for platelet activity that can be determined by conventional autologous hemograms as part of a relatively low-cost CBC test [63, 64]. Normal MPV values in a simple CBC test are between 7.2 and 11.7 fl, and higher MPV values indicate larger platelets that are more active metabolically and enzymatically and have greater anticoagulant potential [64–66]. In addition to being a risk factor for CVD, elevated MPV is also known as a risk factor for DM [8], impaired fasting glucose [67], hyperlipidemia [68], and metabolic syndrome [69]. Most published studies have been conducted to investigate the relationship between MPV, prevalence, severity of diabetes and its complications [70, 71]. Several

studies reported that MPV was significantly increased in DM cases and in patients with poor glycaemic control, and the duration of diabetes was longer. In addition, both groups of diabetic patients have higher MPV values compared to the nondiabetic group [72–75]. Diabetics, especially patients with T2DM, are prone to increased platelet reactivity due to factors such as metabolic disorders (hyperglycemia, hypertriglyceridemia), systemic disorders (such as oxidative stress, inflammation), and insulin resistance [76–78]. Atherothrombosis is a common complication in DM patients. Therefore, patients with T2DM are 2–4 times more likely to develop coronary artery disease and stroke [79]. Possible mechanisms include hyperglycemia or insulin resistance, which increases thromboxane biosynthesis and calcium mobilization and subsequently leads to increased platelet reactivity [80, 81]. Due to metabolic disorders, DM patients have overactive platelets that release more granules and are more prone to thrombotic events, which in turn leads to micro- and macrovascular complications in these patients, which in turn increases the risk of diabetes and increases mortality in these patients [82, 83]. Given the above, it can be concluded that MPV as a noninvasive marker, widely available and low cost can be used as a key prognostic and diagnostic biomarker in DM patients.

5 | Inflammatory Markers Derived From CBC Can Predict DM Events

Generally, the immune system response to inflammation is associated with an increase in WBC and neutrophil as well as a relative decrease in lymphocyte count [49, 50, 84]. Many epidemiological studies have shown an association between chronic inflammation and various inflammatory markers in DM pathogenesis so that chronic inflammation plays an important role in the acceleration and development of micro- and macroangiopathy complications [85–87] (Figure 2). Neutrophil lymphocyte ratio (NLR), is an important marker in systemic inflammation and an indicator of increased risk for cardiovascular events in patients with metabolic syndrome [88, 89] additionally, two other studies showed that NLR is an independent predictor, which is important for cardiac complications in diabetic patients [90, 91]. Elevated NLR levels are associated with poor glycemic control in T2DM patients. Therefore, it can be used as a screening tool in the follow-up of diabetic patients [92, 93]. Also, NLR is superior to other leukocyte parameters due to its stability to various physiological, pathological and physical factors [94, 95]. NLR is a combination of two major components of chronic inflammation (neutrophil increase and lymphocyte depletion). High neutrophil counts indicate a process of nonspecific and persistent destructive inflammation and lymphocyte depletion indicates relatively inadequate immune control as well as an inactive immune pathway [96]. Decrease in lymphocytes is attributed to the increased oxidative damage of DNA and apoptosis of peripheral blood lymphocytes as well as insufficient proliferation of lymphocytes due to low IL2 receptor expression [84, 97, 98]. In addition, NF-κB (Nuclear factor kappa-light-chain-enhancer of activated B cells) can be induced by stimuli such as hyperglycemia and oxidative stress, and the inflammatory response by increasing the expression of adhesion molecules such as ICAM 1 (Intercellular adhesion molecule 1) stimulates pro-inflammatory cytokines and chemokines, and this inflammatory response cascade will eventually lead to an increase in neutrophils [99, 100]. Simultaneously, WBC count is an essential marker in inflammation that elevated levels are associated with vascular complications in patients with T2DM and metabolic syndrome [50, 101–104], as well as increased WBC in atherosclerotic pathogenesis in patients with T2DM, CVD and mortality from coronary heart disease and cerebrovascular disease [105–108]. Additionally, increased arterial stiffness is associated with increased WBC in T2DM patients [109]. Mean platelet volume lymphocyte ratio (MPVLR) is also a new inflammatory marker in diabetic patients that has been recognized in recent studies as an independent predictor of short-term and long-term prognosis in diabetic patients with myocardial infarction (MI) [110, 111]. Increase in MPVLR is followed by decreasing lymphocyte count due to induction of

apoptosis, systemic inflammation, oxidative stress, calcium metabolism deficiency, decrease in nitric oxide, increase in phosphorylation and glycosylation of cellular proteins lead to increased platelet activation and subsequent increase in MPV and proinflammation in diabetic patients [112–115]. Like NLR, and MPVLR, monocyte to high-density lipoprotein ratio (MHR) is also known as a new marker associated with increased inflammation and oxidative stress as well as adverse cardiovascular outcomes [116]. Numerous studies have shown that increased monocytes are associated with insulin resistance (IR), T2DM, coronary artery disease, and micro- and macrovascular complications of diabetes [104, 117–119]. On the other hand, reduction of HDL as an anti-inflammatory lipid and antioxidant that has an antithrombotic effect in inflammatory conditions by inhibiting CD11b integrin activation that is involved in adhesion, migration, and regulation of monocyte/macrophage inflammatory activity leads to accelerated atherosclerosis in diabetic patients [120–122]. Therefore, increasing MHR can be used as a new marker of cardiovascular prognosis or inflammation in T2DM patients and its complications [51, 123]. See (Table 2).

6 | Evaluating Confounding Factors in CBC Parameters Related to Diabetes Mellitus

In the exploration of CBC parameters and their associations with DM, it is essential to recognize and evaluate various confounding factors that may influence these parameters. This section aims to provide a detailed analysis of how co-existing medical conditions and medication use can impact CBC results and subsequently affect the interpretation of their relationship with diabetes.

6.1 | Co-Existing Conditions

Co-existing medical conditions can significantly alter CBC results, leading to potential misinterpretations in the context of DM. For example, infections often trigger an inflammatory response that elevates WBC counts as the body attempts to combat pathogens. In patients with acute infections, leukocytosis (an increased WBC count) may occur, which could be mistakenly attributed to the effects of diabetes if the underlying infection is not properly identified [124]. Additionally, chronic infections can impact the bone marrow's function, subsequently affecting the production of RBCs and platelets. Anemia is another critical co-existing condition that can skew CBC interpretations [125–127]. In diabetic patients, the presence of anemia may lead to lower hemoglobin levels, complicating the assessment of HbA1c, a vital measure of long-term glucose control. Different types of anemia, such as iron deficiency or

TABLE 2 | Summary of inflammatory markers in the patients with diabetes mellitus.

Parameters	M/F (%)	Median/mean age	Patient no.	Statistical significance	p value	Ref.
NLR	34/66	66.60 ± 4.20	145	2.46 ± 1.26	0.001	[93]
MPVLR	36/64	65 [50–78]	42	3.9 (1.4–13.2)	0.02	[115]
MHR	60/40	60.71 ± 9.07	494	10.92 (8.34–14.02)	< 0.001	[51]

Abbreviations: MHR, monocyte high-density lipoprotein ratio; MPVLR, mean platelet volume lymphocyte ratio; NLR, neutrophil-lymphocyte ratio.

anemia of chronic disease, produce distinct effects on CBC readings, making it imperative to differentiate these conditions [128]. Therefore, a thorough understanding and assessment of co-existing medical conditions are crucial for accurately interpreting CBC parameters in relation to diabetes [124, 129].

6.2 | Medication Use

The influence of medication use on CBC parameters is a critical factor in understanding their relationship with DM. Various medications prescribed for diabetes management can have hematological side effects that may complicate CBC interpretations. For instance, metformin, a common first-line treatment for type 2 diabetes, has been associated with vitamin B12 deficiency, which can lead to macrocytic anemia [130, 131]. This condition often results in altered MCV and MCH values, making it challenging to accurately assess a patient's hematological status. Additionally, other pharmacological treatments, such as corticosteroids, can induce leukocytosis, while immunosuppressants may lead to leukopenia, both of which impact WBC counts and overall CBC results. It is essential to document all medications taken by patients to ensure that the influence of these drugs is accounted for when evaluating CBC parameters in the context of diabetes [131, 132]. By thoroughly considering medication use, researchers and clinicians can attain a clearer understanding of how these factors interact with diabetes, thereby enhancing the accuracy of their assessments and improving patient care.

6.3 | Statistical Considerations

Statistical considerations play a crucial role in accurately assessing the relationship between CBC parameters and DM by controlling for confounding factors. Employing multivariate statistical techniques allows researchers to adjust for the influence of co-existing medical conditions and medication use while evaluating the direct impact of diabetes on CBC results. This approach enables the isolation of specific effects attributed to DM, leading to more reliable and valid findings. Additionally, stratification of the analysis based on the presence or absence of certain confounding factors can enhance understanding of how these variables interact with diabetes. For example, examining CBC parameters in diabetic patients with and without anemia can reveal critical insights into the interplay between these conditions. By using appropriate statistical methods, researchers can ensure that their conclusions about the relationship between CBC parameters and diabetes are robust and clinically relevant, ultimately contributing to better-informed clinical practices and improved patient outcomes [133, 134].

7 | Limitations

In this study, which explores the prognostic value of CBC parameters in diabetic patients, it is crucial to acknowledge the body of evidence that presents conflicting or nonsignificant findings. While many studies have emphasized the importance of parameters such as NLR, MPV, RDW, and MHR in predicting

complications or disease progression in diabetes, several others have reported results that challenge these conclusions. For example, certain studies have failed to establish a statistically significant association between NLR and the severity of cardiovascular complications in diabetic populations, suggesting that its predictive power may be limited to specific subgroups or clinical contexts [135–137]. Similarly, MPV, although widely regarded as an indicator of platelet activation, has not shown consistent prognostic significance across all studies, with some investigations reporting no correlation with glycemic control or diabetic complications [138–140].

These discrepancies may stem from several factors, including variability in sample sizes, differences in the demographic and clinical characteristics of the study populations (e.g., age, comorbidities, or diabetes duration), or methodological variations such as the time frame of data collection and the definitions of outcomes. Furthermore, environmental factors, dietary habits, and genetic predispositions, which are often underexplored, could also contribute to the observed inconsistencies [136, 141, 142].

Recognizing these limitations, it becomes clear that while CBC parameters hold promise as prognostic tools, their application should be context-dependent. To solidify their utility, future studies should prioritize large-scale, multi-center trials with standardized methodologies, ensuring better comparability and reproducibility of findings. Additionally, exploring interactions between these parameters and other inflammatory or metabolic markers could provide a more comprehensive understanding of their role in diabetes pathophysiology.

Based on the diagnostic criteria for diabetes mellitus, all studies were included to ensure a comprehensive analysis with high sensitivity. However, a limitation of this study is the variability in the design of the included studies. Some studies focused on comparisons between diabetic and nondiabetic groups, while others examined diabetic patients with various complications. This heterogeneity may influence the interpretation of the results and the generalizability of the findings.

8 | Conclusion and Future Perspective

Nowadays the prognostic and diagnostic value of CBC parameters in DM patients become a hot research topic. It is predicted that CBC testing can be a cheap, accessible, simple, practical, and valuable test that can be used as a new prognostic tool for DM patients in health care centers. Therefore, the physician as the first line of diagnosis and follow-up of DM patients can determine the prognosis and treatment effects more confidently by standard principles of hematology clinical laboratories, especially CBC results. According to the previous studies the four parameters RDW, MPV, WBC, NLR are more relevant and efficient in determining the prognosis of DM patients compared to other CBC parameters. Additionally, we suggest that two new markers, MPVLR and MHR, can be used to determine the more accurate prognosis of DM patients. Additionally, considering the stability of RBC parameters such as Mean corpuscular volume (MCV), Mean corpuscular hemoglobin (MCH), and Mean corpuscular hemoglobin concentration (MCHC) in different populations, we

suggest that these parameters can be used separately or beside other inflammatory and platelet parameters.

Author Contributions

Mojtaba Aghaei: data curation, investigation, methodology, writing – review and editing, writing – original draft, conceptualization. **Seyed Sobhan Bahreiny:** methodology, writing – review and editing. **Zeynab Deris Zayeri:** investigation, methodology, writing – original draft. **Nader Davari:** software, data curation, writing – review and editing. **Mohammad Mehdi Abolhasani:** methodology, investigation. **Najmaldin Saki:** writing – review and editing, conceptualization, supervision.

Acknowledgments

We wish to thank all our colleagues at Ahvaz Jundishapur University of Medical Sciences.

Ethics Statement

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Transparency Statement

The lead author Najmaldin Saki affirms that this manuscript is an honest, accurate, and transparent account of the study being reported, that no important aspects of the study have been omitted, and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

References

1. A. I. Abdallah, A. A. Abdelaziz, E.-S. E.-M. El-Sayed, and T. M. Emran, “Study of Neutrophil Lymphocyte Ratio (NLR) and Platelet Lymphocyte Ratio (PLR) as a Predictor Inflammatory Marker for Diabetic Nephropathy in Type 2 Diabetic Patients,” *Egyptian Journal of Hospital Medicine* 72, no. 7 (2018): 4800–4807.
2. A. H. Mahdizade, S. S. Bahreiny, M.-N. Bastani, et al., “The Influence of CDKAL1 (rs7754840) Gene Polymorphism on Susceptibility to Gestational Diabetes Mellitus in Pregnant Women: A Systematic Review and Meta-Analysis,” *International Journal of Diabetes in Developing Countries* 44, no. 1 (2024): 3–12.
3. M. Aghaei, R. Khademi, S. S. Bahreiny, and N. Saki, “The Need to Establish and Recognize the Field of Clinical Laboratory Science (CLS) as an Essential Field in Advancing Clinical Goals,” *Health Science Reports* 7, no. 8 (2024): e70008.
4. A. Ziaee, A. Ghorbani, S. Kalbasi, A. Hejrati, and S. Moradi, “Association of Hematological Indices With Prediabetes: A Cross-Sectional Study,” *Electronic Physician* 9, no. 9 (2017): 5206–5211.
5. S. S. Bahreiny, A. Ahangarpour, M. Aghaei, R. Mohammadpour Fard, M. A. Jalali Far, and T. Sakharvarz, “A Closer Look at Galectin-3: Its Association With Gestational Diabetes Mellitus Revealed by Systematic Review and Meta-Analysis,” *Journal of Diabetes and Metabolic Disorders* 23, no. 2 (2024): 1621–1633.

6. N. Saki, H. Haybar, and M. Aghaei, “Subject: Motivation Can Be Suppressed, But Scientific Ability Cannot and Should Not Be Ignored,” *Journal of Translational Medicine* 21, no. 1 (2023): 520.
7. J. Barbieri, P. C. Fontela, E. R. Winkelmann, et al., “Anemia in Patients With Type 2 Diabetes Mellitus,” *Anemia* 2015 (2015): 1–7.
8. Z. Hekimsoy, B. Payzin, T. Örnek, and G. Kandoğan, “Mean Platelet Volume in Type 2 Diabetic Patients,” *Journal of Diabetes and Its Complications* 18, no. 3 (2004): 173–176.
9. E. Torfi, S. S. Bahreiny, N. Saki, et al., “Evaluation of Pro-BNP Biomarker in Heart Failure Patients and Its Relationship With Complete Blood Count Parameters: A Case–Control Study,” *Health Science Reports* 7, no. 9 (2024): e70083.
10. A. Salsali and M. Nathan, “A Review of Types 1 and 2 Diabetes Mellitus and Their Treatment With Insulin,” *American Journal of Therapeutics* 13, no. 4 (2006): 349–361.
11. E. Basaran and G. Aktas, “The Relationship of Vitamin D Levels With Hemogram Indices and Metabolic Parameters in Patients With Type 2 Diabetes Mellitus,” *AIMS Medical Science* 11, no. 1 (2024): 47–57.
12. G. Aktas, “Association Between the Prognostic Nutritional Index and Chronic Microvascular Complications in Patients With Type 2 Diabetes Mellitus,” *Journal of Clinical Medicine* 12, no. 18 (2023): 5952.
13. G. Aktas, T. T. Duman, and B. Atak Tel, “Controlling Nutritional Status (CONUT) Score Is a Novel Marker of Type 2 Diabetes Mellitus and Diabetic Microvascular Complications,” *Postgraduate Medicine* 136, no. 5 (2024): 496–503.
14. G. Aktas, “Serum C-Reactive Protein to Albumin Ratio as a Reliable Marker of Diabetic Neuropathy in Type 2 Diabetes Mellitus,” *Biomolecules and Biomedicine* 24, no. 5 (2024): 1380–1386.
15. G. Aktas, S. Yilmaz, D. B. Kantarci, et al., “Is Serum Uric Acid-to-HDL Cholesterol Ratio Elevation Associated With Diabetic Kidney Injury?,” *Postgraduate Medicine* 135, no. 5 (2023): 519–523.
16. C. E. Koro, S. J. Bowlin, N. Bourgeois, and D. O. Fedder, “Glycemic Control From 1988 to 2000 Among U.S. Adults Diagnosed With Type 2 Diabetes,” *Diabetes Care* 27, no. 1 (2004): 17–20.
17. G. Aktas, M. Z. Kocak, T. Taslamacioglu Duman, et al., “Mean Platelet Volume (MPV) as an Inflammatory Marker in Type 2 Diabetes Mellitus and Obesity,” *Bali Medical Journal* 7, no. 3 (2018): 650–653, <https://doi.org/10.15562/bmj.v7i3.806>.
18. T. T. Duman, G. Aktas, B. Atak, and M. Z. Kocak, “Is Mean Platelet Volume to Platelet Ratio a Promising Indicator of Diabetic Regulation in Type 2 Diabetes Mellitus,” *Journal of Medical Research* 4, no. 3 (2018): 137–139.
19. M. Z. Kocak, G. Aktas, E. Erkuş, T. T. Duman, and B. M. Atak, “Mean Platelet Volume to Lymphocyte Ratio as a Novel Marker for Diabetic Nephropathy,” *Jcpssp-Journal of The College of Physicians and Surgeons Pakistan* (2018), <https://doi.org/10.29271/jcpssp.2018.11.844>.
20. G. Aktas, M. Sit, H. Tekce, et al., “Mean Platelet Volume in Nasal Polyps,” *West Indian Medical Journal* 62, no. 6 (2013): 515–5188.
21. M. Torabizadeh, M. Aghaei, N. Saki, M. A. Vahid, S. Bitaraf, and B. Bandar, “The Association of Nasal and Blood Eosinophils With Serum IgE Level in Allergic Rhinitis and Asthma: A Case-Control Study,” *Health Science Reports* 7, no. 11 (2024): e70191.
22. G. Aktas, A. Alcelik, B. K. Tekce, V. Tekelioglu, M. Sit, and H. Savli, “Red Cell Distribution Width and Mean Platelet Volume in Patients With Irritable Bowel Syndrome,” *Gastroenterology Review* 3, no. 3 (2014): 160–163.
23. M. Aghaei, R. Khademi, M. A. J. Far, S. S. Bahreiny, A. H. Mahdizade, and N. Amirrajab, “Genetic Variants of Dectin-1 and Their Antifungal Immunity Impact in Hematologic Malignancies: A Comprehensive Systematic Review,” *Current Research in Translational Medicine* 72, no. 4 (2024): 103460.

24. S. A. Aghapour, M. Torabizadeh, S. S. Bahreiny, et al., "Investigating the Dynamic Interplay Between Cellular Immunity and Tumor Cells in the Fight Against Cancer: An Updated Comprehensive Review," *Iranian Journal of Blood and Cancer* 16, no. 2 (2024): 84–101.
25. S. Buse Balci and G. Aktas, "A Comprehensive Review of the Role of Hemogram Derived Inflammatory Markers in Gastrointestinal Conditions," *Iranian Journal of Colorectal Research* 10, no. 3 (2022): 75–86.
26. G. Aktas, M. Sit, O. Dikbas, et al., "Could Red Cell Distribution Width Be a Marker in Hashimoto's Thyroiditis?," *Experimental and Clinical Endocrinology & Diabetes: Official journal, German Society of Endocrinology [and] German Diabetes Association* 122, no. 10 (2014): 572–574.
27. L. Cakır, G. Aktas, O. B. Mercimek, O. Enginyurt, Y. Kaya, and K. Mercimek, "Are Red Cell Distribution Width and Mean Platelet Volume Associated With Rheumatoid Arthritis," *Biomedical Research* 27, no. 2 (2016): 292–294.
28. Y. Meng, X. L. Cai, S. Cong, et al., "Role of Platelet/Lymphocyte, Neutrophil/Lymphocyte, and Interleukin-37/Interleukin-17 Ratios in the Occurrence and Treatment of Rheumatoid Arthritis," *Immunological Investigations* 53, no. 3 (2024): 464–474, <https://doi.org/10.1080/08820139.2023.2299687>.
29. J. Zhang, Z. Gong, S. Li, et al., "The Value of Neutrophil-to-Lymphocyte Ratio Combined With the Thyroid Imaging Reporting and Data System in the Diagnosis of the Nature of Thyroid Nodules," *Journal of Clinical Laboratory Analysis* 36, no. 5 (2022): e24429, <https://doi.org/10.1002/jcla.24429>.
30. G. Aktas, D. U. M. A. N. Tt, A. T. A. K. Bm, et al., "Irritable Bowel Syndrome Is Associated With Novel Inflammatory Markers Derived From Hemogram Parameters," *Family Medicine & Primary Care Review* 22, no. 2 (2020), <https://doi.org/10.5114/fmpcr.2020.95311>.
31. G. Aktas, "Hematological Predictors of Novel Coronavirus Infection," *Revista da Associação Médica Brasileira* 67, no. Suppl 1 (2021): 1–2.
32. G. Aktas, M. Sit, O. Dikbas, et al., "Elevated Neutrophil-to-Lymphocyte Ratio in the Diagnosis of Hashimoto's Thyroiditis," *Revista da Associação Médica Brasileira* 63 (2017): 1065–1068.
33. I. Karagoz, G. Aktas, H. Yoldas, et al., "Association Between Hemogram Parameters and Survival of Critically Ill Patients," *Journal of Intensive Care Medicine* 34, no. 6 (2019): 511–513.
34. T. C. Evans and D. Jehle, "The Red Blood Cell Distribution Width," *Journal of Emergency Medicine* 9 (1991): 71–74.
35. B. Tesfamariam and R. A. Cohen, "Free Radicals Mediate Endothelial Cell Dysfunction Caused by Elevated Glucose," *American Journal of Physiology-Heart and Circulatory Physiology* 263, no. 2 (1992): H321–H326.
36. G. L. King and M. R. Loeken, "Hyperglycemia-Induced Oxidative Stress in Diabetic Complications," *Histochemistry and Cell Biology* 122 (2004): 333–338.
37. T. S. Perlstein, J. Weuve, M. A. Pfeffer, and J. A. Beckman, "Red Blood Cell Distribution Width and Mortality Risk in a Community-Based Prospective Cohort," *Archives of Internal Medicine* 169, no. 6 (2009): 588–594.
38. Y. Borné, J. G. Smith, O. Melander, B. Hedblad, and G. Engström, "Red Cell Distribution Width and Risk for First Hospitalization Due to Heart Failure: A Population-Based Cohort Study," *European Journal of Heart Failure* 13, no. 12 (2011): 1355–1361.
39. S. Adamsson Eryd, Y. Borné, O. Melander, et al., "Red Blood Cell Distribution Width Is Associated With Incidence of Atrial Fibrillation," *Journal of Internal Medicine* 275, no. 1 (2014): 84–92.
40. G. M. Felker, L. A. Allen, S. J. Pocock, et al., "Red Cell Distribution Width as a Novel Prognostic Marker in Heart Failure," *Journal of the American College of Cardiology* 50, no. 1 (2007): 40–47.
41. M. Tonelli, F. Sacks, M. Arnold, L. Moye, B. Davis, and M. Pfeffer, "Relation Between Red Blood Cell Distribution Width and Cardiovascular Event Rate in People With Coronary Disease," *Circulation* 117, no. 2 (2008): 163–168.
42. L. Gang and W. Lifang, "Association of the Elevated Red Blood Cell Distribution Width With the Risk of Developing Diabetes Mellitus," *Internal Medicine* 55, no. 15 (2016): 1959–1965.
43. S. Tsuboi, K. Miyauchi, T. Kasai, et al., "Impact of Red Blood Cell Distribution Width on Long-Term Mortality in Diabetic Patients After Percutaneous Coronary Intervention," *Circulation Journal* 77, no. 2 (2013): 456–461.
44. V. Veeranna, S. K. Zalawadiya, S. S. Panaich, K. Ramesh, and L. Afonso, "The Association of Red Cell Distribution Width With Glycated Hemoglobin Among Healthy Adults Without Diabetes Mellitus," *Cardiology* 122, no. 2 (2012): 129–132.
45. G. Engström, J. G. Smith, M. Persson, P. M. Nilsson, O. Melander, and B. Hedblad, "Red Cell Distribution Width, Haemoglobin A 1c and Incidence of Diabetes Mellitus," *Journal of Internal Medicine* 276, no. 2 (2014): 174–183.
46. G. Weiss and L. T. Goodnough, "Anemia of Chronic Disease," *New England Journal of Medicine* 352, no. 10 (2005): 1011–1023.
47. D. B. Kell and E. Pretorius, "Serum Ferritin Is an Important Inflammatory Disease Marker, as It Is Mainly a Leakage Product From Damaged Cells," *Metallomics* 6, no. 4 (2014): 748–773.
48. A. D. Nigra, N. E. Monesterolo, J. F. Rivelli, et al., "Alterations of Hemorheological Parameters and Tubulin Content in Erythrocytes From Diabetic Subjects," *International Journal of Biochemistry & Cell Biology* 74 (2016): 109–120.
49. H. Ebrahim, T. Fiseha, Y. Ebrahim, and H. Bisetegn, "Comparison of Hematological Parameters Between Type 2 Diabetes Mellitus Patients and Healthy Controls at Dessie Comprehensive Specialized Hospital, Northeast Ethiopia: Comparative Cross-Sectional Study," *PLoS One* 17, no. 7 (2022): e0272145.
50. A. Klisic, A. Scepanovic, J. Kotur-Stevuljevic, and A. Ninic, "Novel Leukocyte and Thrombocyte Indexes in Patients With Prediabetes and Type 2 Diabetes Mellitus," *European Review for Medical and Pharmacological Sciences* 26, no. 8 (2022): 2775–2781.
51. J. W. Chen, C. Li, Z. H. Liu, et al., "The Role of Monocyte to High-Density Lipoprotein Cholesterol Ratio in Prediction of Carotid Intima-Media Thickness in Patients With Type 2 Diabetes," *Frontiers in Endocrinology* 10 (2019): 191.
52. F. Sefil, K. T. Ulutas, R. Dokuyucu, et al., "Investigation of Neutrophil Lymphocyte Ratio and Blood Glucose Regulation in Patients With Type 2 Diabetes Mellitus," *Journal of International Medical Research* 42, no. 2 (2014): 581–588.
53. C. R. L. Cardoso, N. C. Leite, and G. F. Salles, "Importance of Hematological Parameters for Micro- and Macrovascular Outcomes in Patients With Type 2 Diabetes: The Rio De Janeiro Type 2 Diabetes Cohort Study," *Cardiovascular Diabetology* 20, no. 1 (2021): 133.
54. R. Samidurai, S. varalakshmi Venkatesan, C. B. Kataria, G. Mano, and M. Sundaramalingham, "Complete Blood Count and Cardiovascular Risk Markers in Type 2 Diabetes Mellitus—Findings From a Cross-Sectional Study in South India," *Romanian Journal of Diabetes Nutrition and Metabolic Diseases* 28, no. 1 (2021): 18–26.
55. T. Adane, Z. Getaneh, and F. Asrie, "Red Blood Cell Parameters and Their Correlation With Renal Function Tests Among Diabetes Mellitus Patients: A Comparative Cross-Sectional Study," *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy* 13 (2020): 3937–3946.
56. E. Onalan, N. Gozel, and E. Donder, "Can Hematological Parameters in Type 2 Diabetes Predict Microvascular Complication Development?," *Pakistan Journal of Medical Sciences* 35, no. 6 (2019): 1511–1515.

57. A. Xanthopoulos, G. Giamouzis, A. Melidonis, et al., "Red Blood Cell Distribution Width as a Prognostic Marker in Patients With Heart Failure and Diabetes Mellitus," *Cardiovascular Diabetology* 16, no. 1 (2017): 81.
58. M. Verdoia, A. Schaffer, L. Barbieri, et al., "Diabetes, Glucose Control and Mean Platelet Volume: A Single-Centre Cohort Study," *Diabetes Research and Clinical Practice* 104, no. 2 (2014): 288–294.
59. J. D. McFadyen and Z. S. Kaplan, "Platelets Are Not Just for Clots," *Transfusion Medicine Reviews* 29, no. 2 (2015): 110–119.
60. N. Kakouros, J. J. Rade, A. Kourliouros, and J. R. Resar, "Platelet Function in Patients With Diabetes Mellitus: From a Theoretical to a Practical Perspective," *International Journal of Endocrinology* 2011 (2011): 1–14.
61. A. D. Michelson, "Methods for the Measurement of Platelet Function," *American Journal of Cardiology* 103, no. 3 (2009): 20A–26AA.
62. J. A. Colwell and R. W. Nesto, "The Platelet in Diabetes," *Diabetes Care* 26, no. 7 (2003): 2181–2188.
63. N. S. Nicholson, S. G. Panzer-Knodle, N. F. Haas, et al., "Assessment of Platelet Function Assays," *American Heart Journal* 135, no. 5 (1998): S170–S178.
64. S. G. Chu, R. C. Becker, P. B. Berger, et al., "Mean Platelet Volume as a Predictor of Cardiovascular Risk: A Systematic Review and Meta-Analysis," *Journal of Thrombosis and Haemostasis* 8, no. 1 (2010): 148–156.
65. M. Yilmaz, I. B. Delibas, U. Isaoglu, M. Ingeç, B. Borekci, and P. Ulug, "Relationship Between Mean Platelet Volume and Recurrent Miscarriage: A Preliminary Study," *Archives of Medical Science: AMS* 11, no. 5 (2015): 989–993.
66. H. Demirin, H. Ozhan, T. Uçgun, et al., "Normal Range of Mean Platelet Volume in Healthy Subjects: Insight From a Large Epidemiologic Study," *Thrombosis Research* 128, no. 4 (2011): 358–360.
67. E. Coban, F. Bostan, and M. Ozdogan, "The Mean Platelet Volume in Subjects With Impaired Fasting Glucose," *Platelets* 17, no. 1 (2006): 67–69.
68. E. Coban and B. Afacan, "The Effect of Rosuvastatin Treatment on the Mean Platelet Volume in Patients With Uncontrolled Primary Dyslipidemia With Hypolipidemic Diet Treatment," *Platelets* 19, no. 2 (2008): 111–114.
69. R. Demirtunc, D. Duman, M. Basar, M. Bilgi, M. Teomete, and T. Garip, "The Relationship Between Glycemic Control and Platelet Activity in Type 2 Diabetes Mellitus," *Journal of Diabetes and Its Complications* 23, no. 2 (2009): 89–94.
70. B. Shah, D. Sha, D. Xie, E. R. Mohler, and J. S. Berger, "The Relationship Between Diabetes, Metabolic Syndrome, and Platelet Activity as Measured by Mean Platelet Volume," *Diabetes Care* 35, no. 5 (2012): 1074–1078.
71. E. S. Kim, E. Y. Mo, S. D. Moon, and J. H. Han, "Mean Platelet Volume Is Closely Associated With Serum Glucose Level But Not With Arterial Stiffness and Carotid Atherosclerosis in Patients With Type 2 Diabetes," *Journal of Clinical Endocrinology and Metabolism* 100, no. 9 (2015): 3502–3508.
72. Q. Ding, F. Wang, X. Guo, and M. Liang, "The Relationship Between Mean Platelet Volume and Metabolic Syndrome in Patients With Type 2 Diabetes Mellitus: A Retrospective Study," *Medicine* 100, no. 13 (2021): e25303.
73. A. Ozder and H. H. Eker, "Investigation of Mean Platelet Volume in Patients With Type 2 Diabetes Mellitus and in Subjects With Impaired Fasting Glucose: A Cost-Effective Tool in Primary Health Care?," *International Journal of Clinical and Experimental Medicine* 7, no. 8 (2014): 2292–2297.
74. S. Li, C. Wang, X. W. Zhong, H. Q. Li, X. Q. Fu, and X. W. Ran, "[Variance of Mean Platelet Volume in Subjects With Normal Glucose Tolerance, Impaired Glucose Regulation and Type 2 Diabetic Mellitus and Its Relationship With Diabetic Peripheral Artery Disease]," *Zhonghua Yi Xue Za Zhi* 92, no. 4 (2012): 232–235.
75. H. Ebrahim, F. Asrie, and Z. Getaneh, "Basic Coagulation Profiles and Platelet Parameters Among Adult Type 1 and Type 2 Diabetes Patients at Dessie Referral Hospital, Northeast Ethiopia: Comparative Cross-Sectional Study," *Journal of Blood Medicine* 12 (2021): 33–42.
76. J. L. Ferreira, J. A. Gómez-Hospital, and D. J. Angiolillo, "Review Article: Platelet Abnormalities in Diabetes Mellitus," *Diabetes and Vascular Disease Research* 7, no. 4 (2010): 251–259.
77. D. J. Schneider, "Factors Contributing to Increased Platelet Reactivity in People With Diabetes," *Diabetes Care* 32, no. 4 (2009): 525–527.
78. A. I. Vinik, T. Erbas, T. S. Park, R. Nolan, and G. L. Pittenger, "Platelet Dysfunction in Type 2 Diabetes," *Diabetes Care* 24, no. 8 (2001): 1476–1485.
79. J. A. Dormandy, B. Charbonnel, D. J. Eckland, et al., "Secondary Prevention of Macrovascular Events in Patients With Type 2 Diabetes in the Proactive Study (Prospective Pioglitazone Clinical Trial In Macrovascular Events): A Randomised Controlled Trial," *Lancet* 366, no. 9493 (2005): 1279–1289.
80. W. H. Tang, J. Stitham, S. Gleim, et al., "Glucose and Collagen Regulate Human Platelet Activity Through Aldose Reductase Induction of Thromboxane," *Journal of Clinical Investigation* 121, no. 11 (2011): 4462–4476.
81. G. Davì, I. Catalano, M. Averna, et al., "Thromboxane Biosynthesis and Platelet Function in Type II Diabetes Mellitus," *New England Journal of Medicine* 322, no. 25 (1990): 1769–1774.
82. Y. Zhang and H. Zhou, "Hyper-Reactive Platelets and Type 2 Diabetes," *Zhong Nan Da Xue Xue Bao Yi Xue Ban = Journal of Central South University Medical Sciences* 47, no. 3 (2022): 374–383.
83. S. Jindal, S. Gupta, R. Gupta, et al., "Platelet Indices in Diabetes Mellitus: Indicators of Diabetic Microvascular Complications," *Hematology* 16, no. 2 (2011): 86–89.
84. A. Shiny, Y. S. Bibin, C. S. Shanthirani, et al., "Association of Neutrophil-Lymphocyte Ratio With Glucose Intolerance: An Indicator of Systemic Inflammation in Patients With Type 2 Diabetes," *Diabetes Technology & Therapeutics* 16, no. 8 (2014): 524–530.
85. C. Pitsavos, M. Tampourlou, D. B. Panagiotakos, et al., "Association Between Low-Grade Systemic Inflammation and Type 2 Diabetes Mellitus Among Men and Women From the Attica Study," *Review of Diabetic Studies* 4, no. 2 (2007): 98–104.
86. T. Fujita, S. Hemmi, M. Kajiwar, et al., "Complement-Mediated Chronic Inflammation Is Associated With Diabetic Microvascular Complication," *Diabetes/Metabolism Research and Reviews* 29, no. 3 (2013): 220–226.
87. H. Yilmaz, B. Ucan, M. Sayki, et al., "Usefulness of the Neutrophil-to-Lymphocyte Ratio to Prediction of Type 2 Diabetes Mellitus in Morbid Obesity," *Diabetes & Metabolic Syndrome* 9, no. 4 (2015): 299–304.
88. F. Intiaz, K. Shafique, S. Mirza, Z. Ayoob, P. Vart, and S. Rao, "Neutrophil Lymphocyte Ratio as a Measure of Systemic Inflammation in Prevalent Chronic Diseases in Asian Population," *International Archives of Medicine* 5, no. 1 (2012): 2.
89. E. Buyukkaya, M. F. Karakaş, E. Karakaş, et al., "Correlation of Neutrophil to Lymphocyte Ratio With the Presence and Severity of Metabolic Syndrome," *Clinical and Applied Thrombosis/Hemostasis* 20, no. 2 (2014): 159–163.
90. B. Azab, V. Chainani, N. Shah, and J. T. McGinn, "Neutrophil-Lymphocyte Ratio as a Predictor of Major Adverse Cardiac Events Among Diabetic Population: A 4-Year Follow-Up Study," *Angiology* 64, no. 6 (2013): 456–465.

91. G.-K. Lee, L.-C. Lee, E. Chong, et al., "The Long-Term Predictive Value of the Neutrophil-to-Lymphocyte Ratio in Type 2 Diabetic Patients Presenting With Acute Myocardial Infarction," *QJM: Monthly Journal of the Association of Physicians* 105, no. 11 (2012): 1075–1082.
92. M. Hussain, M. Z. M. Babar, L. Akhtar, and M. S. Hussain, "Neutrophil Lymphocyte Ratio (NLR): A Well Assessment Tool of Glycemic Control in Type 2 Diabetic Patients," *Pakistan Journal of Medical Sciences* 33, no. 6 (2017): 1366.
93. Z. A. Öztürk, M. E. Kuyumcu, Y. Yesil, et al., "Is There a Link Between Neutrophil-Lymphocyte Ratio and Microvascular Complications in Geriatric Diabetic Patients?," *Journal of Endocrinological Investigation* 36 (2013): 593–599.
94. P. H. Gibson, B. L. Croal, B. H. Cuthbertson, et al., "Preoperative Neutrophil-Lymphocyte Ratio and Outcome From Coronary Artery Bypass Grafting," *American Heart Journal* 154, no. 5 (2007): 995–1002.
95. J. Núñez, E. Núñez, V. Bodí, et al., "Usefulness of the Neutrophil to Lymphocyte Ratio in Predicting Long-Term Mortality in ST Segment Elevation Myocardial Infarction," *American Journal of Cardiology* 101, no. 6 (2008): 747–752.
96. B. Azab, J. Daoud, F. B. Naeem, et al., "Neutrophil-To-Lymphocyte Ratio as a Predictor of Worsening Renal Function in Diabetic Patients (3-Year Follow-Up Study)," *Renal Failure* 34, no. 5 (2012): 571–576.
97. F.-Y. Chang and M.-F. Shaio, "Decreased Cell-Mediated Immunity in Patients With Non-Insulin-Dependent Diabetes Mellitus," *Diabetes Research and Clinical Practice* 28, no. 2 (1995): 137–146.
98. Z. Eftekhari, M. Aghaei, and N. Saki, "DNA Damage Repair in Megakaryopoiesis: Molecular and Clinical Aspects," *Expert Review of Hematology* 17, no. 10 (2024): 705–712, <https://doi.org/10.1080/17474086.2024.2391102>.
99. A. K. Mohamed, A. Bierhaus, S. Schiekofer, H. Tritschler, R. Ziegler, and P. P. Nawroth, "The Role of Oxidative Stress and NF- κ B Activation in Late Diabetic Complications," *Biofactors* 10, no. 2–3 (1999): 157–167.
100. G. Caramori, M. Romagnoli, and P. Casolari, "Nuclear Localisation of p65 in Sputum Macrophages but Not in Sputum Neutrophils During COPD Exacerbations," *Thorax* 58, no. 4 (2003): 348–351.
101. B. K. Duffy, H. S. Gurm, V. Rajagopal, R. Gupta, S. G. Ellis, and D. L. Bhatt, "Usefulness of an Elevated Neutrophil to Lymphocyte Ratio in Predicting Long-Term Mortality After Percutaneous Coronary Intervention," *American Journal of Cardiology* 97, no. 7 (2006): 993–996.
102. K. Ohshita, K. Yamane, M. Hanafusa, et al., "Elevated White Blood Cell Count in Subjects With Impaired Glucose Tolerance," *Diabetes Care* 27, no. 2 (2004): 491–496.
103. S. R. De Rooij, G. Nijpels, P. M. Nilsson, et al., "Low-Grade Chronic Inflammation in the Relationship Between Insulin Sensitivity and Cardiovascular Disease (RISC) Population," *Diabetes Care* 32, no. 7 (2009): 1295–1301.
104. P. C. Tong, K.-F. Lee, W.-Y. So, et al., "White Blood Cell Count Is Associated With Macro- and Microvascular Complications in Chinese Patients With Type 2 Diabetes," *Diabetes Care* 27, no. 1 (2004): 216–222.
105. J. C. R. Tsai, S. H. Sheu, H. C. Chiu, et al., "Association of Peripheral Total and Differential Leukocyte Counts With Metabolic Syndrome and Risk of Ischemic Cardiovascular Diseases in Patients With Type 2 Diabetes Mellitus," *Diabetes/Metabolism Research and Reviews* 23, no. 2 (2007): 111–118.
106. E. Ernst, D. E. Hammerschmidt, U. Bagge, A. Matrai, and J. A. Dormandy, "Leukocytes and the Risk of Ischemic Diseases," *Journal of the American Medical Association* 257, no. 17 (1987): 2318–2324.
107. D. W. Brown, W. H. Giles, and J. B. Croft, "White Blood Cell Count," *Journal of Clinical Epidemiology* 54, no. 3 (2001): 316–322.
108. M. S. Sabatine, D. A. Morrow, C. P. Cannon, et al., "Relationship Between Baseline White Blood Cell Count and Degree of Coronary Artery Disease and Mortality in Patients With Acute Coronary Syndromes," *Journal of the American College of Cardiology* 40, no. 10 (2002): 1761–1768.
109. Y. Liu, X. Lai, W. Guo, et al., "Total White Blood Cell Count Mediated the Association Between Increased Arterial Stiffness and Risk of Type 2 Diabetes Mellitus in Chinese Adults," *Arteriosclerosis, Thrombosis, and Vascular Biology* 40, no. 4 (2020): 1009–1015.
110. E. M. Akbas, L. Demirtas, A. Ozcicek, et al., "Association of Epicardial Adipose Tissue, Neutrophil-to-Lymphocyte Ratio and Platelet-To-Lymphocyte Ratio With Diabetic Nephropathy," *International Journal of Clinical and Experimental Medicine* 7, no. 7 (2014): 1794–1801.
111. B. Hudzik, J. Szkodziński, A. Lekston, M. Gierlotka, L. Polonński, and M. Gąsior, "Mean Platelet Volume-to-Lymphocyte Ratio: A Novel Marker of Poor Short- and Long-Term Prognosis in Patients With Diabetes Mellitus and Acute Myocardial Infarction," *Journal of Diabetes and Its Complications* 30, no. 6 (2016): 1097–1102.
112. Y. Le Tulzo, C. Pangault, A. Gacouin, et al., "Early Circulating Lymphocyte Apoptosis in Human Septic Shock Is Associated With Poor Outcome," *Shock* 18, no. 6 (2002): 487–494.
113. X. An, P.-R. Ding, Y.-H. Li, et al., "Elevated Neutrophil to Lymphocyte Ratio Predicts Survival in Advanced Pancreatic Cancer," *Biomarkers* 15, no. 6 (2010): 516–522.
114. L. Demirtas, H. Degirmenci, E. M. Akbas, et al., "Association of Hematological Indices With Diabetes, Impaired Glucose Regulation and Microvascular Complications of Diabetes," *International Journal of Clinical and Experimental Medicine* 8, no. 7 (2015): 11420–11427.
115. S. Bilgin, G. Aktas, G. Kahveci, B. M. Atak, O. Kurtkulagi, and T. T. Duman, "Does Mean Platelet Volume/Lymphocyte Count Ratio Associate With Frailty in Type 2 Diabetes Mellitus?," *Bratislava Medical Journal* 122 (2021): 116–119.
116. M. Yilmaz and H. Kayaçikçik, "A New Inflammatory Marker: Elevated Monocyte to HDL Cholesterol Ratio Associated With Smoking," *Journal of Clinical Medicine* 7, no. 4 (2018): 76.
117. M. Brownlee, "The Pathobiology of Diabetic Complications," *Diabetes* 54, no. 6 (2005): 1615–1625.
118. W. B. Kannel, K. Anderson, and P. W. Wilson, "White Blood Cell Count and Cardiovascular Disease: Insights From the Framingham Study," *Journal of the American Medical Association* 267, no. 9 (1992): 1253–1256.
119. F. Cavalot, P. Massucco, P. Perna, M. Traversa, G. Anfossi, and M. Trovati, "White Blood Cell Count Is Positively Correlated With Albumin Excretion Rate in Subjects With Type 2 Diabetes," *Diabetes Care* 25, no. 12 (2002): 2354–2355.
120. P. J. Barter, S. Nicholls, K.-A. Rye, G. M. Anantharamaiah, M. Navab, and A. M. Fogelman, "Antiinflammatory Properties of HDL," *Circulation Research* 95, no. 8 (2004): 764–772.
121. G. W. Cockerill, K.-A. Rye, J. R. Gamble, M. A. Vadas, and P. J. Barter, "High-Density Lipoproteins Inhibit Cytokine-Induced Expression of Endothelial Cell Adhesion Molecules," *Arteriosclerosis, Thrombosis, and Vascular Biology* 15, no. 11 (1995): 1987–1994.
122. A. J. Murphy, K. J. Woollard, A. Hoang, et al., "High-Density Lipoprotein Reduces the Human Monocyte Inflammatory Response," *Arteriosclerosis, Thrombosis, and Vascular Biology* 28, no. 11 (2008): 2071–2077.
123. M. Kanbay, Y. Solak, H. U. Unal, et al., "Monocyte Count/HDL Cholesterol Ratio and Cardiovascular Events in Patients With Chronic Kidney Disease," *International Urology and Nephrology* 46 (2014): 1619–1625.
124. C. Nguyen, C. L. Crowe, E. Kuti, et al., "Impact of Pre-Existing Type 2 Diabetes Mellitus and Cardiovascular Disease on Healthcare Resource Utilization and Costs in Patients With COVID-19," *Journal of Health Economics and Outcomes Research* 11, no. 1 (2024): 112.

125. L. Zhu, Z. G. She, X. Cheng, et al., "Association of Blood Glucose Control and Outcomes in Patients With COVID-19 and Pre-Existing Type 2 Diabetes," *Cell Metabolism* 31, no. 6 (2020): 1068–1077.e3.
126. F. Liu, Y. Li, W. Li, et al., "The Role of Peripheral White Blood Cell Counts in the Association Between Central Adiposity and Glycemic Status," *Nutrition & Diabetes* 14, no. 1 (2024): 30.
127. M.-N. Bastani, S. Jalilian, S. S. Bahreiny, et al., "Update Prognostic Potency of Surfactant Protein D (SP-D) in the COVID-19 Landscape: An In-Depth Meta-Analytical Exploration," *Biomarkers in Medicine* 18, no. 24 (2024): 1135–1148.
128. A. T. Soliman, V. De Sanctis, M. Yassin, and N. Soliman, "Iron Deficiency Anemia and Glucose Metabolism," *Acta Bio-Medica: Atenei Parmensis* 88, no. 1 (2017): 112–118.
129. W. Xu, H. Wu, S. Ma, et al., "Correlation Between Peripheral White Blood Cell Counts and Hyperglycemic Emergencies," *International Journal of Medical Sciences* 10, no. 6 (2013): 758–765.
130. M. Arkew, T. Yemane, Y. Mengistu, K. Gemechu, and G. Tesfaye, "Hematological Parameters of Type 2 Diabetic Adult Patients at Debre Berhan Referral Hospital, Northeast Ethiopia: A Comparative Cross-Sectional Study," *PLoS One* 16, no. 6 (2021): e0253286.
131. E. Sayedali, A. E. Yalin, and S. Yalin, "Association Between Metformin and Vitamin B12 Deficiency in Patients With Type 2 Diabetes," *World Journal of Diabetes* 14, no. 5 (2023): 585–593.
132. A. Frenkel, E. Kachko, V. Novack, et al., "The Association of Glucocorticosteroid Treatment With WBC Count in Patients With COPD Exacerbation," *Journal of Clinical Medicine* 8, no. 10 (2019): 1697.
133. E. H. Nah, S. Cho, H. Park, S. Kim, and H. I. Cho, "Associations of Complete Blood Count Parameters With Pancreatic Beta-Cell Function and Insulin Resistance in Prediabetes and Type 2 Diabetes Mellitus," *Journal of Clinical Laboratory Analysis* 36, no. 6 (2022): e24454.
134. K. Essawi, G. Dobie, M. Shaabi, et al., "Comparative Analysis of Red Blood Cells, White Blood Cells, Platelet Count, and Indices in Type 2 Diabetes Mellitus Patients and Normal Controls: Association and Clinical Implications," *Diabetes, Metabolic Syndrome and Obesity* 16 (2023): 3123–3132.
135. H. Ozisik, S. Cetinkalp, A. Suner, G. Ozgen, F. Saygili, and M. Erdogan, "The Relationship Between Neutrophil Lymphocyte Ratio and Diabetes Control in Patients With Type 2 Diabetes Mellitus," *Ege Tip Dergisi* 61, no. 3 (2022): 444–451.
136. A. A. Kenyang, N. Juhan, Y. Z. Zubairi, N. Azizan, and C. M. Ho, "Predictive Factors for Diabetes Mellitus: Insights From Complete Blood Count Analysis," *Sains Malaysiana* 53, no. 10 (2024): 3535–3543.
137. K. Aygün, A. Asma Sakalli, H. S. Küçükerdem, O. Aygün, and Ö. Gökdemir, "Assessment of Neutrophil/Lymphocyte Ratio and Mean Platelet Volume Values in Patients With Diabetes Mellitus Diagnosis: A Case-Control Study," *Medicine* 103, no. 37 (2024): e39661.
138. Y. Sertbas, M. Sertbas, N. Okuroglu, M. A. Ozturk, K. Y. Abacar, and A. Ozdemir, "Mean Platelet Volume Changes Before and After Glycated Hemoglobin (HbA 1c) Improvement in a Large Study Population," *Archives of Medical Science* 4, no. 4 (2017): 711–715.
139. A. Swaminathan, K. Amitkumar, S. Ganapathy, and S. Ayyavoo, "Evaluation of Mean Platelet Volume and Other Platelet Parameters in Subjects With Type-2 Diabetes Mellitus," *National Journal of Physiology, Pharmacy and Pharmacology* 7, no. 1 (2017): 51.
140. V. Cassano, G. Armentaro, D. Iembo, et al., "Mean Platelet Volume (MPV) as New Marker of Diabetic Macrovascular Complications in Patients With Different Glucose Homeostasis: Platelets in Cardiovascular Risk," *Cardiovascular Diabetology* 23, no. 1 (2024): 89.
141. D. Milosevic and V. L. Panin, "Relationship Between Hematological Parameters and Glycemic Control in Type 2 Diabetes Mellitus Patients," *Journal of Medical Biochemistry* 38, no. 2 (2019): 164–171.
142. M. Taimah, N. F. Juber, P. Holland, and H. Brown, "A Systematic Review of the Methodology for Examining the Relationship Between Obstructive Sleep Apnea and Type Two Diabetes Mellitus," *Frontiers in Endocrinology* 15 (2024): 288–294, <https://doi.org/10.1016/j.diabres.2013.12.020>.