

```

import numpy as np

#array creation
array_1 = np.array([1,2,3,4,5])
array_2 = np.array([6,7,8,9,10])
print(array_1)
data = [[1,2,3] , [4,5,6]]
array_3 = np.array(data)
print(array_3)

[1 2 3 4 5]
[[1 2 3]
 [4 5 6]]

#universal funcction - add , mean , std
array_add = np.add(array_1,array_2)
print(array_add)
array1_std = np.std(array_1)
print(array1_std)

[ 7  9 11 13 15]
1.4142135623730951

#indexing and slicing
list = [1,2,3,4,5,6]
print(list)
print(list[3:])
print(list[:-4])
print(list[1:2])

[1, 2, 3, 4, 5, 6]
[4, 5, 6]
[1, 2]
[2, 4, 6]

#broadcasting
b_1 = np.array([[1,2],[3,4]])
b_2 = np.array([10,20])
b_3 = b_1 + b_2
b_3

array([[11, 22],
       [13, 24]])

#linear algebra - np.linalg.det , np.linalg.inv
l_1 = np.array([[1,2],[6,7]])
np.linalg.det(l_1)
np.linalg.inv(l_1)

array([[-1.4,  0.4],
       [ 1.2, -0.2]])

```

```
import random  
number = np.random.randint(0,100)  
number
```

43