

# Assignment 2 Group 26

Vanshita Sharma Kumar

Lucian Trușcă

Xander Poortvliet

2025-10-08

## Excercise 2.1

We first will start with the full multi-regression model

```
sat = read.delim("sat.txt", header=TRUE, sep = ",", stringsAsFactors = FALSE)
```

```
##
## Call:
## lm(formula = total ~ expend + ratio + salary + takers, data = sat)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -90.531 -20.855  -1.746   15.979   66.571
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 1045.9715    52.8698   19.784 < 2e-16 ***
## expend         4.4626    10.5465    0.423  0.674
## ratio        -3.6242     3.2154   -1.127  0.266
## salary         1.6379     2.3872    0.686  0.496
## takers        -2.9045     0.2313  -12.559 2.61e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 32.7 on 45 degrees of freedom
## Multiple R-squared:  0.8246, Adjusted R-squared:  0.809
## F-statistic: 52.88 on 4 and 45 DF,  p-value: < 2.2e-16

## [1] 497.3694
```