

# Lesson 05 Multiple LR. Intro to Logistic Regression

Lusine Zilfimian

March 23 (Monday), 2020

# Contents

- Binominal Logistic Regression

# Contents

- Binominal Logistic Regression
- Multinomial Logistic Regression

# Contents

- Binominal Logistic Regression
- Multinomial Logistic Regression
- Interpretation

# Contents

- Binominal Logistic Regression
- Multinomial Logistic Regression
- Interpretation
- Coding examples in R. Intro to Lab 06.

# Last Lecture ReCap

- Why Not Linear Regression and OLS?

# Last Lecture ReCap

- Why Not Linear Regression and OLS?
- What is the difference between binomial and multinomial LogReg?

# Last Lecture ReCap

- Why Not Linear Regression and OLS?
- What is the difference between binomial and multinomial LogReg?
- Which type of dependent/independent variable is used in LogReg?



## Binary simple case

- Binary response using one predictor


## Binary simple case

- Binary response using one predictor
- When  $p = 2$ , there is only a single linear function to estimate.

## Binary simple case

- Binary response using one predictor
- When  $p = 2$ , there is only a single linear function to estimate.
- The probability:  $\mathbb{P}(y_i = 1) = \frac{e^{\beta_0 + \beta_1 x_i}}{1 + e^{\beta_0 + \beta_1 x_i}}$

## S-shaped curve



## Terminology

- Odds:  $\frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = e^{\beta_0 + \beta_1 x} \in [0; +\infty)$

## Terminology

- Odds:  $\frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = e^{\beta_0 + \beta_1 x} \in [0; +\infty)$
- Log-odds or logit:  $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x \in (-\infty; +\infty)$

## Terminology

- Odds:  $\frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = e^{\beta_0 + \beta_1 x} \in [0; +\infty)$
- Log-odds or logit:  $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x \in (-\infty; +\infty)$
- Odds ratio:  $\frac{\frac{\mathbb{P}(y=1|x=1)}{1-\mathbb{P}(y=1|x=1)}}{\frac{\mathbb{P}(y=1|x=0)}{1-\mathbb{P}(y=1|x=0)}}$

## Estimating the Coefficients

- The most common approach – maximum likelihood: choose that value of parameter, under which it is most likely to get the (your) data.



## Estimating the Coefficients

- The most common approach – maximum likelihood: choose that value of parameter, under which it is most likely to get the (your) data.
- In our case, find/choose **estimates** for coefficients such that the **predicted probability** of  $Y$  corresponds as closely as possible to  $Y$ .

## Estimating the Coefficients

- The most common approach – maximum likelihood: choose that value of parameter, under which it is most likely to get the (your) data.
- In our case, find/choose **estimates** for coefficients such that the **predicted probability** of  $Y$  corresponds as closely as possible to  $Y$ .
- Likelihood Function is the joint PD(M)F, considered as a function of the parameter

## Estimating the Coefficients

- The most common approach – maximum likelihood: choose that value of parameter, under which it is most likely to get the (your) data.
- In our case, find/choose **estimates** for coefficients such that the **predicted probability** of  $Y$  corresponds as closely as possible to  $Y$ .
- Likelihood Function is the joint PD(M)F, considered as a function of the parameter
- Likelihood function:

## Estimating the Coefficients

- The most common approach – maximum likelihood: choose that value of parameter, under which it is most likely to get the (your) data.
- In our case, find/choose **estimates** for coefficients such that the **predicted probability** of Y corresponds as closely as possible to Y.
- Likelihood Function is the joint PD(M)F, considered as a function of the parameter
- Likelihood function:
- 

$$L(\beta_0, \beta_1) = \prod_{y_i=1} F(\beta_0 + \beta_1 x_i) \prod_{y_i=0} (1 - F(\beta_0 + \beta_1 x_i)) \rightarrow \max$$

## Estimating the Coefficients

- The most common approach – maximum likelihood: choose that value of parameter, under which it is most likely to get the (your) data.
- In our case, find/choose **estimates** for coefficients such that the **predicted probability** of Y corresponds as closely as possible to Y.
- Likelihood Function is the joint PDF, considered as a function of the parameter
- Likelihood function:
- 

$$L(\beta_0, \beta_1) = \prod_{y_i=1} F(\beta_0 + \beta_1 x_i) \prod_{y_i=0} (1 - F(\beta_0 + \beta_1 x_i)) \rightarrow \max$$

- Likelihood is not a Probability - it can be larger than 1. It is not a PDF either, it is a function of the parameter.

## Estimating the Coefficients

- The Maximum Likelihood Method suggests to find a point that makes our Likelihood Maximal:

## Estimating the Coefficients

- The Maximum Likelihood Method suggests to find a point that makes our Likelihood Maximal:
- Log-Likelihood function:

## Estimating the Coefficients

- The Maximum Likelihood Method suggests to find a point that makes our Likelihood Maximal:
- Log-Likelihood function:
- $l(\beta_0, \beta_1) = \sum_{i=1}^N (y_i \ln \mathbb{P}(x_i) + (1 - y_i) \ln(1 - \mathbb{P}(x_i)))$



## Estimating the Coefficients

- The Maximum Likelihood Method suggests to find a point that makes our Likelihood Maximal:
- Log-Likelihood function:
- $l(\beta_0, \beta_1) = \sum_{i=1}^N (y_i \ln \mathbb{P}(x_i) + (1 - y_i) \ln(1 - \mathbb{P}(x_i)))$
- The points of maximum of  $L(\beta_0, \beta_1)$  and  $\ln L(\beta_0, \beta_1)$  coincide:

## Estimating the Coefficients

- The Maximum Likelihood Method suggests to find a point that makes our Likelihood Maximal:
- Log-Likelihood function:
- $l(\beta_0, \beta_1) = \sum_{i=1}^N (y_i \ln \mathbb{P}(x_i) + (1 - y_i) \ln(1 - \mathbb{P}(x_i)))$
- The points of maximum of  $L(\beta_0, \beta_1)$  and  $\ln L(\beta_0, \beta_1)$  coincide:
- $\underset{\beta}{\operatorname{argmax}} L(\beta_0, \beta_1) = \underset{\beta}{\operatorname{argmax}} \ln L(\beta_0, \beta_1)$

## Estimating the Coefficients

- The Maximum Likelihood Method suggests to find a point that makes our Likelihood Maximal:
- Log-Likelihood function:
- $l(\beta_0, \beta_1) = \sum_{i=1}^N (y_i \ln \mathbb{P}(x_i) + (1 - y_i) \ln(1 - \mathbb{P}(x_i)))$
- The points of maximum of  $L(\beta_0, \beta_1)$  and  $\ln L(\beta_0, \beta_1)$  coincide:
- $\underset{\beta}{\operatorname{argmax}} L(\beta_0, \beta_1) = \underset{\beta}{\operatorname{argmax}} \ln L(\beta_0, \beta_1)$
- The estimates for coefficients are calculated using iterative procedure.

## Binomial, Multiple logistic regression

- Binary response using multiple predictors

## Binomial, Multiple logistic regression

- Binary response using multiple predictors
- Model specification:

## Binomial, Multiple logistic regression

- Binary response using multiple predictors
- Model specification:
- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k$

## Binomial, Multiple logistic regression

- Binary response using multiple predictors
- Model specification:
- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k$
- $\mathbb{P}(y = 1) = \frac{e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}{1 + e^{\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k}}$

# Interpretations (Simple LogReg case)

## Numeric predictors (Simple LogReg case)

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x$ , where  $X$  is numeric

The rate of change in  $\mathbb{P}(x)$  per unit change in  $X$  depends on the current value of  $X$ .



# Interpretations (Simple LogReg case)

## Numeric predictors (Simple LogReg case)

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x$ , where  $X$  is numeric
- Increasing  $X$  by one unit changes the log odds by  $\beta_1$ , or equivalently it multiplies the odds by  $e^{\beta_1}$ .

The rate of change in  $\mathbb{P}(x)$  per unit change in  $X$  depends on the current value of  $X$ .

# Interpretations (Simple LogReg case)

## Numeric predictors (Simple LogReg case)

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x$ , where  $X$  is numeric
- Increasing  $X$  by one unit changes the log odds by  $\beta_1$ , or equivalently it multiplies the odds by  $e^{\beta_1}$ .
- If  $\beta_1 > 0$ ,  $\uparrow x \Rightarrow \uparrow \mathbb{P}(x)$

The rate of change in  $\mathbb{P}(x)$  per unit change in  $X$  depends on the current value of  $X$ .


# Interpretations (Simple LogReg case)

## Numeric predictors (Simple LogReg case)

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 x$ , where  $X$  is numeric
- Increasing  $X$  by one unit changes the log odds by  $\beta_1$ , or equivalently it multiplies the odds by  $e^{\beta_1}$ .
- If  $\beta_1 > 0$ ,  $\uparrow x \Rightarrow \uparrow \mathbb{P}(x)$
- If  $\beta_1 < 0$ ,  $\uparrow x \Rightarrow \downarrow \mathbb{P}(x)$

The rate of change in  $\mathbb{P}(x)$  per unit change in  $X$  depends on the current value of  $X$ .

## Numeric predictors (Simple LogReg case)



$$F(\beta_0 + \beta_1 x) = \frac{1}{1 + e^{-(\beta_0 + \beta_1 x)}} = \frac{1}{1 + e^0} = 0.5$$

$$\beta_0 + \beta_1 x = 0 \Rightarrow x = -\beta_0 / \beta_1$$

Marginal effect:

$$\frac{\partial \Pr(y=1)}{\partial x_j} = F(\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k) * \beta_j$$

## Categorical predictors

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 X,$

## Categorical predictors

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 X,$
- $X$  is binary variable

## Categorical predictors

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 X,$
- $X$  is binary variable
- $\ln \frac{\mathbb{P}(y=1|x=1)}{1-\mathbb{P}(y=1|x=1)} - \ln \frac{\mathbb{P}(y=1|x=0)}{1-\mathbb{P}(y=1|x=0)} = \beta_1$

## Categorical predictors

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 X,$
- $X$  is binary variable
- $\ln \frac{\mathbb{P}(y=1|x=1)}{1-\mathbb{P}(y=1|x=1)} - \ln \frac{\mathbb{P}(y=1|x=0)}{1-\mathbb{P}(y=1|x=0)} = \beta_1$
- $e^{\beta_1} = \frac{\frac{\mathbb{P}(y=1|x=1)}{1-\mathbb{P}(y=1|x=1)}}{\frac{\mathbb{P}(y=1|x=0)}{1-\mathbb{P}(y=1|x=0)}} = \text{Odds ratio}$



## Categorical predictors

- $\ln \frac{\mathbb{P}(y=1)}{1-\mathbb{P}(y=1)} = \beta_0 + \beta_1 X,$
- $X$  is binary variable
- $\ln \frac{\mathbb{P}(y=1|x=1)}{1-\mathbb{P}(y=1|x=1)} - \ln \frac{\mathbb{P}(y=1|x=0)}{1-\mathbb{P}(y=1|x=0)} = \beta_1$
- $e^{\beta_1} = \frac{\frac{\mathbb{P}(y=1|x=1)}{1-\mathbb{P}(y=1|x=1)}}{\frac{\mathbb{P}(y=1|x=0)}{1-\mathbb{P}(y=1|x=0)}} = \text{Odds ratio}$
- By changing  $x$  from 0 to 1, the odds ratio of  $y = 1$  will be changed by  $\frac{1}{e^{\beta_1}}$  times

# Multinomial logistic regression

- Response variable has more than two classes:

# Multinomial logistic regression

- Response variable has more than two classes:
- $p = 1, 2, \dots, P - 1$

# Multinomial logistic regression

- Response variable has more than two classes:
- $p = 1, 2, \dots, P - 1$
- $\ln \frac{\mathbb{P}(Y=1|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k$

# Multinomial logistic regression

- Response variable has more than two classes:
- $p = 1, 2, \dots, P - 1$
- $\ln \frac{\mathbb{P}(Y=1|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k$
- $\ln \frac{\mathbb{P}(Y=2|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k$

# Multinomial logistic regression

- Response variable has more than two classes:
- $p = 1, 2, \dots, P - 1$
- $\ln \frac{\mathbb{P}(Y=1|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k$
- $\ln \frac{\mathbb{P}(Y=2|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k$
- ...

# Multinomial logistic regression

- Response variable has more than two classes:
- $p = 1, 2, \dots, P - 1$
- $\ln \frac{\mathbb{P}(Y=1|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k$
- $\ln \frac{\mathbb{P}(Y=2|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k$
- $\dots$
- $\ln \frac{\mathbb{P}(Y=P-1|X=x)}{1-\mathbb{P}(Y=P|X=x)} = \beta_{(P-1)0} + \beta_{(P-1)1}X_1 + \dots + \beta_{(P-1)k}X_k$

## Multinomial logistic regression

- $\mathbb{P}(Y = 1|X = x) = \frac{e^{\beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$



## Multinomial logistic regression

- $\mathbb{P}(Y = 1|X = x) = \frac{e^{\beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- $\mathbb{P}(Y = 2|X = x) = \frac{e^{\beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$

## Multinomial logistic regression

- $\mathbb{P}(Y = 1|X = x) = \frac{e^{\beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- $\mathbb{P}(Y = 2|X = x) = \frac{e^{\beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- ...

## Multinomial logistic regression

- $\mathbb{P}(Y = 1|X = x) = \frac{e^{\beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- $\mathbb{P}(Y = 2|X = x) = \frac{e^{\beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- ...
- $\mathbb{P}(Y = P|X = x) = \frac{1}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$

## Multinomial logistic regression

- $\mathbb{P}(Y = 1|X = x) = \frac{e^{\beta_{10} + \beta_{11}X_1 + \dots + \beta_{1k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- $\mathbb{P}(Y = 2|X = x) = \frac{e^{\beta_{20} + \beta_{21}X_1 + \dots + \beta_{2k}X_k}}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- ...
- $\mathbb{P}(Y = P|X = x) = \frac{1}{1 + \sum_{l=1}^{P-1} e^{\beta_{l0} + \beta_{l1}X_1 + \dots + \beta_{lk}X_k}}$
- Derive these formulas for  $p=3$  case.

## Goodness of fit

- Confusion matrix

		Predicted	
		Negative (0)	Positive (1)
Actual	Negative (0)	TN	FP
	Positive (1)	FN	TP

## Goodness of fit

- Confusion matrix

		Predicted	
		Negative (0)	Positive (1)
Actual	Negative (0)	TN	FP
	Positive (1)	FN	TP

- $Accuracy = \frac{TP+TN}{Total}$

## Goodness of fit

- Confusion matrix

		Predicted	
		Negative (0)	Positive (1)
Actual	Negative (0)	TN	FP
	Positive (1)	FN	TP

- $Accuracy = \frac{TP+TN}{Total}$
- $Sensitivity = \frac{TP}{TP+FN}$

## Goodness of fit

- Confusion matrix

		Predicted	
		Negative (0)	Positive (1)
Actual	Negative (0)	TN	FP
	Positive (1)	FN	TP

- $Accuracy = \frac{TP+TN}{Total}$
- $Sensitivity = \frac{TP}{TP+FN}$
- $Specificity = \frac{TN}{TN+FP}$



## Goodness of fit

- How to choose the threshold?

Predicted	Actual1	Actual2	Actual3	Actual4	Actual5
0.1000000	0	1	0	0	0
0.1888889	0	1	0	0	1
0.2777778	0	1	0	0	0
0.3666667	0	1	0	0	1
0.4555556	0	1	0	1	0
0.5444444	0	1	1	1	1
0.6333333	0	1	1	1	0
0.7222222	0	1	1	1	1
0.8111111	0	1	1	1	0
0.9000000	0	1	1	1	1

## Goodness of fit

- How to choose the threshold?

Predicted	Actual1	Actual2	Actual3	Actual4	Actual5
0.1000000	0	1	0	0	0
0.1888889	0	1	0	0	1
0.2777778	0	1	0	0	0
0.3666667	0	1	0	0	1
0.4555556	0	1	0	1	0
0.5444444	0	1	1	1	1
0.6333333	0	1	1	1	0
0.7222222	0	1	1	1	1
0.8111111	0	1	1	1	0
0.9000000	0	1	1	1	1

- Which one is the worst case?

## Goodness of fit

- How to choose the threshold?

Predicted	Actual1	Actual2	Actual3	Actual4	Actual5
0.1000000	0	1	0	0	0
0.1888889	0	1	0	0	1
0.2777778	0	1	0	0	0
0.3666667	0	1	0	0	1
0.4555556	0	1	0	1	0
0.5444444	0	1	1	1	1
0.6333333	0	1	1	1	0
0.7222222	0	1	1	1	1
0.8111111	0	1	1	1	0
0.9000000	0	1	1	1	1

- Which one is the worst case?
- Maximum cutoff value  $\Rightarrow$  all records will be classified as 0.


## Goodness of fit

- How to choose the threshold?

Predicted	Actual1	Actual2	Actual3	Actual4	Actual5
0.1000000	0	1	0	0	0
0.1888889	0	1	0	0	1
0.2777778	0	1	0	0	0
0.3666667	0	1	0	0	1
0.4555556	0	1	0	1	0
0.5444444	0	1	1	1	1
0.6333333	0	1	1	1	0
0.7222222	0	1	1	1	1
0.8111111	0	1	1	1	0
0.9000000	0	1	1	1	1


- Which one is the worst case?
- Maximum cutoff value  $\Rightarrow$  all records will be classified as 0.
- Minimum cutoff value  $\Rightarrow$  all records will be classified as 1.

## Goodness of fit: ROC and AUC



- ROC - the trade-off between True Positive Rate (Sensitivity) and False Positive Rate (1 - Specificity)

## Goodness of fit: ROC and AUC



- ROC - the trade-off between True Positive Rate (Sensitivity) and False Positive Rate (1 - Specificity)
- AUC – Area under ROC

## Goodness of fit: Tests

- Likelihood ratio Test

## Goodness of fit: Tests

- Likelihood ratio Test
- $LR = 2(\loglik(m_1) - \loglik(m_0)) \sim \chi^2$



## Goodness of fit: Tests

- Likelihood ratio Test
- $LR = 2(\loglik(m_1) - \loglik(m_0)) \sim \chi^2$
- $H_0 : L_{reduced} > L_{current}$

## Goodness of fit: Tests

- Likelihood ratio Test
- $LR = 2(\loglik(m_1) - \loglik(m_0)) \sim \chi^2$
- $H_0 : L_{reduced} > L_{current}$
- $H_1 : L_{reduced} < L_{current}$

## Goodness of fit: Tests

- Likelihood ratio Test
- $LR = 2(\loglik(m_1) - \loglik(m_0)) \sim \chi^2$
- $H_0 : L_{reduced} > L_{current}$
- $H_1 : L_{reduced} < L_{current}$
- Hosmer-Lemeshow Test

## Goodness of fit: Tests

- Likelihood ratio Test
- $LR = 2(\loglik(m_1) - \loglik(m_0)) \sim \chi^2$
- $H_0 : L_{reduced} > L_{current}$
- $H_1 : L_{reduced} < L_{current}$
- Hosmer-Lemeshow Test
- $H_0 : \text{Observed proportion} = \text{Expected Proportion}$

## Goodness of fit: Tests

- Likelihood ratio Test
- $LR = 2(\loglik(m_1) - \loglik(m_0)) \sim \chi^2$
- $H_0 : L_{reduced} > L_{current}$
- $H_1 : L_{reduced} < L_{current}$
- Hosmer-Lemeshow Test
- $H_0 : \text{Observed proportion} = \text{Expected Proportion}$
- $H_1 : \text{Observed proportion} \neq \text{Expected Proportion}$

# Ideas for final Project

- Play with Goodness of fit

# Ideas for final Project

- Play with Goodness of fit
- LogReg for ordinal response dependent variable.

## Coding examples in R