

Stochastic Processes, Session 13 – Group Work

2D Point Processes

Problem 1: Binomial and Poisson Random Variables

Find the commands to draw binomial and Poisson distributed random numbers in Matlab, and read the help.

- Generate i.i.d. samples from a binomial distribution with $N = 10$ (number of trials) and $p = 0.3$. Generate a histogram and compare it with the probability mass function of the distribution (*Hint*: check the command `binopdf`).
- Generate i.i.d. samples from a poisson distribution with mean $\mu = 3$. Generate a histogram and compare it with the probability mass function of the distribution (*Hint*: check the command `poisspdf`).
- Next, write a program that plots the probability mass function for a binomial distribution with configurable number of trials N and success probability $p = 3/N$. Plot it for larger and larger values of N , and compare the results to the probability mass function generated in b).

Are you surprised by the results? In fact, the poisson distribution with mean μ can be seen as the limit of a binomial distribution with number of trials N and success probability $p = \mu/N$ when $N \rightarrow \infty$.

Problem 2: Binomial and Poisson Point Processes

Consider a 2D point process given by $X \sim \text{BinomialPP}(S, 12, f)$ with $S = [0, 2] \times [0, 2]$ and

$$f(x_1, x_2) = \begin{cases} 1/4, & (x_1, x_2) \in S \\ 0, & \text{otherwise.} \end{cases}$$

- Simulate and plot realizations of the point process X .
- Now, consider a region $B_1 = [0, 1] \times [0, 1] \subseteq S$. Discuss what the distribution of region count $N_X(B_1)$ is, write up its probability mass function and plot it.
- Extend your program to compute the region count $N_X(B_1)$ for each realization. Draw a large number of realizations of the point process and plot a histogram of the obtained region counts. Does the histogram fit with the results of the previous question?
- Next, include in your program code for computing the region count of $B_2 = S \setminus B_1$. Draw a large number of realizations and make a scatter plot of the obtained region counts $N_X(B_1)$ and $N_X(B_2)$. Can you observe any correlation between the two region counts?

Consider now instead a second point process given by $Y \sim \text{PoissonPP}(S, \varrho_Y)$. This is a homogeneous Poisson point process, defined also on S , and with a constant intensity function $\varrho_Y(y_1, y_2) = \varrho_0$.

- e) Write a program to generate and plot realizations of the point process Y , for any value of ϱ_0 .
- f) Set the value of ϱ_0 so that the process Y has, in average, the same number of points as X .
- g) Again, consider a region $B_1 = [0, 1] \times [0, 1] \subseteq S$. What is the distribution of region count $N_Y(B_1)$ now? Write up its probability mass function and plot it.
- h) Draw a large number of realizations of the point process and plot a histogram of the obtained region counts $N_Y(B_1)$. Does the histogram fit with the results of the previous question?
- i) With the same definition of $B_2 = S \setminus B_1$ as before, make a scatter plot of the obtained region counts $N_Y(B_1)$ and $N_Y(B_2)$. Compare your results to those you obtained for process X .

Problem 3: Inhomogeneous Poisson Point Process

Consider the inhomogeneous 2D poisson point process $X \sim \text{PoissonPP}(S, \varrho_x)$, with $S = [0, 1] \times [0, 1]$ and intensity function

$$\varrho_x(\mathbf{x}) = \varrho_x(x_1, x_2) = 30 \cdot x_1 \mathbb{1}[0 \leq x_1 \leq 1] \cdot \mathbb{1}[0 \leq x_2 \leq 1].$$

- a) What is the expected number of points falling in S ?
- b) Qualitatively, how will realizations of X look like? Sketch (by hand) a few figures of these realizations.
- c) Write a script in Matlab to draw realizations of this inhomogeneous Poisson point process. *Hint:* The inverse transformation method can be used for this. Also notice that x_1 and x_2 are independent.
- d) Now, change the intensity function to be $\tilde{\varrho}_x(\mathbf{x}) = \varrho_x(\mathbf{x}) \mathbb{1}[x_1^2 + x_2^2 \leq 1]$, where $\varrho_x(\mathbf{x})$ is the old intensity function given above. Sketch first the new space \tilde{S} . Then simulate this modified Poisson point process \tilde{X} .

Problem 4: Shot Noise and Campbell's theorem

For this problem you need to download the script `shotNoise.m` from Moodle.

- a) Go through `shotNoise.m` and discuss every line. Write up the shot noise model and the intensity function assumed in the script.
- b) Run the script and discuss the output. Try different values for ϱ_0 and observe what happens. Write down your observations.
- c) Discuss the effect of ϱ_0 on the estimated mean and on the Normal Probability Plot. Try to connect what you see to other theorems?
- d) Derive the expected value for the $Z(t)$ using Campbell's theorem and plot it on top of the estimated mean.