

# TITLE

AUTHORS

**Abstract.** Abstract

**Key words.** key words

**AMS subject classifications.** AMS ref numbers

## 1. Introduction.

### 1.1. Contributions.

### 1.2. Previous studies / State-of-the-art.

### 1.3. Organization of the paper.

### 1.4. Notation.

2. .

3. .

$$(3.1) \quad \begin{aligned} \min_{\beta, \tau} \quad & \frac{1}{2} \sum_{n=1}^N (F_n(\beta) \tau - \tilde{w}_n)^2 + \frac{1}{2} \lambda_\alpha \|\tau\|_2^2 + \lambda_\beta \mathcal{R}(\beta) \\ \text{s.t.} \quad & \tau \in \mathbb{R}_+^M, \beta \in ]0, 1[^M \end{aligned}$$

where  $\mathcal{R}$  is a regularization that is proper, convex and lower-semicontinuous (e.g. the total variation  $\mathcal{R}(\mathbf{x}) = \|\mathbf{D}\mathbf{x}\|_1$  with  $\mathbf{D}$  the discrete gradient matrix defined as the identity matrix with coefficients -1 on its subdiagonal).

Equation (3.1) has the typical form of a variable projection problem [1]. Indeed, it can be formulated as

$$(3.2) \quad \begin{aligned} \min_{\beta, \tau} \quad & \frac{1}{2} \left\| \begin{pmatrix} F(\beta) \\ \sqrt{\lambda_\alpha} \text{Id} \end{pmatrix} \tau - \begin{pmatrix} \tilde{w} \\ \mathbf{0} \end{pmatrix} \right\|_2^2 + \lambda_\beta \mathcal{R}(\beta) \\ \text{s.t.} \quad & \tau \in \mathbb{R}_+^M, \beta \in ]0, 1[^M \end{aligned}$$

4. .

## 5. Conclusion. Conclusion

## Appendix A. Proof of blablabla.

## REFERENCES

- [1] D. P. O'LEARY AND B. W. RUST, *Variable projection for nonlinear least squares problems*, Comput. Optim. Appl., 54 (2012), pp. 579–593, <https://doi.org/10.1007/s10589-012-9492-9>.