



Started on	Wednesday, 29 October 2025, 9:04 PM
State	Finished
Completed on	Wednesday, 29 October 2025, 9:10 PM
Time taken	5 mins 59 secs
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

Question 1 | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include<stdio.h>
2  int main(){
3      int n;
4      scanf("%d",&n);
5      int arr[n];
6      for(int i=0;i<n;i++){
7          scanf("%d",&arr[i]);
8      }
9      int k;
10     scanf("%d",&k);
11     for(int i=0;i<n-1;i++){
12         for(int j=0;j<n-i-1;j++){
13             if(arr[j]>arr[j+1]){
14                 int temp=arr[j];
15                 arr[j]=arr[j+1];
16                 arr[j+1]=temp;
17             }
18         }
19     }
20     int count=0;
21     for(int i=n-1;i>=0;i--){
22         for(int j=i-1;j>=0;j--){
23             if(arr[i]-arr[j]==k){
24                 count++;
25             }
26         }
27     }
28     if(count>=1){
29         printf("1");
30     }
31     else{
32         printf("0");

```

```
33 | }  
34 | }
```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)