


# Confidence Intervals

Interval Estimate



# Problem with point estimate

- Sometimes samples don't give quite the right result.
- We have seen how you can use point estimators to estimate the precise value of the population mean, variance, or proportion, but the trouble is, how can you be certain that your estimate is completely accurate?
- We need another way of estimating population statistics, one that allows for uncertainty.


# Confidence Intervals (CI)



# Confidence Intervals (CI)



# Confidence Intervals (CI)



We need to find a and b such that  $P(a < \mu < b) = 0.95$   
(a,b) is called the confidence interval

# Finding Confidence Interval

1. Choose your population statistic
2. Find its sampling distribution
3. Decide on the level of confidence
4. Find the confidence limits

# Cheat sheet for CI

Population statistic	Population distribution	Conditions	Confidence interval
$\mu$	Normal	You know what $\sigma^2$ is n is large or small $\bar{x}$ is the sample mean	$\left( \bar{x} - c \frac{\sigma}{\sqrt{n}}, \bar{x} + c \frac{\sigma}{\sqrt{n}} \right)$
$\mu$	Non-normal	You know what $\sigma^2$ is n is large (at least 30) $\bar{x}$ is the sample mean	$\left( \bar{x} - c \frac{\sigma}{\sqrt{n}}, \bar{x} + c \frac{\sigma}{\sqrt{n}} \right)$
$\mu$	Normal or non-normal	You don't know what $\sigma^2$ is n is large (at least 30) $\bar{x}$ is the sample mean $s^2$ is the sample variance	$\left( \bar{x} - c \frac{s}{\sqrt{n}}, \bar{x} + c \frac{s}{\sqrt{n}} \right)$
p	Binomial	n is large $p_s$ is the sample proportion $q_s$ is $1 - p_s$	$\left( p_s - c \sqrt{\frac{p_s q_s}{n}}, p_s + c \sqrt{\frac{p_s q_s}{n}} \right)$

# Confidence Interval

Level of confidence	Value of c
90%	1.64
95%	1.96
99%	2.58

CI  $\rightarrow$  statistic  $\pm$  (margin of error)

margin of error =  $c \times$  (standard deviation of statistic)



# Practice Problem

Q) We took a sample of 50 gems and found that in the sample, the proportion of red gems is 0.25. Construct a 99% confidence interval for the proportion of red gems in the population

# Practice Problem

Q) We took a sample of 50 gems and found that in the sample, the proportion of red gems is 0.25. Construct a 99% confidence interval for the proportion of red gems in the population

Sol: The CI for the population proportion is given by  $\left( p_s - c\sqrt{\frac{p_s q_s}{n}}, p_s + c\sqrt{\frac{p_s q_s}{n}} \right)$

# Practice Problem

Q) We took a sample of 50 gems and found that in the sample, the proportion of red gems is 0.25. Construct a 99% confidence interval for the proportion of red gems in the population

Sol: The CI for the population proportion is given by  $\left( p_s - c\sqrt{\frac{p_s q}{n}}, p_s + c\sqrt{\frac{p_s q}{n}} \right)$

Here  $p_s = 0.25$ ,  $q = 0.75$ ,  $n = 50$  and for 99% CI,  $c = 2.58$

# Practice Problem

Q) We took a sample of 50 gems and found that in the sample, the proportion of red gems is 0.25. Construct a 99% confidence interval for the proportion of red gems in the population

Sol: The CI for the population proportion is given by  $\left( p_s - c\sqrt{\frac{p_s q}{n}}, p_s + c\sqrt{\frac{p_s q}{n}} \right)$

Here  $p_s = 0.25$ ,  $q = 0.75$ ,  $n = 50$  and for 99% CI,  $c = 2.58$

Substituting we get (0.092, 0.408)