

Example Soil Testing Report

Sample Details

- Farmer / Client: Ramesh Kumar
- Sample ID: ST-2026-001
- Sample Collection Date: 12 July 2026
- Report Date: 17 July 2026
- Location: Vellore, Tamil Nadu
- Field Area: 1.5 acres
- Crop Planned: Groundnut
- Sample Depth: 0–15 cm
- Lab Reference: Demonstration report format only

Soil Test Results

Parameter	Result	Unit	Rating	Typical interpretation
pH	7.4	-	Neutral to slightly alkaline	Suitable for most field crops
Electrical Conductivity (EC)	0.32	dS/m	Normal	No salinity concern
Organic Carbon	0.48	%	Medium	Add organic matter regularly
Available Nitrogen (N)	245	kg/ha	Low	Nitrogen supplementation needed
Available Phosphorus (P2O5)	38	kg/ha	Medium	Maintenance dose recommended
Available Potassium (K2O)	290	kg/ha	High	Lower potash dose may be enough
Sulfur (S)	11	mg/kg	Medium	Monitor for oilseed crops
Zinc (Zn)	0.54	mg/kg	Low	Zinc application recommended
Iron (Fe)	4.8	mg/kg	Adequate	Usually sufficient
Manganese (Mn)	3.1	mg/kg	Adequate	Normal
Copper (Cu)	0.62	mg/kg	Adequate	Normal
Boron (B)	0.38	mg/kg	Low	Boron may improve flowering and pod set

Interpretation

The soil is generally in workable condition with normal salinity and near-neutral reaction. Organic carbon is moderate, but available nitrogen is low, which suggests a need for balanced fertilization with emphasis on nitrogen.

Phosphorus is in the medium range and potassium is high, so phosphorus can be applied at a maintenance level while potash can be reduced after considering crop demand. Zinc and boron are low, so micronutrient correction is advisable for better crop growth and yield.

Example Fertilizer Recommendation

For a groundnut crop under irrigated conditions, an example recommendation based on this sample could be:

- Apply 5 to 6 tons of well-decomposed farmyard manure per acre before sowing.
- Apply nitrogen in a moderate dose as per local university or department recommendation.
- Apply phosphorus at the recommended basal dose.
- Reduce potassium application if previous crop residues or irrigation water already contribute potassium.
- Apply zinc sulfate once before sowing or as per soil-test-based advisory.
- Apply boron carefully at a low dose only; excess boron can be harmful.
- Confirm final dosage with a local agricultural officer or soil testing laboratory because crop, irrigation, season, and soil type change the recommendation.

Field Advice

- Retest the field after one or two seasons if major nutrient corrections are made.
- Incorporate crop residues and compost to improve organic carbon.
- Avoid applying micronutrients without measured need.
- Take 8 to 10 subsamples from the field and mix them well for future testing.

Remarks

This document is a **sample format** created for demonstration and learning purposes. It is not an official laboratory certificate and should not be used as a real agronomic prescription without verified field sampling and lab analysis.