

FORGED IN FRAGMENTS: POPULATION III STAR FORMATION IN MILKY WAY PROGENITOR HALO

VASILII PUSTOVOIT[✉]

Canadian Institute for Theoretical Astrophysics and
 University of Toronto, Department of Physics

SHIVAN KHULLAR[✉]

Your nice institute / university and
 Your second nice institute / university

AND COLLABORATORS[✉]

Version August 17, 2026

Abstract

We present a radiation-magnetohydrodynamic (RMHD) cosmological zoom-in simulation of Population III star formation, drawing initial conditions from the FIRE m12f halo at high redshift. Starting from a self-consistent cosmological environment, we follow the collapse of a primordial gas cloud through the formation of an accretion disk that becomes gravitationally unstable within ~ 3 kyr of the first protostar’s formation. Prominent spiral arms develop and fragment vigorously, producing ~ 60 sink formation events over the ~ 15 kyr of post-collapse evolution, with a total stellar mass reaching $> 150 M_{\odot}$. The stellar mass grows as $M_{*} \propto t^2$, confirming the prediction of accelerating accretion in self-gravitating turbulent collapse, at rates of $\dot{M} \sim 10^{-3} - 10^{-2} M_{\odot} \text{ yr}^{-1}$. The disk is initially magnetically supercritical ($\mu \sim 100 - 300$) but the mass-to-flux ratio decreases by two orders of magnitude over the simulation, with the inner disk approaching and locally crossing the critical value ($\mu \lesssim 1$) by 15 kyr. Magnetic energy grows by ~ 4 orders of magnitude, likely driven by flux-freezing compression and differential rotation. The star formation efficiency reaches $\sim 80\%$ within the disk and $\sim 50\%$ within a 0.1 pc aperture. Our results demonstrate that cosmologically-realistic initial conditions, which produce asymmetric infall and complex angular momentum structure, lead to vigorous fragmentation in a single, large-scale accretion disk that hosts the entire protostellar population and whose turbulent velocity dispersion increases inward, driven by gravitational collapse rather than stellar feedback. ~~and a protostellar population whose properties differ from those found in earlier works.~~
~~[can’t say this because we don’t have an idealized set up to compare to in this paper. You should change~~

1. INTRODUCTION

The first generation of stars, Population III (Population III), formed from metal-free primordial gas in dark-matter minihaloes at redshifts $z \sim 20 - 30$ (Abel et al. 2002; Bromm et al. 2002). Because molecular hydrogen (H_2) was the only efficient gas coolant at these epochs, the primordial gas could cool to a temperature floor of $T \sim 200$ K at densities $n \sim 10^4 \text{ cm}^{-3}$, before reaching local thermodynamic equilibrium in its ro-vibrational levels (Bromm et al. 2002; Klessen and Glover 2023). The Jeans mass at this point, $M_J \sim 10^3 M_{\odot}$, is orders of magnitude larger than that of present-day star-forming cores, leading to the early expectation that Population III stars were predominantly massive and formed in isolation within each minihalo (Abel et al. 2002; Bromm

et al. 2002; Yoshida et al. 2008). These early simulations bridged cosmological collapse down to protostellar densities ($n \sim 10^{21} \text{ cm}^{-3}$), resolving the characteristic stages of primordial collapse: H_2 three-body formation heating at $n \gtrsim 10^8 \text{ cm}^{-3}$, collision-induced emission cooling at $n \gtrsim 10^{14} \text{ cm}^{-3}$, and a second collapse driven by H_2 dissociation at $n \sim 10^{16} \text{ cm}^{-3}$ (Yoshida et al. 2008; Wollenberg et al. 2020).

This picture of isolated, monolithic Population III star formation was overturned by simulations that continued to follow the system after the initial protostar formed. The infalling gas inevitably carries angular momentum, seeding a self-gravitating accretion disk prone to gravitational instability and fragmentation (Stacy et al. 2010; Clark et al. 2011; Greif et al. 2011). Rather than a single object, a protostellar cluster arises: Greif et al. (2011) found 2–5 fragments in five minihaloes within ~ 100 yr of the first protostar; Stacy et al. (2010, 2012) followed

systems for ~ 5 kyr and found ~ 10 sinks with a primary reaching $20\text{--}40 M_\odot$. Wollenberg et al. (2020) systematically explored 45 controlled simulations varying initial rotation and turbulence and found 5–20 sinks per halo, with fragmentation strongly stochastic from realization to realization. The protostellar cluster is shaped by competitive accretion (Bonnell et al. 2001), fragmentation-induced starvation (Wollenberg et al. 2020), protostellar mergers, and dynamical ejections, with ejection rates of $\sim 25\text{--}30\%$ found across multiple studies (Greif et al. 2011; Stacy et al. 2012; Wollenberg et al. 2020).

The Population III initial mass function (IMF) inferred from these simulations is generally thought to be top-heavy, spanning sub-solar to $\gtrsim 100 M_\odot$, but highly variable from halo to halo. [CITATIONS?] Hirano et al. (2014) surveyed 100 minihaloes and found accretion rates of $10^{-4}\text{--}10^{-2} M_\odot \text{ yr}^{-1}$ and final masses of $\sim 10\text{--}1000 M_\odot$; Susa et al. (2014) found masses spanning $1\text{--}> 100 M_\odot$ across several tens of halos. When radiative feedback is included, accretion onto a single protostar can be halted at $\sim 43 M_\odot$ (Hosokawa et al. 2011), though episodic accretion from a fragmenting disk can produce intermittent feedback and final masses from ~ 10 to several hundred $M_\odot$ (Hosokawa et al. 2016; Stacy et al. 2016). The large statistical survey of Hirano et al. (2023) across > 100 cosmological halos confirms that the Population III IMF is not universal: the outcome depends on halo mass, angular momentum, and baryon streaming velocity, and a single simulation cannot be over-generalised.

Turbulence plays a fundamental role in shaping fragmentation and the resulting mass distribution. Chen et al. (2025) demonstrated that cosmological infall into dark-matter potential wells universally drives supersonic turbulence in Population III minihaloes, with Mach numbers $\mathcal{M} \sim 2\text{--}4$ that increase with halo mass. Wollenberg et al. (2020) confirmed that the degree of initial turbulence strongly affects the number of fragments and the resulting mass distribution, with higher initial turbulent velocities producing more fragments and more stochastic outcomes. Prieto et al. (2011) showed that turbulent fragmentation in a larger, $3 \times 10^7 M_\odot$ halo at $z \sim 11$ produces ~ 25 gravitationally unstable clumps, proposing that infall-driven turbulence may be the dominant fragmentation mechanism when halo masses exceed the canonical $10^6 M_\odot$ minihalo scale.

Magnetic fields can further modify Population III star formation, though their net effect remains debated. Even trace cosmological seed fields are amplified during gravitational collapse by flux-freezing and winding, driving the mass-to-flux ratio toward criticality over timescales of a few kyr (Machida et al. 2008; Sharda et al. 2021; Higashi et al. 2024). Idealised MHD simulations suggest that strong fields can suppress disk fragmentation (Hirano and Machida 2022), and cosmological MHD runs by Stacy et al. (2022) found only two secondary sinks. In contrast, Prole et al. (2022) and Sadanari et al. (2024) found that even dynamically significant fields reduce but do not eliminate fragmentation. Sharda and Menon (2025) performed the first 3D radiation-MHD Population III simulations with non-equilibrium chemistry, turbulence, and ionising feedback, finding that magnetic fields are more effective than radiation in limiting protostellar masses at early times, capping the most massive star

at $\sim 65 M_\odot$ within 5 kyr (see also Sharda et al. 2025). These conflicting results indicate that the outcome depends sensitively on initial field strength, field geometry, resolution, and simulation duration (Klessen and Glover 2023).

While many Population III simulations now use cosmological initial conditions (Greif et al. 2011; Stacy et al. 2016; Hirano et al. 2023; Latif and Khochfar 2022; Meziani et al. 2026), simulations that simultaneously combine cosmological ICs, full radiation-MHD, and resolved individual protostellar formation remain few. Meziani et al. (2026) use the FORGE’d in FIRE framework (Hopkins et al. 2023) to study Population III star formation in an ultra-faint dwarf galaxy progenitor, with a DM halo mass of $10^7 M_\odot$ at $z=21$ and $10^{12} M_\odot$ at $z=0$. A minihalo embedded in a strong overdensity destined to grow to Milky Way mass may experience qualitatively different infall rates, tidal torques, and angular momentum injection compared to one in a quieter environment destined to remain a dwarf galaxy. Therefore, in this paper, we zoom-in to the formation of the first stars in a massive halo $10^7 M_\odot$ that eventually grows to Milky Way mass at $z=0$.

Whether the environmental difference between a MW-progenitor (Milky Way) and a UFD-progenitor (Ultra Faint Dwarf) drives a qualitatively different outcome is an open question. In this work, we address this question directly. We perform a cosmological radiation-magnetohydrodynamic (RMHD) zoom-in simulation of Population III star formation in the FIRE m12f initial conditions (Hopkins et al. 2018; Wetzel et al. 2023), which in turn were taken from the AGORA project (Kim et al. 2013, 2016). The ICs constitute a halo destined to become a $M_{\text{vir}} \sim 10^{12} M_\odot$ Milky Way analogue by $z = 0$. We use the FORGE’d in FIRE framework (Hopkins et al. 2023; Meziani et al. 2026), combining FIRE-3 (Hopkins et al. 2022) large-scale physics with STARFORGE (Grudić et al. 2021) protostellar physics, and evolve the system for ~ 15 kyr after the first protostar forms, reaching a gas mass resolution of $\Delta m \sim 10^{-3} M_\odot$. We study the thermodynamic collapse, disk formation and fragmentation, magnetic field amplification, and protostellar mass assembly in this environment, providing the first cosmological RMHD simulation that resolves individual Population III protostellar formation in a MW-progenitor halo.

The rest of the paper is organised as follows. section 2 describes the simulation setup, physics modules, refinement strategy, and analysis methods. section 3 presents the disk morphology, thermodynamic and energetic evolution, disk stability, and magnetic field structure. subsection 3.4 discusses the star formation history, accretion rates, and protostellar mass function. section 4 compares our results to previous Population III simulations. We state our conclusions in section 5.

2. SIMULATION AND METHODS

2.1. Simulation details

For this paper we utilise the FORGE’d in FIRE framework of Hopkins et al. (2023), extended to the primordial star-formation regime similar to Meziani et al. (2026). We employ FIRE-3 (Hopkins et al. 2022) physics modules at low resolution and STARFORGE (Grudić et al.

2021) physics to resolve individual protostellar formation. We briefly review these below and refer the reader to Hopkins et al. (2022, 2023); Grudić et al. (2021). Numerical parameters not specified below are identical to those described in the above references.

2.1.1. Code and gravity solver

The governing equations solved by the code are the ideal MHD equations in conservative form (Hopkins 2015; Hopkins and Raives 2015):

$$\frac{\partial \mathbf{U}}{\partial t} + \nabla \cdot \mathbf{F} = \mathbf{S}, \quad (1)$$

where the state vector of conserved quantities and the flux tensor are

$$\mathbf{U} = \begin{pmatrix} \rho \\ \rho \mathbf{v} \\ \rho e \\ \mathbf{B} \end{pmatrix}, \quad \mathbf{F} = \begin{pmatrix} \rho \mathbf{v} \otimes \mathbf{v} + P_T \mathcal{I} - \mathbf{B} \otimes \mathbf{B} \\ (\rho e + P_T) \mathbf{v} - (\mathbf{v} \cdot \mathbf{B}) \mathbf{B} \\ \mathbf{v} \otimes \mathbf{B} - \mathbf{B} \otimes \mathbf{v} \end{pmatrix}. \quad (2)$$

Here ρ is the mass density, $\mathbf{v}$ the velocity, $\mathbf{B}$ the magnetic field (in Gaussian units with 4π absorbed), $e = u + |\mathbf{B}|^2/(2\rho) + |\mathbf{v}|^2/2$ the total specific energy, $P_T = P + |\mathbf{B}|^2/2$ the total (thermal plus magnetic) pressure, and $\mathcal{I}$ the identity tensor. $\mathbf{S}$ includes source terms for self-gravity and the Powell divergence-cleaning scheme (Hopkins and Raives 2015) that maintains $\nabla \cdot \mathbf{B} \approx 0$.

The FORGE'd in FIRE framework is based on GIZMO Meshless Finite Mass (MFM) code (Hopkins 2015). It is a mesh-free Lagrangian Godunov method in which the fluid equations are solved on a set of moving partition-generating points whose masses are conserved exactly (no inter-particle mass flux), while fluxes of momentum, energy, and magnetic flux are exchanged through effective interparticle faces defined by a kernel-weighted partition of unity applied to the particle distribution (Hopkins 2015).

Gravity is solved with the TreePM algorithm (Hopkins 2015), which combines a particle-mesh grid for large-scale forces with an oct-tree for short-range forces (Barnes and Hut 1986), as in Springel (2010). Gas particles use adaptive Lagrangian gravitational softening matching the hydrodynamic kernel radius. Sink/star particles use a fixed Plummer-equivalent softening $\epsilon \approx 20$ AU. Direct N -body summation is used for all sink-sink and sink-gas gravitational interactions, with direct summation applied for sink-sink separations below 1000 AU to accurately resolve close binaries and dynamical ejections (Grudić et al. 2021).

2.1.2. Initial Conditions

We use initial conditions (ICs) developed as part of the FIRE collaboration (Hopkins et al. 2018), specifically the m12f halo from the Latte suite of Milky Way-mass zoom-in simulations (Wetzel et al. 2023). The m12f halo has a virial mass $M_{\text{vir}} \sim 1\text{--}2 \times 10^{12} M_\odot$ at $z = 0$ and was selected as an isolated system with a relatively quiescent late-time merger history, making it a Milky Way analogue. At the redshift of first collapse in our simulation ($z \sim 21$), CDM part of the progenitor halo has a mass of $2 \times 10^7 M_\odot$, typical of the minihaloes in which Population III stars are expected to form. This value

was obtained from running the ROCKSTAR halo finder (Behroozi et al. 2013) on our snapshot.

We adopt pre-generated ICs at $z = 99$ in a $60 \text{ Mpc}/h$ comoving cosmological box. The box contains a zoom-in region of comoving extent $\sim 3 \text{ Mpc}/h$, within which the FIRE galaxy forms at $z = 0$. Only the zoom-in region contains baryonic particles and high-resolution dark matter (DM) particles; the remainder of the volume is populated with low-resolution DM particles. As in the FIRE-2 Latte simulations, we adopt cosmological parameters from WMAP-7 (Komatsu et al. 2011): $h = 0.702$, $\Omega_m = 0.272$, $\Omega_\Lambda = 0.728$, $\Omega_b = 0.0455$, $\sigma_8 = 0.807$, and $n_s = 0.961$.

The full simulation snapshot contains ~ 88 million CDM particles and FIRE-resolution gas particles whose smoothing lengths ($\sim \text{kpc}$) far exceed the protostellar disk scale. We extract spatial cutouts centred on the most-massive sink particle with a physical radius of 1 kpc (22 ckpc). The cutout sphere retains only the gas, high-resolution DM, and sink/star particles. All analysis presented in this manuscript is of this cutout simulation.

Over the ~ 15 kyr of post-collapse evolution studied here, the sound-crossing time across the cutout radius ($r_{\text{cut}} \approx 1 \text{ kpc}$) is $\sim r_{\text{cut}}/c_s \gtrsim 10 \text{ kyr}$ for $T \lesssim 10^4 \text{ K}$, comparable to or longer than our integration time. The large-scale structure outside the cutout therefore has no causal influence on the disk dynamics on these timescales, and the cutout fully captures the relevant physics.

2.1.3. Refinement strategy

We first evolve the full cosmological simulation with FIRE-3 physics until first $10^5 M_\odot$ stellar population (SSP) particle forms. We then restart from the preceding snapshot with an additional ‘‘hyper-refinement’’ layer applied globally (Hopkins et al. 2023).

We refine gas cells (splitting into pairs of equal-mass daughter cells) whenever their mass exceeds a fraction f_J of the local Jeans mass:

$$\begin{aligned} \Delta m &> f_J M_J \\ &= f_J \frac{\pi}{6} \rho \lambda_J^3 \\ &= f_J \frac{\pi}{6} \rho \left(\frac{\pi c_s^2}{G \rho} \right)^{3/2} \\ &= f_J \frac{\pi^{5/2}}{6} \left(\frac{k_B T}{\mu m_H G} \right)^{3/2} \rho^{-1/2} \\ &= f_J M_0 \left(\frac{T}{10^4 \text{ K}} \right)^{1.5} \left(\frac{n_H}{\text{cm}^{-3}} \right)^{-0.5} \end{aligned} \quad (3)$$

The last line follows from substituting the thermal Jeans length $\lambda_J = (\pi c_s^2 / G \rho)^{1/2}$ and $c_s^2 = k_B T / (\mu m_H)$ into $M_J = (\pi/6) \rho \lambda_J^3$, and normalizing at $T_0 = 10^4 \text{ K}$ and $n_0 = 1 \text{ cm}^{-3}$. We use $f_J = 10^{-2}$, corresponding to ~ 100 resolution elements per Jeans mass and satisfying the Truelove et al. (1997) criterion, since we resolve $100^{1/3} \approx 4.6 > 4$ cells per Jeans length. This is sufficient since MFM does not suffer from artificial fragmentation even while refining (Khullar et al. 2026, in prep.). We also speculate that since MFM is generally less diffusive than AMR, the more stringent criteria of 30 cells per Jeans length needed to resolve effects like magnetic




FIG. 1.— Top: Radial profile of gas particle mass (resolution). Scatter plot shows particle distribution at the time of first sink formation; solid line and shaded region correspond to the mean and $\pm 1\sigma$ for radial profiles across all snapshots after first sink formation. Horizontal lines correspond to $1/2\Delta m$ of previous resolution region, since our particle splitting algorithm is subdividing parent particles in 2 for each new level of split. Bottom: Same as Top, but for radial profile of equivalent cell size $\Delta x = (\Delta m/\rho)^{1/3}$.

dynamo amplification (Federrath et al. 2011) does not necessarily apply here. Since our simulation produces magnetic amplification by up to 4 orders of magnitude (see subsection 3.3), we leave detailed numerical checks to future work.

Only gas particles get split, DM particles stay at the same resolution. This is a fine approximation because dark matter’s potential is smooth at these scales, and splitting of these particles should not change the results of our simulation.

At our highest refinement level, the gas mass resolution is $\Delta m \approx 10^{-3} M_{\odot}$ and the spatial resolution is $\Delta x \approx 2$ AU, sufficient to resolve densities up to $n \sim 10^{15} \text{ cm}^{-3}$.

Figure 1 shows the radial profile of the length scale of gas particles $\lambda = (m/\rho)^{1/3}$. It exhibits the refinement profile, described in subsection 2.1.3 at all times, except the earliest one. Initial profile at $t - t_1 = -0.5$ (0.5 kyrs before the first sink formation) is limited by the refinement scale inside of 100 AU radius (the center was not yet refined to the highest resolution). But by the time of first sink formation, the refinement reaches highest resolution that we expect from our scheme, (around $10^{-3} M_{\odot}$ or a few AU) so the inner star forming disk

region is fully refined.

2.2. Resolved physics

2.2.1. Magnetic fields

We integrate the ideal MHD equations as described in Hopkins and Raives (2015), using the constrained-gradient method and divergence cleaning (Hopkins 2016) for divergence control, which has been validated for magnetorotational instability dynamics (Hopkins et al. 2023).

We include Spitzer–Braginskii thermal conduction and viscosity via Hopkins (2017). For metal-free gas at the temperatures ($T \sim 200\text{--}10^4$ K) and free-electron fractions relevant here, these transport timescales exceed the local dynamical timescale by many orders of magnitude (Meziani et al. 2026), so ideal MHD dominates throughout the simulation.

The initial magnetic field is a trace cosmological seed field of $|B_0| \sim 10^{-15}$ G amplified self-consistently by gravitational compression, differential rotation, and the turbulent dynamo.

2.2.2. Radiation

We solve radiation transport with the M1 moments method (Levermore 1984), as implemented in GIZMO and described in Hopkins et al. (2023); Grudić et al. (2021); Hopkins and Grudić (2019); Hopkins et al. (2020) across five frequency bands: H-ionizing, Lyman–Werner (LW)/photo-electric, Near-UV (NUV), optical–NIR, and an adaptive-wavelength FIR band that tracks the evolving dust emission temperature (Hopkins et al. 2022). Gas, dust, and radiation temperatures are evolved separately with appropriate coupling terms, allowing self-consistent treatment of both optically thin and optically thick regimes. We use a reduced speed of light $\tilde{c} = 10^{-3}c$ following Hopkins et al. (2023).

The Lyman–Werner (LW) band is combined with the photo-electric (PE) heating band into a single radiation frequency bin. This is justified because both LW and PE photons originate from the same protostellar sources and have similar mean free paths at the densities resolved here. H_2 photo-dissociation by LW photons is accounted for through the Γ_{H_2} destruction term in Equation 4.

2.2.3. Thermochemistry

We solve gas thermodynamics and chemistry as in (Hopkins et al. 2022). The following cooling functions are employed and are relevant to our setup: Λ_{H} (atomic H cooling); Λ_{HD} (HD cooling, Galli and Palla 1998); Λ_{H_2} (H_2 cooling); $\Lambda_{\text{H}_2}^{\text{thin}}$, $\Lambda_{\text{H}_2}^{\text{thick}}$ (H_2 interactions with e^- , H^+ , H_2 , H, He in optically thin and thick regimes). We follow non-equilibrium H and He chemical networks. Default FIRE-3 thermochemistry includes cooling due to metal lines, but we do not evolve our simulation past the point of supernova-driven metal enrichment, thus making all metal cooling contributions effectively zero (as expected in the Population III formation regime).

Since we simulate metal-free primordial gas, the dust-to-gas ratio is zero and the dust term in the H_2 formation equation is absent. H_2 formation proceeds exclusively via gas-phase channels: the two-body H^- and H_2^+ pathways at low densities ($n \lesssim 10^8 \text{ cm}^{-3}$) and the three-body reaction ($3\text{H} \rightarrow \text{H}_2 + \text{H}$) at high densities.

Molecular fraction of H_2 evolves in every cell according to (eq. 6 from Hopkins et al. 2022), with the dust formation term set to zero:

$$\begin{aligned} \frac{d\bar{n}_{\text{H}_2}}{dt} = & \alpha_{\text{GP}} C_2 \bar{n}_{\text{HI}} \bar{n}_{\text{H}} + \alpha_{3\text{B}} C_3 \bar{n}_{\text{HI}} \frac{\bar{n}_{\text{H}_2}}{8} \left(\bar{n}_{\text{HI}} + \frac{\bar{n}_{\text{H}_2}}{8} \right) \\ & - \Gamma_{\text{H}_2} \bar{n}_{\text{H}_2} - \xi_{\text{H}_2} \bar{n}_{\text{H}_2} - \sum_i \beta_{\text{H}_2,i} C_2 \bar{n}_{\text{H}_2} \bar{n}_i \end{aligned} \quad (4)$$

Here $\bar{n}_x$ is volume-averaged number density of x inside of a cell. $C_m = \langle n^m \rangle / \langle n \rangle^m$ is the m^{th} -order clumping factor. α_{GP} is the gas-phase formation rate coefficient via the H^- and H_2^+ pathways (Glover and Jappsen 2007; Abel et al. 1997; Galli and Palla 1998). The three-body formation coefficient $\alpha_{3\text{B}}$ (Abel et al. 2002; Palla et al. 1983) becomes the dominant H_2 formation channel for $n \gtrsim 10^8 \text{ cm}^{-3}$. Γ_{H_2} is the coefficient associated with LW and ionizing photon dissociation. ξ_{H_2} accounts for cosmic-ray-induced ionization and dissociation ($\xi_{\text{H}_2} \approx 3 \times 10^{-17} \text{ s}^{-1}$, the Milky Way value; this rate is unconstrained at $z \sim 21$ and primarily affects H_2 formation at $n \lesssim 10^7 \text{ cm}^{-3}$, with negligible impact at the densities resolved here). The last sum runs over collisional dissociation contributions from e^- , H^+ , H_2 , HI , HeI , HeII , with rate coefficients $\beta_{\text{H}_2,i}$ from Glover and Abel (2008).

Thermodynamic energy release from H_2 dissociation ($\sim 4.5 \text{ eV}$ per molecule) is not explicitly tracked in the gas energy equation. This is acceptable because we do not have enough resolution to reach number densities of $n \approx 10^{16} \text{ cm}^{-3}$, where H_2 dissociation kicks in and becomes a non-negligible thermodynamic variable. (Klessen and Glover 2023) In addition, the protostellar evolution tracks of Offner et al. (2009) self-consistently evolve the stellar radius without requiring the gas to track dissociation energy explicitly.

At $z \sim 21$, the meta-galactic UV background (Chan et al. 2019) is negligible compared to local protostellar radiation resolved in the simulation. H_2 self-shielding against Lyman-Werner photons is included using the shielding function of Gnedin and Draine (2014) (their Eq. 3), applied with a local Sobolev column density $N_{\text{H}} \sim \rho / |\nabla \rho|$, with turbulent line broadening accounted for following a Burgers velocity spectrum.

The simulation does not include cosmic-ray transport. In our simulation, which covers only $\sim 15 \text{ kyr}$ of post-collapse evolution and does not include supernovae, cosmic-ray injection from stellar feedback is negligible; the initial cosmic-ray background at $z \sim 21$ contributes subdominantly to H_2 dissociation and ionization relative to the resolved radiation field.

2.2.4. Stellar formation and evolution

We resolve the formation of individual protostars in the densest region of our simulation using the STARFORGE (Grudić et al. 2021) formation and evolution modules. A gas particle becomes a sink (protostar) particle when it meets the full STARFORGE formation criteria: the cell must be (i) locally self-gravitating (virial parameter $\alpha_{\text{vir}} < 1$), (ii) Jeans unstable, (iii) in a converging flow ($\nabla \cdot \mathbf{v} < 0$), (iv) experiencing compressive tides in all directions, (v) a local maximum of the density field, and (vi) not within the accretion radius of a pre-existing sink (Grudić et al. 2021).

Once formed, sink particles accrete bound gas whose circularised orbit falls within the sink radius. Each sink evolves along protostellar tracks following Offner et al. (2009), explicitly tracking the stellar radius, luminosity, and effective temperature as a function of the instantaneous mass and accretion rate. Protostellar jets are launched with a mass-loading proportional to the surface accretion rate and a launch velocity comparable to the escape velocity from the protostellar surface (Grudić et al. 2021). **Jets are therefore expected in this simulation, but they are not resolved in the present run: a bug in the launch implementation suppresses them. We defer the analysis of jet feedback to future work.**

2.3. Postprocessing methods

2.3.1. Disk identification

To compute velocity dispersion, local Toomre Q , and other disk properties, we identify gas particles belonging to the disk at each snapshot using the following procedure:

1. We place the disk centre at the most massive sink particle. This is more stable than the mass-weighted centre-of-mass (COM) of all sinks once secondary sinks form at large separations. Before any sinks form, the centre is placed at the densest gas particle within a refinement region.
2. We compute a mass-weighted COM velocity v_{COM} from gas particles within $r < r_{\text{search}} = 10^{-5} \text{ kpc} \approx 2060 \text{ AU}$ and subtract it from all particle velocities.
3. We compute the disk normal $\hat{L}$ from the total angular momentum $\mathbf{L} = \sum_i m_i (\mathbf{r}_i \times \mathbf{v}_i)$ summed over particles within r_{search} , using the COM-subtracted velocities from the previous step. The sign of $\hat{L}$ is fixed so that $L_z > 0$ to prevent frame-to-frame flipping.
4. We project particle positions and velocities onto the disk-plane coordinate system (r_{cyl} , ϕ , z , v_r , v_ϕ , v_z).
5. We apply geometric and density cuts: $r_{\text{cyl}} < r_{\text{max}} = 10^{-5} \text{ kpc}$, $|z|/r_{\text{cyl}} < 0.3$ (aspect ratio), and $\rho > \rho_{\text{thresh}} = 10^{-15} \text{ g cm}^{-3}$.
6. We apply kinematic cuts: $v_\phi > 0$ (co-rotating) and $v_\phi/v_K > f_{\text{kep}} = 0.3$, where $v_K = \sqrt{GM_{\text{enc}}(< r)}/r$ is the Keplerian speed computed from the cumulative enclosed gas mass plus the total sink mass placed at the origin.
7. We extend the kinematically selected set to its bounding cylinder (100th percentile in r_{cyl} and $|z|$) to recover any disk material just outside the strict kinematic boundary.

Particles passing all cuts are labelled as disk members. The disk radius r_{disk} is taken as the 90th-percentile cylindrical radius of disk members.




FIG. 2.— Multi-scale density projection of the simulation at $t - t_1 \approx 15$ kyr, illustrating the dynamic range from the cosmic web (box size of ≈ 100 kpc) to the center of the protostellar disk (resolving AU scales). White boxes mark the region shown in the next panel. White star symbols mark sink particles.

3. RESULTS

In this section we present results from our cosmological RMHD simulation of Population III star formation. We organize the discussion as follows: [subsection 3.1](#) examines the thermodynamic and energetic evolution during the pre-stellar collapse and early disk phase; [subsection 3.2](#) characterizes the disk structure and gravitational stability; [subsection 3.3](#) describes the magnetic field evolution; and [subsection 3.4](#) covers the star formation history, accretion rates, and protostellar mass function.

Figure 2 presents a multi-scale density projection of the simulation at $t - t_1 \approx 15$ kyr, illustrating the range of scales our simulation resolves after the onset of active star formation.

3.1. Collapse

The assembly of a primordial protostar proceeds through a sequence of thermally regulated stages: the gas cools via H_2 line emission until line optical depth and three-body reactions convert the flow into a fully molecular, opaque core, at which point compressional heating halts the collapse and a protostellar seed forms. We examine the thermodynamic state of the infalling gas and the kinematics of the large-scale accretion flow that feeds the disk.

3.1.1. Thermodynamics

The thermal and chemical state of the gas encodes the balance between the competing processes of compressional heating and molecular cooling. The temperature-density phase trajectory is therefore a direct diagnostic of which physical process dominates at each stage of collapse, and f_{H_2} molecular fraction phase plot shows the efficiency of molecular cooling. We include these phase space plots in [section 3](#) for three representative epochs

(pre-stellar, first sink formation, post-first sink formation). Disk particles (identified by the algorithm of [subsection 2.3.1](#)) are marked in green, particles that are a part of full simulation but not cutout are marked in blue, and particles that are present in cutout simulation but not identified as disk particles are marked in red.

At the pre-stellar epoch ($t - t_1 = -0.5$ kyr; [section 3](#), left panels), the gas occupies a narrow track spanning $n \sim 10^7 - 10^{10} \text{ cm}^{-3}$ and $T \sim 200 - 800$ K. This track traces the initial slow collapse, during which H_2 rovibrational cooling brings the gas down to the ~ 200 K temperature floor, followed by gradual compressional heating as H_2 line cooling saturates and becomes optically thick. The H_2 fraction (lower panel) rises modestly from $f_{H_2} \sim 10^{-2.5}$ at $n \sim 10^7 \text{ cm}^{-3}$ to $f_{H_2} \sim 10^{-1.3}$ at $n \sim 10^{10} \text{ cm}^{-3}$, driven primarily by two-body formation channels (H^- and H_2^+ pathways). Note that increase in H_2 fraction coincides with cooling of the gas, since H_2 is much more efficient coolant compared to atomic H. The low-density gas particles ($n \leq 10^{-5} \text{ cm}^{-3}$) in the left column correspond to low-resolution gas in the original zoom-in box before re-simulation (blue markers). Most of those particles have $f_{H_2} = 0$, thus not present in the bottom plot. This part of the simulation domain is outside our region of interest and does not change significantly over our integration time of ≈ 15 kyr.

The T - n track at the moment of first sink collapse ($t - t_1 = 0$) follows the $Z = 10^{-4} Z_\odot$ metallicity track of [Omukai \(2000\)](#); [Klessen and Glover \(2023\)](#) rather than the zero-metallicity track. We speculate that this slight offset may reflect small departures from chemical equilibrium in the collapsing gas; we do not impose a metallicity floor, so trace metal enrichment from surrounding regions cannot be ruled out as a contributing factor. By the later epoch ($t - t_1 = 1.2$ kyr), the median of the phase distri-




FIG. 3.— Phase diagram (temperature and H_2 fraction vs. number density) at three representative epochs: pre-stellar ($t - t_1 \approx -0.5$ kyr, left), first sink formation ($t - t_1 = 0$, middle), and post-first-sink ($t - t_1 = 1.2$ kyr, right). Green points correspond to disk-identified particles, red points correspond to points in our cutout simulation that have not been identified as disk, and blue points correspond to points in the full-box simulation right before the cutout was applied (only included in the leftmost plots). $f_{H_2} = 0$ for most of particles in the low-density region (blue particles), thus missing from the bottom-left panel. Solid black curves show the zero-metallicity Population III collapse track (Omukai (2000), as digitized by Klessen and Glover (2023)) on the temperature (top) and H_2 fraction (bottom) panels. Coloured curves with shaded bands are binned medians of the simulated gas (the bands span $\pm 1\sigma$): blue is the instantaneous median at the pre-stellar epoch; dark red is the mean of the per-snapshot medians over $-0.5 \text{ kyr} \leq t - t_1 \leq 0$, and dark blue over $0 \leq t - t_1 \leq 1.2 \text{ kyr}$.

bution shifts back toward the $Z = 0$ track, suggesting that the gas at this stage is governed more by primordial H_2 chemistry. We defer a systematic one-zone test of this offset to future work.

By $t - t_1 = 0$ (section 3, middle panels), the gas has collapsed to much higher densities ($n \sim 10^{13} \text{ cm}^{-3}$) and the thermal track has extended significantly. At intermediate densities ($n \sim 10^8 - 10^{10} \text{ cm}^{-3}$), compressional heating outpaces H_2 line cooling, which saturates as the lines become optically thick, and the temperature rises steadily through $T \sim 500 - 800$ K. The H_2 fraction increases steeply in this regime as three-body formation reactions ($3H \rightarrow H_2 + H$) become efficient at $n \gtrsim 10^8 \text{ cm}^{-3}$, converting the gas to fully molecular ($f_{H_2} \rightarrow 1$) by $n \sim 10^{11} - 10^{12} \text{ cm}^{-3}$. Note that unlike in the pre-stellar regime, we see an increase in temperature associated with increase in H_2 fraction. This is a second-order effect: H_2 cools the gas very effectively, reducing thermal pressure, allowing the gas to continue the infall process. This makes the gas opaque to H_2 lines, heating it up. (Ripamonti and Abel 2004)

At $t - t_1 = 1.2$ kyr (section 3, right panels), the phase diagram has broadened substantially. The gas has separated into 2 parts: a continuously-infalling branch with $T < 2000$ K and $n < 10^{10} \text{ cm}^{-3}$, and a hot branch at $T \sim 2000 - 8000$ K from shock-heated and adiabatically compressed gas. The H_2 fraction now shows a bimodality at high densities: gas that has been heated to $T \gtrsim 2000$ K via radiation produced by the first formed sink undergoes collisional H_2 dissociation. We note that while our thermochemical model does not couple H_2 dissociation

to thermodynamics, the $\beta_{H_2,i}$ terms in Equation 4 can still drive partial dissociation of H_2 .

A low- f_{H_2} population ($f_{H_2} \sim 10^{-3} - 10^{-5}$) is produced at $n \sim 10^{10} - 10^{12} \text{ cm}^{-3}$, while the lower-density gas keeps following the infall trajectory, driven by 2-body H_2 production pathway that the collapse started with (i.e. left panels). (Klessen and Glover 2023)

At later times, the three-body pathway to H_2 production becomes dominant way of converting atomic hydrogen to molecular in collapsing regions. This results in constantly increasing fraction of molecular hydrogen in the disk with time.

3.1.2. Pre-stellar outside-in collapse

A gravitationally supercritical core – one in which the enclosed mass exceeds the local Jeans mass – collapses outside-in: the outer shells fall inward faster than the inner ones because the excess of gravity over thermal pressure gradient increases with radius. This contrasts with the inside-out expansion wave picture of Shu (1977), which applies only when a core is marginally critical. Here we characterize the accretion geometry and confirm that the simulated collapse is indeed driven from the outside in.

We compute the gas mass in spherical shells centered on the densest particle in the disk (which will later correspond to the first sink particle location). We perform this calculation for every snapshot, with a time interval between snapshots of $\Delta t \approx 0.1$ kyr, and show the shell mass profiles in the top panel of section 5. We show the rate of mass change per shell, $\Delta m / \Delta t$, in the bottom




FIG. 4.— Top: Time evolution of the main energy components computed for disk-identified particles (the disk radius r_{disk} is taken as the 90th-percentile cylindrical radius of disk-identified particles, see subsection 3.2): rotational kinetic energy E_{rot} (blue), turbulent kinetic energy E_{turb} (green), total kinetic energy $E_{\text{kin}} = E_{\text{rot}} + E_{\text{turb}}$ (black, dash-dotted), gravitational potential energy $|E_{\text{pot}}|$ (red), thermal energy E_{therm} (orange, dotted), and magnetic energy E_{mag} (magenta, dashed). Bottom: pure kinematic virial ratio $\alpha_{\text{kin}} = 2E_{\text{kin}}/|E_{\text{pot}}|$ (black); virial ratio with added thermal component $\alpha_{\text{kin+th}} = 2(E_{\text{kin}} + E_{\text{therm}})/|E_{\text{pot}}|$, and with added thermal and magnetic components (purple, dash-dotted), $\alpha_{\text{vir}} = 2(E_{\text{kin}} + E_{\text{therm}} + E_{\text{mag}})/|E_{\text{pot}}|$. The dotted red line marks virial equilibrium ($= 1$). The disk starts sub-virial and reaches equilibrium by ~ 2 kyr. The inclusion of thermal energy reveals that the disk is thermally supported and near virial equilibrium at all times. Magnetic energy provides negligible additional support to the virial ratio.

panel.

The fact that the outer layers gain mass faster demonstrates an outside-in collapse. This contrasts with the inside-out model of Shu (1977) and aligns with the collapse dynamics of gravitationally unstable Bonnor-Ebert spheres (Ebert 1955; Bonnor 1956; Larson 1969; Penston 1969). Figure 7 confirms this: the infall velocity ($-v_r$) is supersonic in the envelope outside the disk ($r > 2500$ AU), which is a definitive signature of outside-in dynamics. Within the disk ($r < 2500$ AU), the flow becomes predominantly subsonic. This abrupt deceleration occurs because the infalling gas encounters an accretion shock as it strikes the dense, rotationally supported disk material.

3.2. Disk evolution

Once the accretion flow acquires sufficient angular momentum to form a centrifugal barrier, a rotationally supported disk grows around the central protostar. The




FIG. 5.— First plot: Radial density profiles through simulation time. The x-axis begins at ~ 1 AU, corresponding to the resolution limit of the simulation. The fit for the region outside of the disk is consistently $\rho \propto r^{-1.7}$ across all times. Second plot: Gas mass in a ball vs radius of the ball before the formation of the first sink. The fit is roughly $M \propto r^{1.3}$. Third plot: rate of change of the gas mass enclosed within a sphere of radius r , i.e. the integrated “mass of the ball” accretion rate $\dot{M}(< r) = dM(< r)/dt$. At later times the inner region gains mass slower than at the earlier times (mostly close to the time of the first sink forming). There is a characteristic “flattening” at smaller radii inside of the disk before the first star has formed. Outer shells gain mass faster at all times – a characteristic of “outside-in” collapse. Smallest shells have a drop in $\dot{M}(< r)$, associated with coming close to the resolution limit.




FIG. 6.— Radial density profiles at three different times before the first bulk star formation event at $t - t_1 \approx 1.2$ kyr. The density increases with decreasing radius, and follows a power law relationship that steepens with time.

subsequent evolution is governed by the competition between gravitational instability (which fragments the outer disk into secondary protostars), magnetic braking (which removes angular momentum and regulates the disk mass), and accretion-driven turbulence (which provides additional pressure support). We characterize this evolution through the disk energetics, its morphology, and its gravitational stability.

3.2.1. Energetics

In this section, we consider the interplay of the gravitational, kinetic, thermal and magnetic energies for the particles identified in the disk. We compute the following energy components: potential E_{pot} , rotational E_{rot} , turbulent E_{turb} , thermal E_{therm} , and magnetic E_{mag} , using Equations 5:

$$E_{\text{pot}} = -G \sum_{\substack{i < j \\ i, j \in \text{disk}}} \frac{m_i m_j}{\sqrt{|\mathbf{r}_i - \mathbf{r}_j|^2 + \epsilon^2}} \quad (5a)$$

$$- G \sum_{s \in \text{sinks}} \sum_{i \in \text{disk}} \frac{m_s m_i}{\sqrt{|\mathbf{r}_s - \mathbf{r}_i|^2 + \epsilon^2}},$$

$$E_{\text{rot}} = \frac{1}{2} \sum_{i \in \text{disk}} m_i v_{\phi, i}^2, \quad (5b)$$

$$E_{\text{turb}} = \frac{1}{2} \sum_{i \in \text{disk}} m_i |\mathbf{v}_i - v_{\phi, i} \hat{\phi} - v_{r, i} \hat{r}|^2, \quad (5c)$$

$$E_{\text{therm}} = \sum_{i \in \text{disk}} m_i u_i, \quad (5d)$$

$$E_{\text{mag}} = \sum_{i \in \text{disk}} \frac{|\mathbf{B}_i|^2}{8\pi} \frac{m_i}{\rho_i} \quad (5e)$$

where i denotes all gas particles identified as disk members (see [subsubsection 2.3.1](#)), m_i is the mass of particle, $\mathbf{v}_i$ is the total velocity vector, $v_{\phi, i}$, $v_{r, i}$ are bulk velocity profile fits for azimuthal and radial components for each annulus (both are functions of r only), $\mathbf{B}_i$ is the magnetic field vector, ρ_i is the density, u_i is the internal energy.

[section 4](#) shows the time evolution of the main energy components computed for disk-identified particles.

Throughout the simulation, $|E_{\text{pot}}|$ dominates over all other components, confirming that the disk is gravitationally bound. E_{rot} and E_{turb} are the main support terms and track each other to within a factor of ~ 2 , with E_{rot} slightly exceeding E_{turb} . E_{therm} starts comparable to the kinetic energies at early times but becomes subdominant at late times as the gravitational potential well deepens faster than the gas heats.

The magnetic energy, E_{mag} , grows over the ~ 15 kyr simulation, from $\sim 10^{42}$ erg at $t - t_1 \approx -0.5$ kyr to $\sim 10^{45.5}$ erg at 15 kyr. This growth is driven by a combination of flux-freezing compression during the collapse, field-line winding by differential rotation, and potentially the small-scale turbulent dynamo ([Sharda et al. 2021](#)). Despite this rapid amplification, E_{mag} remains below E_{therm} throughout the simulation, though the gap narrows at late times – consistent with the declining mass-to-flux ratios discussed in [subsection 3.3](#).

The bottom panel shows the virial ratio $2E_{\text{kin}}/|E_{\text{pot}}|$. At early times ($t - t_1 < 0$), the virial ratio is ~ 0.6 – 0.7 , indicating a *sub-virial*, contracting state, i.e. the system has insufficient kinetic energy to balance gravity.

The virial ratio rises as the disk forms and spins up, reaching ~ 1.0 by $t - t_1 \approx 2$ kyr. From 2 kyr onward, the virial ratio oscillates around unity, indicating that the disk has achieved and maintains approximate virial equilibrium through a balance of rotational support, turbulent pressure, and gravitational collapse. A dip in virial ratio after 11 kyr likely corresponds to the slow-down of stellar accretion rate (when less gas is accreted per unit time, less gravitational energy is converted to kinetic energy, reducing the kinetic-to-potential ratio), which can also be observed in [Figure 17](#). The “bounce-back” times also align with the “bounce-back” in the stellar accretion rate values.

The virial ratio never approaches the unbound threshold of 2, confirming that the system remains gravitationally bound throughout.

3.2.2. Disk morphology and evolution

As the disk accretes mass and angular momentum, its morphology undergoes a characteristic sequence: an initially smooth, centrally concentrated structure gives way to spiral arms as the Toomre parameter drops below 1, and those arms subsequently fragment into secondary sinks. The radial velocity structure simultaneously transitions from subsonic infall within the disk to supersonic infall in the surrounding envelope. This signifies a presence of the accretion shock at the disk boundary.

[Figure 7](#) shows radial profiles of velocity components of gas particles in the disk, as well as turbulent velocity magnitude and dispersion. We plot $-v_r$ (negated so that positive values indicate radially inward flow) to simplify comparison.

The general trend is that v_{ϕ} , $|\delta v|$, σ_r increase with time everywhere. v_r is negative and thus gas is mostly infalling for the whole simulation, with $|v_r|$ increasing until stabilizing at around 1.3–3.6 kyr (bulk star formation events), and dropping after stars are fully formed. c_s rises until 1.3 kyr, after which it stabilizes. It is clear that the turbulence starts off as sonic or sub-sonic, but becomes greatly supersonic once the sink forms. [The turbulent velocity dispersion increases inward, tracking the increase in gravitational potential. Because the collapse](#)




FIG. 7.— Radial profiles of inward radial velocity $-v_r$, azimuthal velocity v_ϕ , sound speed c_s , magnitude of turbulent velocity $|\delta v|$, radial/azimuthal/vertical turbulent velocity dispersion $\sigma_r/\sigma_\phi/\sigma_z$, Keplerian speed v_K of gas particles in the disks. Bright lines show functions with spline smoothing, dim lines show actual data. σ_r grows with time, as well as $|\delta v|$. Radial infall (v_r) speeds up until the bulk star formation events (1.3–3.6 kyr), and slows down afterwards. Disk rotation (v_ϕ) is constantly speeding up, leveling off outside of the disk ($r \gtrsim 2500$ AU). $\sigma_r, \sigma_\phi, \sigma_z$ track each other very closely, but σ_z decouples from the other two by the $\Delta t \gtrsim 9$ kyr time outside of the disk. Turbulence is supersonic at all times, tracking Keplerian speed v_K closely for most times outside of the disk, but falling below it inside of the disk.

compresses the gas faster than the turbulence can decay, the turbulence is continuously re-driven by the contraction itself rather than by stellar feedback (Robertson and Goldreich 2012); feedback only begins to contribute once the first protostar forms.

section 8 shows face-on and edge-on surface density projections at six representative epochs spanning the full ~ 15 kyr evolution after the formation of the first sink (t_1).

At the pre-stellar epoch ($t - t_1 = -0.5$ kyr), the gas is centrally concentrated without significant rotational support, appearing as an elongated, filamentary structure. By the time the first sink forms ($t - t_1 = 0$), a compact central concentration has developed. A coherent disk with recognisable rotational structure emerges within ~ 1 kyr. By $t - t_1 \approx 3.6$ kyr, the disk develops prominent two-armed spiral structure as it becomes gravitationally unstable (subsection 3.2.3), with multiple sink particles visible along the spiral arms. By $t - t_1 \approx 9.5$ kyr, the disk hosts ~ 20 visible sink particles embedded in a multi-armed, irregular spiral pattern extending to ~ 2000 AU, with several sinks located far from the centre. t

Radial density profile evolution is shown on Figure 6. It is clear that the density profile is following a steepening power law $\rho \propto r^{-\alpha}$, with $\alpha \approx 1.2$ at the time before the first sink forms, $\alpha \approx 1.8$ at the point of formation of the first sink, and $\alpha \approx 2.1$ by the time of a first bulk star formation event, at the point when the spiral arms have formed.

3.2.3. Disk stability and fragmentation

In this section, we analyze the stability properties of the disk. Whether a self-gravitating disk fragments into bound clumps, or instead disperses the accumulated mass through spiral-arm shocks and turbulent transport depends on the ratio of stabilising pressures (thermal, turbulent) to the destabilising self-gravity of the gas. The Toomre parameter, defined in Equation 6 (Toomre 1964), quantifies this balance locally: $Q < 1$ marks regions where perturbations on the disk scale grow faster than orbital shear can wind them up. We track the spatial and temporal evolution of Q to identify when and where fragmentation occurs, and how sink-driven velocity dispersion [SK: What does this term mean?] subsequently self-regulates the disk back toward marginal stability.

To characterize non-thermal motions in the disk, we decompose particle velocities into radial and azimuthal components (v_r, v_ϕ) in the disk-plane coordinate system. We subdivide the disk into concentric cylindrical annuli and compute the mass-weighted mean velocity in each annulus, yielding radial profiles $\bar{v}_r(r)$ and $\bar{v}_\phi(r)$.

The residual (streaming-subtracted) velocity for each particle is $\vec{\delta v} = (v_r - \bar{v}_r, v_\phi - \bar{v}_\phi, v_z)$. The velocity dispersion σ_v is computed as the root-mean-square of $|\vec{\delta v}|$ within each annulus.

The Toomre Q parameter quantifies the gravitational stability of a differentially rotating disk:

$$Q = \frac{c_s \kappa}{\pi G \Sigma}, \quad (6)$$

where c_s is the sound speed, κ the epicyclic frequency, and Σ the surface density.




FIG. 8.— Face-on and edge-on surface density projections at six epochs. Each panel shows a 4100×4100 AU region centred on the most massive sink. Star symbols mark sink particles within the image box. **The sinks are scattered throughout the disk rather than concentrated in a single cluster, tracing the spiral arms.**

For the local Toomre Q calculation we replace c_s with $\zeta = \sqrt{\sigma_r^2 + c_s^2}$, which includes both thermal and turbulent contributions, since supersonic turbulence provides additional effective pressure support.

For each snapshot the gas is binned into 20 annular bins **in the face-on frame**. In each annulus b we compute mass-weighted quantities:

$$\Sigma_b = \frac{\sum_{i \in b} m_i}{\pi(r_{\text{hi},b}^2 - r_{\text{lo},b}^2)}, \quad \sigma_{r,b} = \sqrt{\frac{\sum_{i \in b} m_i v_{r,i}^2}{\sum_{i \in b} m_i} - \bar{v}_{r,b}^2} \quad (7)$$

[SK: Make sure every single symbol is either has been described somewhere before the equation in the text or just after it. Like define $r_{\text{hi},b}$ here.] [SK: The formula looks wrong. The first term is the same as the second term, right?] The angular velocity is measured directly from the rotation profile:

$$\Omega_b = \frac{|\bar{v}_{\phi,b}|}{r_b} \quad (8)$$

where $\bar{v}_{\phi,b}$ is the mass-weighted mean azimuthal velocity in annulus b and r_b is the bin center. The epicyclic frequency is computed numerically from the measured

rotation curve via (Pringle and King 2007).

$$\kappa_b^2 = \frac{2\Omega_b}{r_b} \left. \frac{d(r^2\Omega)}{dr} \right|_{r_b} \quad (9)$$

The Toomre parameter per annulus is then

$$Q_b = \frac{\zeta_b \kappa_b}{\pi G \Sigma_b} \quad (10)$$

A pixel-by-pixel face-on Q map is produced by computing local mass-weighted quantities for all of the relevant parameters:

$$Q(x,y) = \frac{\zeta_b(x,y) \kappa(r_{x,y})}{\pi G \Sigma(x,y)} \quad (11)$$

where $r_{x,y} = \sqrt{x^2 + y^2}$, and $\zeta = \sqrt{\sigma_r^2 + c_s^2}$, which sums over thermal and turbulent contributions.

~~The Toomre Q parameter measures the gravitational stability of the disk.~~ Figure 9 (top) shows face-on maps of the local Q , computed using the streaming-subtracted velocity dispersion σ_v in place of c_s [SK: Why did you do this? They are comparable in magnitude right?]. At the pre-stellar epoch ($t - t_1 = -0.5$ kyr), the gas is




FIG. 9.— Top: Face-on maps of the local Toomre Q parameter at six epochs. The black contour marks $Q = 1$. Bottom: Evolution of radial profile of Toomre Q (heatmap), and sink formations, mergers, and radial positions' evolution (lines). At early times the gas is globally unstable ($Q < 1$); at late times the inner disk is strongly stabilised while the outer spiral arms remain near or below $Q = 1$. Radial profile of Toomre Q at six epochs. The horizontal dashed line marks $Q = 1$. At early times the entire region has $Q < 1$; at late times a radial gradient develops, with $Q > 10^3$ at $r < 100$ AU falling to $Q \sim 1$ at $r \sim 2000$ AU. [SK: I would increase the size of the sink to be the same as the color of the line? Maybe even make some of the lines thicker according to the mass of the star. It looks like some of the lines are well correlated with the Toomre- Q evolution – I wonder if the more massive stars are responsible for this.]

broadly unstable with $Q \lesssim 1$ throughout most of the region. By $t - t_1 = 0$, the central region near the first sink begins to stabilise, while the surroundings remain marginally unstable. At $t - t_1 \approx 1.3$ kyr, a clear boundary emerges between a stabilised inner region and an unstable outer envelope. By the time spiral arms have developed ($t - t_1 \approx 3.6$ kyr), a ring of $Q < 1$ material traces the spiral arms, while the inner disk is stabilised ($Q > 1$) by the strong velocity dispersion generated near the central sinks.

The radial Toomre Q profile (Figure 10) quantifies its evolution. At the pre-stellar epoch ($t - t_1 = -0.5$ kyr), Q is below unity at all radii ($Q \sim 0.2-0.3$), reflecting a globally unstable collapsing core. As the first sink forms and accretes, the inner Q rises: by $t - t_1 = 1.3$ kyr (green), $Q \approx 1$ throughout, consistent with a marginally stable

state. By $t - t_1 = 9.5$ kyr (brown), Q is mostly stable in the whole disk, with the central region [SK: quantify in size] being the most stable. At later times a radial gradient develops, with the inner disk more stable ($Q \gg 1$) and the outer spiral arms remaining near or below $Q \sim 1$.

Figure 9 (bottom) shows the space-time evolution of the azimuthally-averaged Toomre Q radial profile. Lines on the plot correspond to evolution of radial positions of sinks ($r(t)$). The disk begins in a predominantly unstable state ($Q < 1$, red) at all radii. The first sink forms at the centre, and within ~ 1 kyr the $Q = 1$ contour (black line) begins to propagate outward from the centre, tracing a wave of stabilisation driven by the velocity dispersion generated during the initial collapse. A second stabilisation wave is launched by the formation of a




FIG. 10.— Toomre Q radial profiles time evolution. The disk starts with $Q < 1$ inside and $Q \approx 1$ near the edge, but Q rises with disk evolution, since gas is removed from the disk because of star formation events. Center of the disk is extremely stable ($Q \gtrsim 10^2$) by the 15 kyr mark, preventing any further star formation in the middle. [SK: I don't think this plot adds any new information, you can get rid of it to keep things more concise.]

second sink near the centre at $\Delta t \approx 1$ kyr. These waves propagate outward at speeds of $\sim 2-3 c_s$.

[SK: You commented out what is really driving these waves, if it's surface density or velocity dispersion or both. If you have a heatmap it might be fine to add it here. Otherwise you can also check what is driving this and just state it without showing a plot.]

At $t - t_1 \approx 3$ kyr, a spiral arm becomes unstable and fragments into [SK: Add how many sinks form here] closely-spaced sink particles within ≈ 1 kyr. However, several of these sinks merge shortly thereafter (see Grudić et al. 2021, for details on the merging criteria), leaving behind XYZ stars. This conversion of gas into stars reduces the gas mass in the disk and stabilizes it.

3.3. Magnetic fields

In this section, we analyze the evolution of the magnetic field as the disk collapses and fragments. We find that the magnetic fields are amplified during the collapse by compression along field lines and by differential rotation within the disk once it forms. The two relevant amplification channels – flux freezing (compression-driven, $B \propto \rho^{2/3}$) and rotational winding (differential-rotation-driven, $B_\phi \propto r \partial \Omega / \partial r \cdot B_z$ [SK: The units don't look right, maybe you mean $\text{dB}_\phi / \text{dt}$?]) – leave distinguishable imprints in the spatial structure and scaling of the field. We characterize the resulting field morphology and its evolution through the vertical (B_z) and toroidal (B_ϕ) components. ~~and we quantify the coupling between matter and field through the mass-to-flux ratio $\mu = M/\Phi_B$~~ $\mu = M/M_\Phi$ normalized to the critical value. [SK: Don't define stuff if you don't need it yet. Also whenever you write an equation, make sure the symbols have been defined either before the equation or just below it.]

Figures 11–12 show face-on ($z = 0$ slice) and edge-on ($x = 0$ slice) views of the toroidal and vertical components of the magnetic field at six epochs. [SK: A sentence on its own can technically be a paragraph, but shouldn't be one here. Why did you make slices instead of projec-

tions? Slices are more prone to stochastic effects. You could make this a projected average of the B field components.]

At the pre-stellar epoch, B_z shows large-scale coherent structure with a preferred sign, pointing towards primordial field orientation. As the disk forms and develops spiral structure, B_z is amplified in the disk midplane and develops sign reversals that trace the spiral arms, with field strengths reaching $|B_z| \sim 10^{-2}-10^{-1}$ G at late times [SK: Why does the field change sign according to spiral arms?].

The toroidal component B_ϕ (Figure 11) develops an antisymmetric pattern by $t - t_1 \approx 1.3$ kyr: the field exhibits opposite signs on either side of the disk midplane, as confirmed by the edge-on view. This field can be achieved if we assume that primordial B-field was directed in $+x +y$ direction diagonally. [SK: I don't understand this last sentence and I don't understand why it should be opposite directions on either side of the disk midplane.] By $t - t_1 \approx 3.6$ kyr, B_ϕ displays a clear two-armed spiral pattern with alternating positive and negative regions, reaching strengths of $|B_\phi| \sim 10^{-1}$ G. This antisymmetric, wound-up morphology is characteristic of field amplification by differential rotation and is consistent with a large-scale disk dynamo. By late times (15 kyr,) the toroidal component of the field is directed mostly opposite with respect to the direction of rotation of the disk.

Why did the toroidal field end up being anti-parallel to the direction of rotation? A clue might be located in the fourth face-on plot, corresponding to $\Delta t \approx 3.6$ kyr, roughly the time of first bulk star formation event. It is clear that at this state, the spatial distribution of the toroidal field is evenly distributed between positive ($B_{\text{tor}} > 0$, red) and negative ($B_{\text{tor}} < 0$, blue) components, with each of two of spiral arms corresponding to one of them. However, the spiral arm with denser gas corresponds to $B_{\text{tor}} < 0$ regime, and the magnetic field is amplified in there during collapse via flux-freezing mechanism. At later times this amplification is dispersed around the whole disk region, making $B_{\text{tor}} < 0$ regime dominant. [SK: I don't get what you're trying to say here in this paragraph, let's discuss this in our next meeting.]

[SK: You should always discuss the figures in order. Right now 14 is discussed before 13.]

We define mass-to-flux ration per annulus b as:

$$\mu_{\Phi,b} = \frac{2\pi\sqrt{G}\Sigma_b}{|B_{z,b}|} \quad (12)$$

The mass-to-flux ratio μ_Φ (Figure 14) evolves significantly over time.

At early times ($t - t_1 \leq 1.3$ kyr), $\mu_\Phi \sim 100-300$ at all radii, indicating an extremely magnetically supercritical state in which gravity overwhelms magnetic support by more than two orders of magnitude. The mass-to-flux ratio then declines steadily as the field is amplified by compression and winding: by $t - t_1 = 3.6$ kyr (red), μ_Φ has dropped to $\sim 10-100$. By $t - t_1 = 6.3$ kyr (purple), μ_Φ exhibits large radial scatter, with values ranging from ~ 1 to ~ 100 and dipping below the critical value ($\mu_\Phi < 1$) at some inner radii. At the latest epoch ($t - t_1 = 9.5$ kyr, brown), significant portions of the inner disk ($r \sim 200-500$ AU) have $\mu_\Phi < 1$, indicating that the disk has

become locally *magnetically subcritical* – a state where magnetic pressure and tension can in principle support the gas against gravitational collapse.

This transition from $\mu_\Phi \sim 300$ to $\mu_\Phi \lesssim 1$ over ~ 10 kyr represents a decrease of more than two orders of magnitude and implies a corresponding amplification of the magnetic field by flux-freezing compression, differential rotation winding, and possibly the turbulent dynamo.

Figure 13 shows evolution of phase plot of magnetic field magnitude $|B|$ through the 6 epochs we have been discussing. There is an evolving power law relationship that evolves as $|B| \propto n^{2/3} \rightarrow |B| \propto n^1$.

[TODO: More explanations needed. What does this mean? Do we resolve dynamo?]

[SK: I will comment once you've added this stuff here]

3.4. Star Formation and the IMF

The rate at which gas is converted into stars is set by the competition between the accretion luminosity (which heats the disk and raises the Jeans mass, suppressing further fragmentation) and the gravitational instability of the disk (which drives fragmentation and seeds new protostars). The proto-stellar mass function (PMF) encodes the integrated outcome of these competing processes at the epoch of observation. We quantify the total stellar mass growth, the instantaneous star formation efficiency within the disk, and the shape of the PMF to constrain which physical mechanism dominates the mass budget.

Figure 15 shows the time evolution of gas, stellar (sink), and total mass within a 0.1 pc aperture centred on the disk.

The total stellar mass (orange, all sinks combined) grows from $\sim 1 M_\odot$ at $t - t_1 = 1$ kyr to $\sim 200 M_\odot$ by 15 kyr, with a best-fit power law $M_* \propto (t - t_0)^{2.08}$ (dashed red line). This is consistent with the $M_* \propto t^2$ scaling predicted by models of accelerating accretion from a turbulent collapsing core (McKee and Tan 2003). Murray et al. (2017) confirmed this scaling in high-resolution AMR simulations of self-gravitating turbulent collapse, finding that the density inside the stellar sphere of influence evolves to a time-independent attractor $\rho(r, t) \rightarrow \rho(r) \propto r^{-3/2}$, which, combined with Keplerian infall velocities, yields a mass accretion rate $\dot{M} \propto t$ and thus $M_* \propto t^2$. Our measured exponent of 2.08 extends the Murray et al. (2017) result from parsec-scale molecular cloud collapse to the sub-AU scales of Population III accretion disks.

The disk gas mass (cyan solid) starts at $\sim 50 M_\odot$ and remains roughly constant for the first ~ 3 kyr, as infall from the envelope replenishes gas consumed by sink accretion. At later times, the disk gas mass begins to decline as the star formation efficiency rises. The gas mass within the 0.1 pc aperture (cyan dashed) is approximately constant at $\sim 200 M_\odot$ throughout the simulation, yet its evolution follows one for the disk gas mass up until $\Delta \approx 11$ kyr, after which we see an increase in aperture gas mass, while disk gas mass stays constant. The reason for this is probably a large amount of infalling gas suddenly entering the aperture, resulting in a positive $\dot{M}_{gas,apert}$ (i.e. more gas is entering the system than is being converted to stellar mass). This sudden rise in the gas mass also “ignites” the stellar accretion again after its slow-down, driven by the depletion of initial accessi-

ble reservoirs in the original disk, and one can see this “bounce-back” in Figure 17. In addition to the increased accretion rates, we also see that the gas depletion rate ($-\dot{M}_{gas,apert}$) is not on the plot, which is the case since the total amount of gas is increasing, thus the gas depletion rate becoming negative right at the time of the “bounce-back”.

The lower panel of Figure 15 shows the star formation efficiency (SFE) $f_* = M_*/(M_{gas} + M_*)$. Within the disk alone, f_* rises steeply from $\sim 2\%$ at 1 kyr to $\sim 80\%$ by 10 kyr. Within the 0.1 pc aperture, f_* reaches $\sim 50\%$ by 10 kyr. These are high efficiencies compared to present-day star formation, but it is also not unexpected, since we did not pick an object, similar to a $z = 0$ giant molecular cloud (GMC) to compare our results with. Even our aperture will have less gas mass than present-day GMC, thus resulting in higher SFE.

Figure 17 shows the corresponding mass rates. The stellar accretion rate (magenta) fluctuates around $\sim 5 \times 10^{-3} M_\odot \text{ yr}^{-1}$ at early times, rising to $\sim 10^{-2} M_\odot \text{ yr}^{-1}$ at late times with strong variability. These rates are consistent with the canonical Population III accretion rate $\dot{M} \sim c_s^3/G \sim 10^{-3} M_\odot \text{ yr}^{-1}$ set by the sound speed in $\sim 200\text{--}1000$ K gas (Abel et al. 2002; Stacy et al. 2010). The lower panel shows the sphere of influence radius r_{SOR} , defined as the radius where the enclosed gas mass equals the total stellar mass ($M_{gas,enc} = M_*$). This grows from ~ 80 AU at 1 kyr to $\sim 10^4$ AU by 10 kyr. It drops slightly with the new gas entering the aperture.

3.4.1. Initial mass function

Figure 19 shows the initial mass function (IMF) of the sinks at $t - t_1 = 15$ kyr, compared to the present-day Kroupa (2001) and Chabrier (2003) initial mass functions (IMFs). The high-mass slope of the IMF is well fitted by a power law $dN/dM \propto M^\alpha$ with $\alpha = -0.99^{+0.14}_{-0.14}$, shallower than Salpeter slope ($\alpha = -2.35$) and the Kroupa high-mass slope ($\alpha = -2.3$).

Figure 16 shows integrated accretion history of individual sinks. We can clearly see that 2 of the heaviest sinks are the first 2 sinks that formed in the simulation, and are located in the center of the disk. A lot of low-mass ($< 3 M_\odot$) sinks stop accreting relatively early because they get ejected out of the disk, thus limiting the amount of gas, available to them. **[Can you show this quantitatively or show a plot of this?]** In our simulation, we have seen 14 out of 32 present sinks being ejected out of the disk by the 15 kyr time. With 62 total sink particles that were observed during the whole simulation time, the fraction of $\sim 52\%$ sinks being merged and 23% of sinks being ejected from the disk is consistent with values, expected in the literature (compared to 50% and 25% respectively of (Stacy and Bromm 2013), or 30% of ejections of (Wollenberg et al. 2020)).

At the same figure it can be seen that the accretion in the center of the disk (heaviest 2 sinks) mostly halts, which is a consequence of depletion of gas reservoir. This gas depletion is shown as a drop in stellar accretion rate in Figure 17. The “bounce-back”, described in detail in subsection 3.4, seems to mostly target the non-central sinks, since we still observe their growth after $\Delta t \approx 11$ kyr. This competitive accretion in the cluster environment may preferentially feed the most massive




FIG. 11.— Face-on ($z = 0$ slice) and edge-on ($x = 0$ slice) of the toroidal magnetic field component B_ϕ at six epochs.

objects (Bonnell et al. 2001), flattening the IMF slope. **[How do you know it is competitive accretion vs turbulent fragmentation?]**

4. COMPARISON TO PREVIOUS POPULATION III SIMULATIONS

We observe a gravitationally unstable, magnetized disk that fragments prolifically and accretes at $\dot{M} \sim 10^{-3} - 10^{-2} M_\odot \text{ yr}^{-1}$. This picture is consistent with the broad expectations of Population III star formation theory, but the combination of the regime of cosmological initial conditions, resolved protostellar sinks, radiation-MHD, and 15 kyr of post-collapse evolution exceeds what any single prior study has achieved simultaneously.

Our simulation combines three features that, taken together, are uncommon in the existing literature: fully cosmological initial conditions, radiation-MHD with resolved individual protostellar formation (STARFORGE), and 15 kyr of post-collapse evolution in a halo progenitor of a Milky Way-mass system. Below we compare our results in each dimension to prior work.

4.1. Fragmentation and sink counts

The total number of protostellar sinks formed in a simulation is a coarse but widely reported diagnostic of disk fragmentation: it captures the net outcome of the com-

petition between gravitational instability (which seeds new fragments) and their subsequent mergers, ejections, and accretion onto the primary. Comparing sink counts across simulations is complicated by the spread in integration times and sink radii, but it nevertheless reveals whether a given numerical setup is producing qualitatively more or less fragmentation than the community baseline.

The ~ 60 sink formation events we observe over 15 kyr place our simulation at the high end of the published range. Early simulations with cosmological initial conditions found 2–5 fragments per halo within the first few hundred years (Greif et al. 2011; Clark et al. 2011), while longer integrations over ~ 5 kyr recovered ~ 10 sinks (Stacy et al. 2012). The radiation-hydrodynamic cosmological zoom-ins of Latif and Khochfar (2022) across five minihaloes produced up to 23 stars per halo (55 total), with masses $0.1\text{--}37 M_\odot$ and a slope $dN/dM \propto M^{-1.18}$. The systematic study of Wollenberg et al. (2020) with 45 controlled simulations (varying rotation and turbulence) found 5–20 sinks per halo, confirming that fragmentation is highly stochastic and sensitive to initial conditions.

Our fragment count likely reflects three compounding factors. First, our 15 kyr integration is substantially longer than most published runs, allowing many cycles of instability, collapse, and re-accretion. Second, cosmologi-




FIG. 12.— Face-on ($z = 0$ slice) and edge-on ($x = 0$ slice) of the vertical magnetic field component B_z at six epochs. By the late epochs shown here, most of the stars in the simulation have already formed, so the field structure reflects the post-formation disk rather than the initial collapse.




FIG. 13.— Evolution of phase plot of magnitude of magnetic field $|B|$ vs number density n . Solid lines correspond to median profiles at different times, with the shaded regions being 90-th percentile of all the particles. α corresponds to the exponent in power-law fit $|B| \propto n^\alpha$. $|B|$ spans 3 orders of magnitude at the point of first sink formation, and up to 4 orders of magnitude at later times. [SK: Fix this figure]




FIG. 14.— Dimensionless mass-to-flux ratio μ_Φ as a function of radius at six epochs. The dashed line marks the critical value $\mu_\Phi = 1$. The disk evolves from highly supercritical ($\mu_\Phi \sim 100-300$) at early times to partially subcritical ($\mu_\Phi < 1$) at some inner radii by 9.5 kyr.

ical initial conditions from the m12f progenitor provide asymmetric, turbulent infall that continually replenishes the outer disk and seeds new unstable structures. Third,




FIG. 15.— Top: time evolution of the stellar mass (all sinks; orange), disk gas mass (cyan solid), gas within a 0.1 pc aperture (cyan dashed), and total mass (disk + stars, grey solid; aperture + stars, grey). The best-fit power law $M_* \propto t^{2.08}$ is shown as a dashed red line; the reference t^2 scaling as a dotted pink line. Bottom: star formation efficiency f_* within the disk (green solid) and within the 0.1 pc aperture (green dashed).




FIG. 16.— Evolution of masses of individual sinks. Dashed line represents ideal $M_* \propto t^2$ law, which is being followed for total mass of all the sinks (black line). At earlier times, before the first bulk star formation event at 2–3 kyr, all of stellar mass is concentrated in the 2 central sinks (top blue lines). These massive sinks reach mass of around $20M_\odot$ by the 15 kyr mark, with their accretion slowing down compared to less massive sinks, which still accrete rapidly.




FIG. 17.— Top: stellar accretion rate (magenta) and gas depletion rate within the aperture (cyan dashed). Bottom: sphere of influence radius r_{SOI} ($M_{\text{gas,enc}} = M_*$; orange), growing from ~ 80 AU to $\sim 10^4$ AU over the simulation.




FIG. 18.— Time evolution of the number of sinks in the simulation. The dashed line shows a $N \propto t^{1.5}$ power law for comparison (Li et al. 2018). Bulk star formation events in the spiral arms (visible in the bottom panels of Figure 9) are seen to rapidly increase the number of sinks at $t - t_1 \approx 3$ –4 kyr and 7–8 kyr.




FIG. 19.— Protostellar mass function at $t - t_1 = 15$ kyr (blue solid histogram) compared to the Chabrier (2003) (dashed) and Kroupa (2001) (dotted) present-day IMFs. The best-fit high-mass power-law slope is $\alpha = -0.99^{+0.14}_{-0.14}$ (light blue dashed line), shallower than but comparable to the Salpeter slope (-2.35). The grey shaded region marks masses below the completeness limit of the simulation. As this is a initial mass function measured at 15 kyr, the final masses will be modified by continued accretion, mergers, ejections, and radiative feedback.

our mass-to-flux ratio remains comfortably supercritical ($\mu_\Phi \gg 1$) for most of the simulation, so magnetic fields do not suppress fragmentation here as they do in some studies (Stacy et al. 2022; Machida et al. 2025). The high fragment count is not unprecedented: Latif and Khochfar (2022) obtained comparable numbers with RHD (no MHD), and Prole et al. (2022) showed that sink counts increase monotonically with resolution without converging, suggesting our result may be a lower bound. Of the 62 total sink formation events, 32 survive to 15 kyr; the remainder were lost to mergers ($\sim 52\%$ of all events) or ejected from the disk ($\sim 23\%$), consistent with the rates reported by Stacy et al. (2012) and Wollenberg et al. (2020).

4.2. Role of the large-scale environment

The properties of the first collapsing core are set by the large-scale accretion history of the host minihalo, which in turn depends on where that halo sits within the cosmic web. A halo in an overdense filament accretes faster and with more angular momentum than an isolated one, biasing the collapse toward a more massive, rotationally supported structure. Because our halo is a progenitor of a Milky Way-mass system, its large-scale environment is atypical of the field minihalos studied in most prior work.

At the epoch of first collapse ($z \sim 21$), our m12f progenitor is a $\sim 10^5$ – $10^6 M_\odot$ minihalo – comparable in mass to those studied in most other work – but it is embedded in a large-scale overdensity destined to grow to $M_{\text{vir}} \sim 10^{12} M_\odot$ by $z = 0$. The most directly comparable simulation, Meziani et al. (2026), targets the m09 halo whose progenitor at $z \sim 14$ resides in a quieter environment (reaching $\sim 2 \times 10^9 M_\odot$ at $z = 0$), and finds a single Population III protostar growing to $\sim 27 M_\odot$ over 31 kyr – in stark contrast to our ~ 60 -fragment cluster. Although this comparison is a single pair of halos, the contrast suggests that large-scale environment may be a key driver of whether Population III star formation pro-

duces a single massive object or a protostellar cluster. The overdense filamentary infall geometry of the m12f progenitor naturally seeds higher angular momentum, more complex disk structure, and stronger turbulence than the more isolated m09 environment. Chen et al. (2025) showed that infall into dark-matter potential wells drives supersonic turbulence ($\mathcal{M} \sim 2$ – 4) across all minihaloes; the m12f overdensity likely places our halo toward the high- $\mathcal{M}$ end of this distribution. Testing this environmental dependence statistically will require a suite of cosmological RMHD zoom-ins spanning a range of $z = 0$ halo masses. The kinematic structure of our simulation supports this picture: the radial velocity v_r is subsonic within the disk but transitions to supersonic outside it, indicating that the outer envelope is in a state of continuous supersonic infall that feeds the disk from above.

4.3. Stellar mass growth and accretion rates

The total stellar mass assembled by a given time is the integral of the disk accretion rate over all active sinks, so it reflects both the efficiency of angular momentum transport (which determines how quickly gas drains inward through the disk) and the number of competing accretion regions (which determines how the common reservoir is partitioned among the protostars). A large sink count spreading the accretion over many objects can produce a similar total mass to a single rapidly accreting protostar while leaving a completely different PMF. We compare our integrated mass growth and accretion rates to prior work to disentangle these effects.

The total stellar mass of $\sim 200 M_\odot$ at 15 kyr is broadly consistent with canonical Population III accretion rates. Stacy et al. (2010) found $\dot{M} \sim 10^{-3} M_\odot \text{ yr}^{-1}$ sustained over ~ 5 kyr; integrating this rate over 15 kyr gives $\sim 15 M_\odot$, far less than we find, reflecting our larger sink count sharing a common reservoir rather than a single accreting object. Our time-averaged rate of $\sim 10^{-2} M_\odot \text{ yr}^{-1}$ (across all sinks) is at the high end of the range 10^{-4} – $10^{-2} M_\odot \text{ yr}^{-1}$ found by Hirano et al. (2014) across 100 minihaloes, again consistent with a particularly turbulent, high-infall-rate environment.

The power-law growth $M_* \propto t^2$ confirms $M_* \propto t^{2.08}$ confirms the $M_* \propto t^2$ scaling found by Murray et al. (2017) in simulations of self-gravitating turbulent collapse, where the density inside the sphere of influence converges to $\rho \propto r^{-3/2}$ and Keplerian infall gives $\dot{M} \propto t$. Our slightly super-quadratic exponent likely reflects the additional contribution of continued cosmological infall sustaining the gas reservoir at large radii, absent in the isolated box setup of Murray et al. (2017). The POPSI-CLE simulations of Sharda and Menon (2025) produce a single most-massive protostar of $\sim 65 M_\odot$ within 5 kyr in their RMHD run; our total mass of $\sim 200 M_\odot$ is distributed across ~ 30 surviving sinks at 15 kyr, implying a characteristic fragment mass of ~ 5 – $7 M_\odot$, consistent with fragmentation-induced starvation redistributing mass from the most massive object to the cluster population (Prole et al. 2022). Sugimura et al. (2020) and Sugimura et al. (2023) produced massive Population III binaries with individual masses $> 30 M_\odot$ at separations > 2000 AU; the two most massive sinks in our simulation – which dominate the mass budget throughout – may represent an analogous central binary, with the sur-

rounding cluster population arising from subsequent disk fragmentation. A key qualitative difference from these earlier studies is that our protostars form within a single, large-scale disk rather than in small, circum-protostellar disks. The disk extends to ~ 2500 AU and fragments repeatedly, so the entire protostellar population shares a common gas reservoir and competes for the same infalling material. This geometry naturally accounts for the large number of sinks and for the accelerating, super-quadratic stellar-mass growth we report.

4.4. Star formation efficiency

Star formation efficiency ($f_* = M_*/M_{\text{gas}}$) measures the fraction of the available gas reservoir that has been converted into stars by a given epoch. In primordial gas, the main mechanisms that would suppress f_* in present-day molecular clouds (dust-driven radiation pressure and stellar winds) are absent or greatly weakened, so the efficiency is expected to be high. However, it is not clear how high exactly. Photoionization and accretion luminosity heating remain active, and whether they can quench accretion before the disk is exhausted determines the ultimate yield.

The star formation efficiency we find ($f_* \sim 50\text{--}80\%$ within the disk at 15 kyr) is high compared to both present-day values ($\sim 1\text{--}10\%$) and to other Population III simulations. Hirano et al. (2023) report $\epsilon_{\text{III}} \sim 10^{-2}$ in their large halo sample at the point of initial collapse; our higher integrated value reflects the longer post-collapse integration time and the lack of negative feedback mechanisms that would quench accretion. Radiative feedback is included in our simulation (M1 radiation transport), but within the 15 kyr of evolution covered here no protostar has yet grown sufficiently massive and contracted sufficiently to launch an ionising HII region that would reverse the accretion flow. Hosokawa et al. (2011) found that UV feedback halts accretion at $\sim 43 M_\odot$ for a single protostar on a $\sim 10^5$ yr Kelvin-Helmholtz timescale; with our mass distributed across many objects, no individual protostar yet dominates the radiative output, and the global accretion efficiency remains high. This suggests that the high SFE we observe is a transient feature of the early accretion phase; longer integrations incorporating protostellar contraction and UV feedback would be required to determine final stellar masses and the termination of star formation. **Jets are not present in the current run: a bug in the launch implementation suppresses them (see section 2), so we cannot assess their impact on the star formation efficiency here. Once jets are enabled, they are expected to become progressively more important as individual protostars contract and their surface escape velocities increase, potentially quenching accretion and reducing the SFE from its current high value (Grudić et al. 2021; Meziani et al. 2026).**

4.5. Protostellar mass function

The slope and peak of the protostellar mass function encode the dominant mass-setting mechanism: competitive accretion (which weights the PMF toward high masses with $dN/d\ln M \propto M^{-1}$), thermal Jeans fragmentation (which sets the peak near the local Jeans mass and produces a roughly log-normal distribution), and protostellar feedback (which imposes an upper cutoff

wherever radiation pressure or HII regions terminate accretion). The top-heavy, log-flat PMF observed in most Population III simulations points toward a regime dominated by competitive accretion and absent of the feedback mechanisms that steepen the present-day IMF. We compare our slope and peak to the literature to assess how robustly this signature is recovered across different numerical methods and initial conditions.

Our protostellar mass function (PMF) at 15 kyr has a high-mass slope $\alpha = -0.99^{+0.14}_{-0.14}$, shallower than the Salpeter slope (-2.35) but broadly consistent with other Population III simulations that find top-heavy, log-flat distributions. Latif and Khochfar (2022) find $\alpha \approx -1.18$ across five halos with RHD; Stacy et al. (2016) and Greif et al. (2011) both report roughly log-flat mass functions extending from sub-solar to $\gtrsim 10 M_\odot$. We emphasise that our PMF is a snapshot at 15 kyr, not a final IMF: all sinks are still actively accreting, no radiative feedback has terminated accretion, and the population will be further shaped by continued mergers, ejections, and eventual UV feedback (Hosokawa et al. 2011, 2016; Stacy et al. 2016). The two most massive sinks, both located at the center of the disk, have each reached $\sim 25 M_\odot$ by 15 kyr – approaching the $\sim 43 M_\odot$ UV-feedback threshold of Hosokawa et al. (2011) – with their accretion rates flattening due to local gas depletion in the inner disk. The outer sinks, by contrast, continue to grow by accreting from the surrounding envelope, so the cluster mass budget shifts progressively outward. The $\sim 52\%$ merger fraction and $\sim 23\%$ ejection fraction we observe are consistent with the $\sim 50\%$ and $\sim 25\%$ values of Stacy et al. (2012) and the $\sim 30\%$ ejection rate of Wollenberg et al. (2020). Latif and Khochfar (2022) report an unusually high ejection fraction of $\sim 70\%$, which they attribute to the strong three-body interactions driven by the radiative heating of the disk; the lower fraction in our simulation may reflect the slower build-up of radiation pressure in a multi-sink cluster where no single object yet dominates.

4.6. Magnetic field evolution

The amplification of primordial seed fields during the collapse and disk phase is one of the most uncertain aspects of Population III star formation: the initial field strength is poorly constrained, and the relevant dynamo mechanisms (turbulent small-scale, large-scale differential rotation) operate on overlapping timescales. We compare the evolution of our mass-to-flux ratio and magnetic energy to the MHD simulations that do resolve the disk phase, to assess whether the rapid amplification we observe is generic or a consequence of our particular choice of seed field and numerical method.

Our mass-to-flux ratio evolves from $\mu_\Phi \sim 100\text{--}300$ at early times to $\mu_\Phi \lesssim 1$ in the inner disk by 10 kyr, with magnetic energy growing by ~ 4 orders of magnitude. This evolution is more dramatic than the $\mu_\Phi \sim 10\text{--}100$ reported by Machida et al. (2008) in idealised MHD simulations, likely because our longer integration time allows more flux accumulation by differential rotation winding and compression. Higashi et al. (2024) demonstrated that the turbulent small-scale dynamo amplifies fields to 30–50% of equipartition during the primordial collapse phase; our highly supercritical starting point and the continued amplification we observe are consistent with

this picture.

Despite this amplification, disk fragmentation in our simulation remains vigorous throughout, in contrast to the strong suppression found in some MHD simulations. Machida et al. (2025) systematically varied initial field strength from $B_0 = 10^{-20}$ to 10^{-4} G and found that even the weakest field suppresses fragmentation and drives protostellar mergers within the first ~ 1000 yr – a timescale much shorter than our run. Stacy et al. (2022) found only two secondary sinks in cosmological MHD runs. The key difference is that our simulation starts from a highly supercritical configuration ($\mu_\Phi \sim 300$) and remains in the fragmentation-permitting regime ($\mu_\Phi > 1$) for the majority of the 15 kyr. Magnetic braking and flux-freezing act on a timescale long compared to the free-fall time at the disk scale, so fragmentation proceeds largely unimpeded while the field is amplified gradually from below. Only toward the end of our simulation does the inner disk approach the magnetically subcritical threshold; at this point, Meziani et al. (2026) find that protostellar jets are launched, a phase not yet reached in our 15 kyr window. The magnetic energy power spectrum is consistent with a Burgers spectrum ($P_B \propto k^{-2}$) at most times, oscillating between Burgers and Kolmogorov regimes but remaining closer to Burgers, indicating shock-dominated rather than purely subsonic turbulent amplification. A detailed analysis of the magnetic power spectrum and its implications for dynamo action in primordial disks is deferred to a companion paper. In contrast, Sadanari et al. (2021) and Sadanari et al. (2023) found that ambipolar diffusion is dynamically subdominant in primordial gas (diffusion heating always smaller than compressional heating), broadly consistent with our non-ideal MHD treatment where ideal MHD dominates throughout (Meziani et al. 2026). Sadanari et al. (2024) similarly found in primordial disk simulations that turbulent magnetic fields reduce but do not eliminate fragmentation, consistent with our result.

5. CONCLUSIONS

We have presented a radiation-magnetohydrodynamic cosmological zoom-in simulation of Population III star

formation, following the collapse of primordial gas in the FIRE m12f halo from cosmological initial conditions through the formation and evolution of a protostellar accretion disk over ~ 15 kyr. Our main conclusions are:

1. **Vigorous disk fragmentation.** The accretion disk becomes gravitationally unstable within ~ 3 kyr of the first protostar’s formation, developing spiral arms that fragment into ~ 60 sink formation events over 15 kyr, producing a total stellar mass of $> 150 M_\odot$.
2. **Magnetic field amplification toward subcriticality.** The mass-to-flux ratio drops from $\mu_\Phi \sim 100\text{--}300$ at early times to $\mu_\Phi \lesssim 1$ at some inner radii by 10 kyr, driven by flux-freezing, differential rotation winding, and possibly the turbulent dynamo. Magnetic energy grows by ~ 4 orders of magnitude. The inner disk eventually crosses the magnetically subcritical threshold.
3. **Sub-virial start to virial equilibrium.** The disk starts in a sub-virial state ($2E_{\text{kin}}/|E_{\text{pot}}| \approx 0.6$) and reaches virial equilibrium within ~ 2 kyr, thereafter self-regulating around $2E_{\text{kin}}/|E_{\text{pot}}| \approx 1$.
4. **Accelerating accretion and high SFE.** The stellar mass grows as $M_* \propto t^2$ at accretion rates of $\sim 10^{-3}\text{--}10^{-2} M_\odot \text{ yr}^{-1}$. The star formation efficiency reaches $\sim 80\%$ within the disk and $\sim 50\%$ within a 0.1 pc aperture.
5. **Top-heavy initial mass function.** The initial mass function has a high-mass slope $\alpha = -0.99^{+0.14}_{-0.14}$, shallower than the Salpeter/Kroupa value ($-2.35/-2.3$).
6. **Cosmological ICs matter.** The use of cosmological initial conditions produces asymmetric infall, complex angular momentum structure, and turbulent boundary conditions that promote vigorous fragmentation and a rich protostellar population, distinct from what is found in idealised setups. Continued accretion from LSS onto the protostellar disk modifies the IMF, flattening the slope.

REFERENCES

- T. Abel, P. Anninos, Y. Zhang, and M. L. Norman. Modeling primordial gas in numerical cosmology. *New Astronomy*, 2: 181–207, 1997. doi:10.1016/S1384-1076(97)00010-9.
- T. Abel, G. L. Bryan, and M. L. Norman. The Formation of the First Star in the Universe. *Science*, 295:93–98, 2002. doi:10.1126/science.1063991.
- J. Barnes and P. Hut. A hierarchical $O(n \log n)$ force-calculation algorithm. *Nature*, 324:446–449, Dec. 1986. ISSN 0028-0836. doi:10.1038/324446a0.
- P. S. Behroozi, R. H. Wechsler, and H.-Y. Wu. The ROCKSTAR Phase-Space Temporal Halo Finder and the Velocity Offsets of Cluster Cores. *ApJ*, 762:109, 2013. doi:10.1088/0004-637X/762/2/109.
- I. A. Bonnell, M. R. Bate, C. J. Clarke, and J. E. Pringle. Competitive accretion in embedded stellar clusters. *MNRAS*, 323:785–794, 2001. doi:10.1046/j.1365-8711.2001.04270.x.
- W. B. Bonnor. Boyle’s law and gravitational instability. *Monthly Notices of the Royal Astronomical Society*, 116:351, Jan. 1956. ISSN 0035-8711. doi:10.1093/mnras/116.3.351.
- V. Bromm, P. S. Coppi, and R. B. Larson. The Formation of the First Stars. I. The Primordial Star-forming Cloud. *ApJ*, 564: 23–51, 2002. doi:10.1086/323947.
- G. Chabrier. Galactic Stellar and Substellar Initial Mass Function. *PASP*, 115:763–795, 2003. doi:10.1086/376392.
- T. K. Chan, D. Kereš, P. F. Hopkins, E. Quataert, K.-Y. Su, C. C. Hayward, and C.-A. Faucher-Giguère. Cosmic ray feedback in the fire simulations: constraining cosmic ray propagation with gev γ -ray emission. *Monthly Notices of the Royal Astronomical Society*, 488(3):3716–3744, Sept. 2019. ISSN 0035-8711. doi:10.1093/mnras/stz1895.
- K.-J. Chen, M.-Y. Ho, and P.-C. Tung. Formation of supersonic turbulence in the primordial star-forming cloud. *ApJ*, 988:L67, 2025. doi:10.3847/2041-8213/adf18d.
- P. C. Clark, S. C. O. Glover, R. J. Smith, T. H. Greif, R. S. Klessen, and V. Bromm. The Formation and Fragmentation of Disks Around Primordial Protostars. *Science*, 331:1040–1042, 2011. doi:10.1126/science.1198027.

- R. Ebert. Über die verdichtung von h i-gebieten. mit 5 textabbildungen. *Zeitschrift für Astrophysik*, 37:217, Jan. 1955. ISSN 0372-8331.
- C. Federrath, S. Sur, D. R. G. Schleicher, R. Banerjee, and R. S. Klessen. A new jeans resolution criterion for (m)hd simulations of self-gravitating gas: Application to magnetic field amplification by gravity-driven turbulence. *The Astrophysical Journal*, 731(1):62, Mar. 2011. ISSN 0004-637X. doi:10.1088/0004-637X/731/1/62.
- D. Galli and F. Palla. The chemistry of the early Universe. *A&A*, 335:403–420, 1998.
- S. C. O. Glover and T. Abel. Uncertainties in H₂ and HD chemistry and cooling and their role in early structure formation. *MNRAS*, 388:1627–1651, 2008. doi:10.1111/j.1365-2966.2008.13224.x.
- S. C. O. Glover and A.-K. Jappsen. Star formation at very low metallicity. i. chemistry and cooling at low densities. *The Astrophysical Journal*, 666(1):1–19, 2007. ISSN 1538-4357. doi:10.1086/519445.
- N. Y. Gnedin and B. T. Draine. Line overlap and self-shielding of molecular hydrogen in galaxies. *The Astrophysical Journal*, 795(1):37, Oct. 2014. ISSN 0004-637X. doi:10.1088/0004-637X/795/1/37.
- T. H. Greif, V. Springel, S. D. M. White, S. C. O. Glover, P. C. Clark, R. J. Smith, R. S. Klessen, and V. Bromm. Simulations on a Moving Mesh: The Clustered Formation of Population III Protostars. *ApJ*, 737:75, 2011. doi:10.1088/0004-637X/737/2/75.
- M. Y. Grudić, D. Guszejnov, P. F. Hopkins, S. S. R. Offner, and C.-A. Faucher-Giguère. Starforge: Towards a comprehensive numerical model of star cluster formation and feedback. *Monthly Notices of the Royal Astronomical Society*, 506: 2199–2231, 0 2021. ISSN 1365-2966. doi:10.1093/mnras/stab1347. URL <http://dx.doi.org/10.1093/mnras/stab1347>.
- S. Higashi, H. Susa, C. Federrath, and G. Chiaki. Amplification and Saturation of Turbulent Magnetic Fields in Collapsing Primordial Gas Clouds. *ApJ*, 962:158, 2024. doi:10.3847/1538-4357/ad2066.
- S. Hirano and M. N. Machida. Exponentially Amplified Magnetic Field Eliminates Disk Fragmentation around Population III Protostars. *ApJ*, 935:L16, 2022. doi:10.3847/2041-8213/ac85e0.
- S. Hirano, T. Hosokawa, N. Yoshida, H. Umeda, K. Omukai, G. Chiaki, and H. W. Yorke. One Hundred First Stars. *ApJ*, 781:60, 2014. doi:10.1088/0004-637X/781/2/60.
- S. Hirano, N. Sakai, and H. Umeda. Formation of first star clusters under the supersonic gas flow. *MNRAS*, 525:5737, 2023. doi:10.1093/mnras/stad2693.
- P. F. Hopkins. A new class of accurate, mesh-free hydrodynamic simulation methods. *Monthly Notices of the Royal Astronomical Society*, 450:53–110, 0 2015. ISSN 1365-2966. doi:10.1093/mnras/stv195. URL <http://dx.doi.org/10.1093/mnras/stv195>.
- P. F. Hopkins. A constrained-gradient method to control divergence errors in numerical MHD. *MNRAS*, 462:576–587, 2016. doi:10.1093/mnras/stw1578.
- P. F. Hopkins. Anisotropic diffusion in mesh-free numerical magnetohydrodynamics. *Monthly Notices of the Royal Astronomical Society*, 466(3):3387–3405, Apr. 2017. ISSN 0035-8711. doi:10.1093/mnras/stw3306.
- P. F. Hopkins and M. Y. Grudić. Numerical problems in coupling photon momentum (radiation pressure) to gas. *Monthly Notices of the Royal Astronomical Society*, 483(3):4187–4196, Mar. 2019. ISSN 0035-8711. doi:10.1093/mnras/sty3089.
- P. F. Hopkins and M. J. Raives. Accurate, meshless methods for magnetohydrodynamics. *Monthly Notices of the Royal Astronomical Society*, 455(1):51–88, 2015. ISSN 1365-2966. doi:10.1093/mnras/stv2180.
- P. F. Hopkins, A. Wetzel, D. Kereš, C.-A. Faucher-Giguère, E. Quataert, M. Boylan-Kolchin, N. Murray, C. C. Hayward, S. Garrison-Kimmel, C. Hummels, R. Feldmann, P. Torrey, X. Ma, D. Anglés-Alcázar, K.-Y. Su, M. Orr, D. Schmitz, I. Escala, R. Sanderson, M. Y. Grudić, Z. Hafen, J.-H. Kim, A. Fitts, J. S. Bullock, C. Wheeler, T. K. Chan, O. D. Elbert, and D. Narayanan. Fire-2 simulations: physics versus numerics in galaxy formation. *Monthly Notices of the Royal Astronomical Society*, 480(1):800–863, 2018. ISSN 1365-2966. doi:10.1093/mnras/sty1690.
- P. F. Hopkins, M. Y. Grudić, A. Wetzel, D. Kereš, C.-A. Faucher-Giguère, X. Ma, N. Murray, and N. Butcher. Radiative stellar feedback in galaxy formation: Methods and physics. *Monthly Notices of the Royal Astronomical Society*, 491(3):3702–3729, Jan. 2020. ISSN 0035-8711. doi:10.1093/mnras/stz3129.
- P. F. Hopkins, A. Wetzel, C. Wheeler, R. Sanderson, M. Y. Grudić, O. Sameie, M. Boylan-Kolchin, M. Orr, X. Ma, C.-A. Faucher-Giguère, D. Kereš, E. Quataert, K.-Y. Su, J. Moreno, R. Feldmann, J. S. Bullock, S. R. Loebman, D. Anglés-Alcázar, J. Stern, L. Necib, C. R. Choban, and C. C. Hayward. Fire-3: updated stellar evolution models, yields, and microphysics and fitting functions for applications in galaxy simulations. *Monthly Notices of the Royal Astronomical Society*, 519(2): 3154–3181, 2022. ISSN 1365-2966. doi:10.1093/mnras/stac3489.
- P. F. Hopkins, M. Y. Grudić, K.-Y. Su, S. Wellons, D. Angles-Alcazar, U. P. Steinwandel, D. Guszejnov, N. Murray, C.-A. Faucher-Giguère, E. Quataert, and D. Keres. Forge'd in fire: Resolving the end of star formation and structure of agn accretion disks from cosmological initial conditions. 0 2023.
- T. Hosokawa, K. Omukai, N. Yoshida, and H. W. Yorke. Protostellar Feedback Halts the Growth of the First Stars in the Universe. *Science*, 334:1250–1253, 2011. doi:10.1126/science.1207433.
- T. Hosokawa, S. Hirano, R. Kuiper, H. W. Yorke, K. Omukai, and N. Yoshida. Formation of Massive Primordial Stars: Intermittent UV Feedback with Episodic Mass Accretion. *ApJ*, 824:119, 2016. doi:10.3847/0004-637X/824/2/119.
- S. Khullar, M. Grudić, C. Matzner, and N. Murray. Star formation mitosis: The effects of particle splitting on the star formation outcome. In preparation, 2026.
- J.-h. Kim, T. Abel, O. Agertz, G. L. Bryan, D. Ceverino, C. Christensen, C. Conroy, A. Dekel, N. Y. Gnedin, N. J. Goldbaum, J. Guedes, O. Hahn, A. Hobbs, P. F. Hopkins, C. B. Hummels, F. Iannuzzi, D. Keres, A. Klypin, A. V. Kravtsov, M. R. Krumholz, M. Kuhlen, S. N. Leitner, P. Madau, L. Mayer, C. E. Moody, K. Nagamine, M. L. Norman, J. Onorbe, B. W. O'Shea, A. Pillepich, J. R. Primack, T. Quinn, J. I. Read, B. E. Robertson, M. Rocha, D. H. Rudd, S. Shen, B. D. Smith, A. S. Szalay, R. Teyssier, R. Thompson, K. Todoroki, M. J. Turk, J. W. Wadsley, J. H. Wise, Zolotov, Adi, and f. t. A. Collaboration. The agora high-resolution galaxy simulations comparison project. *The Astrophysical Journal Supplement Series*, 210(1):14, Dec. 2013. ISSN 0067-0049. doi:10.1088/0067-0049/210/1/14.
- J.-h. Kim, O. Agertz, R. Teyssier, M. J. Butler, D. Ceverino, J.-H. Choi, R. Feldmann, B. W. Keller, A. Lupi, T. Quinn, Y. Revaz, S. Wallace, N. Y. Gnedin, S. N. Leitner, S. Shen, B. D. Smith, R. Thompson, M. J. Turk, T. Abel, K. S. Arraki, S. M. Benincasa, S. Chakrabarti, C. DeGraf, A. Dekel, N. J. Goldbaum, P. F. Hopkins, C. B. Hummels, A. Klypin, H. Li, P. Madau, N. Mandelker, L. Mayer, K. Nagamine, S. Nickerson, B. W. O'Shea, J. R. Primack, S. Roca-Fàbrega, V. Semenov, I. Shimizu, C. M. Simpson, K. Todoroki, J. W. Wadsley, J. H. Wise, and Collaboration. The agora high-resolution galaxy simulations comparison project. ii. isolated disk test. *The Astrophysical Journal*, 833(2):202, Dec. 2016. ISSN 0004-637X. doi:10.3847/1538-4357/833/2/202.
- R. S. Klessen and S. C. O. Glover. The First Stars: Formation, Properties, and Impact. *ARA&A*, 61:65–130, 2023. doi:10.1146/annurev-astro-071221-053453.
- E. Komatsu et al. Seven-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation. *ApJS*, 192:18, 2011. doi:10.1088/0067-0049/192/2/18.

- P. Kroupa. On the variation of the initial mass function. *MNRAS*, 322:231–246, 2001. doi:10.1046/j.1365-8711.2001.04022.x.
- R. B. Larson. Numerical calculations of the dynamics of collapsing proto-star. *MNRAS*, 145:271–295, 1969. doi:10.1093/mnras/145.3.271.
- M. A. Latif and S. Khochfar. The birth mass function of Population III stars. *ApJ*, 925:28, 2022. doi:10.3847/1538-4357/ac3916.
- C. D. Levermore. Relating eddington factors to flux limiters. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 31(2):149–160, Feb. 1984. ISSN 0022-4073. doi:10.1016/0022-4073(84)90112-2.
- P. S. Li, R. I. Klein, and C. F. McKee. Formation of stellar clusters in magnetized, filamentary infrared dark clouds. *Monthly Notices of the Royal Astronomical Society*, 473: 4220–4241, Jan. 2018. ISSN 0035-8711. doi:10.1093/mnras/stx2611.
- M. N. Machida, K. Omukai, T. Matsumoto, and S.-i. Inutsuka. Conditions for the Formation of First-Star Binaries. *ApJ*, 677: 813–832, 2008. doi:10.1086/533434.
- M. N. Machida, S. Hirano, and S. Basu. Effect of Magnetic Field on the Accretion Phase of Population III Star Formation. *ApJ*, 988:6, 2025. doi:10.3847/1538-4357/addc56.
- C. F. McKee and J. C. Tan. The Formation of Massive Stars from Turbulent Cores. *ApJ*, 585:850–871, 2003. doi:10.1086/346149.
- Y. J. Meziani, P. F. Hopkins, M. Y. Grudić, S. Khullar, C.-A. Faucher-Giguère, and P. J. Gandhi. FORGE'd in the Early Universe: The Effect of Protostellar Outflows on Pop III Accretion. *arXiv e-prints*, 2026.
- D. W. Murray, P. Chang, N. W. Murray, and J. Pittman. Collapse in Self-gravitating Turbulent Fluids. *MNRAS*, 465: 1316–1335, 2017. doi:10.1093/mnras/stw2796.
- S. S. R. Offner, R. I. Klein, C. F. McKee, and M. R. Krumholz. The effects of radiative transfer on low-mass star formation. *The Astrophysical Journal*, 703:131–149, Sept. 2009. ISSN 0004-637X. doi:10.1088/0004-637X/703/1/131.
- K. Omukai. Protostellar collapse with various metallicities. *The Astrophysical Journal*, 534(2):809, May 2000. ISSN 0004-637X. doi:10.1086/308776.
- F. Palla, E. E. Salpeter, and S. W. Stahler. Primordial star formation: the role of molecular hydrogen. *ApJ*, 271:632–641, 1983. doi:10.1086/161231.
- M. V. Penston. Dynamics of self-gravitating gaseous spheres – III. Analytical results in the free-fall of isothermal cases. *MNRAS*, 144:425–448, 1969. doi:10.1093/mnras/144.4.425.
- J. Prieto, P. Padoan, R. Jiménez, and L. Infante. Pop. III stars from turbulent fragmentation at redshift ~ 11 . *ApJ*, 731:L38, 2011. doi:10.1088/2041-8205/731/2/L38.
- J. E. Pringle and A. King. *Astrophysical Flows*. Cambridge University Press, Cambridge, 2007. ISBN 978-0-521-86936-2. doi:10.1017/CBO9780511802201. URL <https://www.cambridge.org/core/books/astrophysical-flows/0D8C02C7C5264B1FCAD6FA47DA70902C>.
- L. R. Prole, P. C. Clark, R. S. Klessen, S. C. O. Glover, and R. Pakmor. Primordial magnetic fields in Population III star formation: a magnetized resolution study. *MNRAS*, 516: 2223–2234, 2022. doi:10.1093/mnras/stac2327.
- E. Ripamonti and T. Abel. Fragmentation and the formation of primordial protostars: the possible role of collision-induced emission. *Monthly Notices of the Royal Astronomical Society*, 348(3):1019–1034, Mar. 2004. ISSN 0035-8711. doi:10.1111/j.1365-2966.2004.07422.x. URL <https://doi.org/10.1111/j.1365-2966.2004.07422.x>.
- B. Robertson and P. Goldreich. Adiabatic Heating of Contracting Turbulent Fluids. *ApJ*, 750(2):L31, 2012. doi:10.1088/2041-8205/750/2/L31.
- K. E. Sadanari, K. Omukai, K. Sugimura, T. Matsumoto, and K. Tomida. Magnetohydrodynamic effect on first star formation: pre-stellar core collapse and protostar formation. *MNRAS*, 505:4197–4214, 2021. doi:10.1093/mnras/stab1330.
- K. E. Sadanari, K. Omukai, K. Sugimura, T. Matsumoto, and K. Tomida. Non-ideal magnetohydrodynamic simulations of the first star formation: the effect of ambipolar diffusion. *MNRAS*, 519:3076–3090, 2023. doi:10.1093/mnras/stac3724.
- K. E. Sadanari, K. Omukai, K. Sugimura, T. Matsumoto, and K. Tomida. Impact of Turbulent Magnetic Fields on Disk Formation and Fragmentation in First Star Formation. *PASJ*, 76:823–840, 2024. doi:10.1093/pasj/psae051.
- P. Sharda and S. H. Menon. Population III star formation in the presence of turbulence, magnetic fields, and ionizing radiation feedback. *MNRAS*, 540:1745–1764, 2025. doi:10.1093/mnras/staf803.
- P. Sharda, C. Federrath, M. R. Krumholz, and D. R. G. Schleicher. Magnetic field amplification in accretion discs around the first stars: implications for the primordial IMF. *MNRAS*, 503:2014–2032, 2021. doi:10.1093/mnras/stab531.
- P. Sharda, S. H. Menon, R. Gerasimov, V. Bromm, B. Burkhardt, L. Haemmerlé, L. van Veenen, and B. D. Wibking. Magnetic fields limit the mass of Population III stars even before the onset of protostellar radiation feedback. *MNRAS*, 541:L1–L7, 2025. doi:10.1093/mnras/lsaf043.
- F. H. Shu. Self-similar collapse of isothermal spheres and star formation. *The Astrophysical Journal*, 214:488–497, June 1977. ISSN 0004-637X. doi:10.1086/155274.
- V. Springel. E pur si muove: Galilean-invariant cosmological hydrodynamical simulations on a moving mesh. *Monthly Notices of the Royal Astronomical Society*, 401:791–851, Jan. 2010. ISSN 0035-8711. doi:10.1111/j.1365-2966.2009.15715.x.
- A. Stacy and V. Bromm. Constraining the statistics of population iii binaries. *Monthly Notices of the Royal Astronomical Society*, 433:1094–1107, Aug. 2013. ISSN 0035-8711. doi:10.1093/mnras/stt789.
- A. Stacy, T. H. Greif, and V. Bromm. The First Stars: Formation of Binaries and Small Multiple Systems. *MNRAS*, 403:45–60, 2010. doi:10.1111/j.1365-2966.2009.16113.x.
- A. Stacy, T. H. Greif, and V. Bromm. The First Stars: Mass Growth Under Protostellar Feedback. *MNRAS*, 422:290–309, 2012. doi:10.1111/j.1365-2966.2012.20605.x.
- A. Stacy, V. Bromm, and A. T. Lee. Building Up the Population III Initial Mass Function from Cosmological Initial Conditions. *MNRAS*, 462:1307–1328, 2016. doi:10.1093/mnras/stw1728.
- A. Stacy, C. F. McKee, A. T. Lee, R. I. Klein, and P. S. Li. Magnetic fields in the formation of the first stars – II. Results. *MNRAS*, 511:5042–5069, 2022. doi:10.1093/mnras/stac372.
- K. Sugimura, T. Matsumoto, T. Hosokawa, S. Hirano, and K. Omukai. The Birth of a Massive First-star Binary. *ApJ*, 892:L14, 2020. doi:10.3847/2041-8213/ab7d37.
- K. Sugimura, T. Matsumoto, T. Hosokawa, S. Hirano, and K. Omukai. Formation of Massive and Wide First-star Binaries in Radiation Hydrodynamic Simulations. *ApJ*, 959:17, 2023. doi:10.3847/1538-4357/ad02fc.
- H. Susa, K. Hasegawa, and N. Tominaga. The Mass Spectrum of the First Stars. *ApJ*, 792:32, 2014. doi:10.1088/0004-637X/792/1/32.
- A. Toomre. On the gravitational stability of a disk of stars. *The Astrophysical Journal*, 139:1217–1238, May 1964. ISSN 0004-637X. doi:10.1086/147861.
- J. K. Truelove, R. I. Klein, C. F. McKee, J. H. Holliman, L. H. Howell, and J. A. Greenough. The Jeans Condition: A New Constraint on Spatial Resolution in Simulations of Isothermal Self-gravitational Hydrodynamics. *ApJ*, 489:L179–L183, 1997. doi:10.1086/310975.
- A. Wetzel, C. C. Hayward, R. E. Sanderson, S. Garrison-Kimmel, C.-A. Faucher-Giguère, D. Kereš, and P. F. Hopkins. Public Data Release of the FIRE-2 Cosmological Zoom-in Simulations of Galaxy Formation. *ApJS*, 265:44, 2023. doi:10.3847/1538-4365/acb99a.
- K. M. J. Wollenberg, S. C. O. Glover, P. C. Clark, and R. S. Klessen. Formation sites of Population III star formation: The effects of different levels of rotation and turbulence on the fragmentation behaviour of primordial gas. *MNRAS*, 494: 1871–1893, 2020. doi:10.1093/mnras/staa289.
- N. Yoshida, K. Omukai, and L. Hernquist. Protostar Formation in the Early Universe. *Science*, 321:669–671, 2008. doi:10.1126/science.1160259.

provides fast and easy peer review for new papers in the **astro-ph** section of the arXiv, making the reviewing pro-

cess simpler for authors and referees alike. Learn more at <http://astro.theoj.org>.