

# DFT - convolution

Dr. Sampath Kumar

Associate Professor

Department of ECE

MIT, Manipal

1) Find the 4 Point DFT of  $x(n)$

$$x(n) = \{2, 1, 2, -1\}$$

$$X(k) = \sum_{n=0}^{4-1} x(n) e^{-j2\pi kn/4}, \quad 0 \leq k \leq 4-1$$

$$= 2 + e^{-j\pi k/2} + 2e^{-j\pi k} - e^{-j3\pi k/2}$$

Alternate method (using Twiddle factor)

$$\begin{bmatrix} X(0) \\ X(1) \\ X(2) \\ X(3) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & W_4^1 & W_4^2 & W_4^3 \\ 1 & W_4^2 & W_4^4 & W_4^6 \\ 1 & W_4^3 & W_4^6 & W_4^9 \end{bmatrix} \begin{bmatrix} x(0) \\ x(1) \\ x(2) \\ x(3) \end{bmatrix}$$

• Ans: 4, -2j, 4, 2j

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -j & -1 & j \\ 1 & -1 & 1 & -1 \\ 1 & j & -1 & -j \end{bmatrix}$$

Derive the 4 Point IDFT of  $X(k) = [ \quad ]$

$$x(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k) e^{j \frac{2\pi nk}{N}}, \quad n=0, 1, \dots, N-1$$

Alternate method (Twiddle factor)

$$x(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k) W_N^{*nk}, \quad n=0, 1, \dots, N-1$$

(29) If we have  $\alpha(n) = [8, 1, 2, 2]$

In this case find 4 point DFT  $X(k)$

then  $x(-n) = x(-n + N)$   $\hookrightarrow$   $N$  point DFT

## ↳ N Point DFT

So in this example

$$x(-1) = 8$$

$$50 \quad x(-1+4) = x(3) = 8$$

Now  $x(n) = [1, 2, 2, 8]$

Then proceed to find 4-Point DFT  $X(k)$ .

# Applications of DFT – IDFT tools

- Frequency (spectrum) analysis
- Power spectrum estimations
- Linear filtering
- Presence of computationally efficient algorithms

# Multiplication of two DFT and circular convolution

- Let

$$X_1(k) = \sum_{n=0}^{N-1} x_1(n) e^{-j2\pi nk/N}, \quad k = 0, 1, \dots, N-1$$

$$X_2(k) = \sum_{n=0}^{N-1} x_2(n) e^{-j2\pi nk/N}, \quad k = 0, 1, \dots, N-1$$

We have

$$X_3(k) = X_1(k) X_2(k), \quad k = 0, 1, \dots, N-1$$

Let us determine the relationship between  $x_3(n)$  and the sequences  $x_1(n)$  and  $x_2(n)$ .

The IDFT of  $\{X_3(k)\}$  is

$$\begin{aligned}x_3(m) &= \frac{1}{N} \sum_{k=0}^{N-1} X_3(k) e^{j2\pi km/N} \\&= \frac{1}{N} \sum_{k=0}^{N-1} X_1(k) X_2(k) e^{j2\pi km/N}\end{aligned}$$

$$x_3(m) = \frac{1}{N} \sum_{k=0}^{N-1} \left[ \sum_{n=0}^{N-1} x_1(n) e^{-j2\pi kn/N} \right] \left[ \sum_{l=0}^{N-1} x_2(l) e^{-j2\pi kl/N} \right] e^{j2\pi km/N}$$

$$= \frac{1}{N} \sum_{n=0}^{N-1} x_1(n) \sum_{l=0}^{N-1} x_2(l) \left[ \sum_{k=0}^{N-1} e^{j2\pi k(m-n-l)/N} \right]$$

$$x_3(m) = \frac{1}{N} \sum_{n=0}^{N-1} x_1(n) \sum_{l=0}^{N-1} x_2(l) \left[ \sum_{k=0}^{N-1} e^{j2\pi k(m-n-l)/N} \right]$$

$$\sum_{k=0}^{N-1} a^k = \begin{cases} N, & a = 1 \\ \frac{1 - a^N}{1 - a}, & a \neq 1 \end{cases}$$

where  $a$  is defined as

$$a = e^{j2\pi(m-n-l)/N}$$

$$\sum_{k=0}^{N-1} a^k = \begin{cases} N, & l = ((m - n))_N \\ 0, & \text{otherwise} \end{cases}$$

$$x_3(m) = \sum_{n=0}^{N-1} x_1(n) x_2((m - n))_N, \quad m = 0, 1, \dots, N - 1 \quad \text{circular convolution.}$$

$$x_1(n) \circledast x_2(n) \xrightleftharpoons[N]{\text{DFT}} X_1(k) X_2(k)$$



*Thank  
you*

