

# Properties of DFT contd...

Dr. Sampath Kumar

Associate Professor

Department of ECE

MIT, Manipal

# Circular time shift of a sequence

$$x((n-l))_N \xleftrightarrow[N]{\text{DFT}} X(k)e^{-j2\pi kl/N}$$

$$\text{DFT}\{x((n-l))_N\} = \sum_{n=0}^{N-1} x((n-l))_N e^{-j2\pi kn/N}$$

$$= \sum_{n=0}^{l-1} x((n-l))_N e^{-j2\pi kn/N}$$

$$+ \sum_{n=l}^{N-1} x(n-l) e^{-j2\pi kn/N}$$

$$\sum_{n=l}^{N-1} x(n-l) e^{-j2\pi kn/N} = \sum_{m=0}^{N-1-l} x(m) e^{-j2\pi k(m+l)/N}$$

Put  $n-l = m$

But  $x((n-l))_N = x(N-l+n)$ .

$$\sum_{n=0}^{l-1} x((n-l))_N e^{-j2\pi kn/N} = \sum_{n=0}^{l-1} x(N-l+n) e^{-j2\pi kn/N}$$

$$= \sum_{m=N-l}^{N-1} x(m) e^{-j2\pi k(m+l)/N}$$

Put  $N-l+n = m$

$$\begin{aligned} \text{Therefore, DFT}\{x((n-l))\} &= \sum_{m=0}^{N-1} x(m) e^{-j2\pi k(m+l)/N} \\ &= X(k) e^{-j2\pi kl/N} \end{aligned}$$

## Circular frequency shift.

$$x(n)e^{j2\pi ln/N} \xleftrightarrow[N]{\text{DFT}} X((k-l))_N$$

This is the dual to the circular time-shifting property and its proof is similar

## Complex-conjugate properties

$$x^*(n) \xleftrightarrow[N]{\text{DFT}} X^*((-k))_N = X^*(N - k)$$

Proof is an exercise

The IDFT of  $X^*(k)$  is

$$\frac{1}{N} \sum_{k=0}^{N-1} X^*(k) e^{j2\pi kn/N} = \left[ \frac{1}{N} \sum_{k=0}^{N-1} X(k) e^{j2\pi k(N-n)/N} \right]$$

Therefore,

$$x^*((-n))_N = x^*(N - n) \xleftrightarrow[N]{\text{DFT}} X^*(k)$$

## Circular correlation.

$$\tilde{r}_{xy}(l) \xleftrightarrow[N]{\text{DFT}} \tilde{R}_{xy}(k) = X(k)Y^*(k)$$

where  $\tilde{r}_{xy}(l)$  is the (unnormalized) circular crosscorrelation sequence, defined as

$$\tilde{r}_{xy}(l) = \sum_{n=0}^{N-1} x(n)y^*((n-l))N$$

circular autocorrelation of  $x(n)$ ,

$$\tilde{r}_{xx}(l) \xleftrightarrow[N]{\text{DFT}} \tilde{R}_{xx}(k) = |X(k)|^2$$

## Multiplication of two sequences.

$$x_1(n)x_2(n) \xleftrightarrow[N]{\text{DFT}} \frac{1}{N} X_1(k) \otimes X_2(k)$$

Its proof follows simply by interchanging the roles of time and frequency in the expression for the circular convolution of two sequences.

## Parseval's Theorem.

$$\sum_{n=0}^{N-1} x(n)y^*(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k)Y^*(k)$$

*Proof* The property follows immediately from the circular correlation property

$$\sum_{n=0}^{N-1} x(n)y^*(n) = \tilde{r}_{xy}(0)$$

and

$$\begin{aligned}\tilde{r}_{xy}(l) &= \frac{1}{N} \sum_{k=0}^{N-1} \tilde{R}_{xy}(k) e^{j2\pi kl/N} \\ &= \frac{1}{N} \sum_{k=0}^{N-1} X(k)Y^*(k) e^{j2\pi kl/N}\end{aligned}$$

Evaluate IDFT at  $l = 0$

special case where  $y(n) = x(n)$

$$\sum_{n=0}^{N-1} |x(n)|^2 = \frac{1}{N} \sum_{k=0}^{N-1} |X(k)|^2$$



**TABLE 7.2** Properties of the DFT

Property	Time Domain	Frequency Domain
Notation	$x(n), y(n)$	$X(k), Y(k)$
Periodicity	$x(n) = x(n + N)$	$X(k) = X(k + N)$
Linearity	$a_1 x_1(n) + a_2 x_2(n)$	$a_1 X_1(k) + a_2 X_2(k)$
Time reversal	$x(N - n)$	$X(N - k)$
Circular time shift	$x((n - l))_N$	$X(k)e^{-j2\pi kl/N}$
Circular frequency shift	$x(n)e^{j2\pi ln/N}$	$X((k - l))_N$
Complex conjugate	$x^*(n)$	$X^*(N - k)$
Circular convolution	$x_1(n) \circledast x_2(n)$	$X_1(k)X_2(k)$
Circular correlation	$x(n) \circledast y^*(-n)$	$X(k)Y^*(k)$
Multiplication of two sequences	$x_1(n)x_2(n)$	$\frac{1}{N} X_1(k) \circledast X_2(k)$
Parseval's theorem	$\sum_{n=0}^{N-1} x(n)y^*(n)$	$\frac{1}{N} \sum_{k=0}^{N-1} X(k)Y^*(k)$



## The DFT as a Linear Transformation

$$X(k) = \sum_{n=0}^{N-1} x(n) W_N^{kn}, \quad k = 0, 1, \dots, N-1$$

$$x(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k) W_N^{-kn}, \quad n = 0, 1, \dots, N-1$$

$$W_N = e^{-j2\pi/N} \quad \text{which is an } N\text{th root of unity.}$$

We note that the computation of each point of the DFT can be accomplished by  $N$  complex multiplications and  $(N-1)$  complex additions. Hence the  $N$ -point DFT values can be computed in a total of  $N^2$  complex multiplications and  $N(N-1)$  complex additions.

# DFT matrix

$$\mathbf{x}_N = \begin{bmatrix} x(0) \\ x(1) \\ \vdots \\ x(N-1) \end{bmatrix}, \quad \mathbf{X}_N = \begin{bmatrix} X(0) \\ X(1) \\ \vdots \\ X(N-1) \end{bmatrix}$$

$$X(k) = \sum_{n=0}^{N-1} x(n) W_N^{kn}, \quad k = 0, 1, \dots, N-1$$

$$x(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k) W_N^{-kn}, \quad n = 0, 1, \dots, N-1$$

$$W_N = e^{-j2\pi/N}$$

$$\mathbf{W}_N = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & W_N & W_N^2 & \dots & W_N^{N-1} \\ & W_N^2 & W_N^4 & \dots & W_N^{2(N-1)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & W_N^{N-1} & W_N^{2(N-1)} & \dots & W_N^{(N-1)(N-1)} \end{bmatrix}$$

$$\mathbf{X}_N = \mathbf{W}_N \mathbf{x}_N$$

$$\mathbf{x}_N = \mathbf{W}_N^{-1} \mathbf{X}_N$$

$$\mathbf{x}_N = \frac{1}{N} \mathbf{W}_N^* \mathbf{X}_N$$

$$\begin{bmatrix} X(0) \\ X(1) \\ \vdots \\ X(N-1) \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & W_N & W_N^2 & \dots & W_N^{N-1} \\ & W_N^2 & W_N^4 & \dots & W_N^{2(N-1)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & W_N^{N-1} & W_N^{2(N-1)} & \dots & W_N^{(N-1)(N-1)} \end{bmatrix} \begin{bmatrix} x(0) \\ x(1) \\ \vdots \\ x(N-1) \end{bmatrix}$$

where  $\mathbf{W}_N^*$  denotes the complex conjugate of the matrix  $\mathbf{W}_N$

$$\mathbf{W}_N^{-1} = \frac{1}{N} \mathbf{W}_N^*$$

$$\mathbf{W}_N \mathbf{W}_N^* = N \mathbf{I}_N$$

orthogonal (unitary) matrix.

Compute the DFT of the four-point sequence

$$x(n) = (0 \quad 1 \quad 2 \quad 3)$$

$W_N^{kn} \Rightarrow$  Twiddle factor

eg:  $W_4^1 = \left( e^{-j\frac{2\pi}{4}} \right)^1 = \underline{-j}$

$$W_8^1 = \left( e^{-j\frac{2\pi}{8}} \right)^1 = \underline{\underline{\frac{1}{\sqrt{2}} - \frac{j}{\sqrt{2}}}}$$

**Solution.** The first step is to determine the matrix  $\mathbf{W}_4$ . By exploiting the periodicity property of  $\mathbf{W}_4$  and the symmetry property

$$W_N^{k+N/2} = -W_N^k$$

$$\begin{aligned} \mathbf{W}_4 &= \begin{bmatrix} W_4^0 & W_4^0 & W_4^0 & W_4^0 \\ W_4^0 & W_4^1 & W_4^2 & W_4^3 \\ W_4^0 & W_4^2 & W_4^4 & W_4^6 \\ W_4^0 & W_4^3 & W_4^6 & W_4^9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & W_4^1 & W_4^2 & W_4^3 \\ 1 & W_4^2 & W_4^0 & W_4^2 \\ 1 & W_4^3 & W_4^2 & W_4^1 \end{bmatrix} & \mathbf{X}_4 = \mathbf{W}_4 \mathbf{x}_4 = \begin{bmatrix} 6 \\ -2 + 2j \\ -2 \\ -2 - 2j \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -j & -1 & j \\ 1 & -1 & 1 & -1 \\ 1 & j & -1 & -j \end{bmatrix} \end{aligned}$$

The IDFT of  $\mathbf{X}_4$  may be determined by conjugating the elements in  $\mathbf{W}_4$  to obtain  $\mathbf{W}_4^*$

*Thank  
you*

