

Energy Resolution of Runge-Kutta Reconstruction of Padded Microscopically Simulated Tracks

Second Iteration, More Data

Martin Vavřík

martin.vavrik@cvut.cz
IEAP CTU PRAGUE

December 15, 2023



Outline

- 1 Grid-like track simulation
- 2 Direction independent resolution
- 3 Direction dependent resolution



- 1 Grid-like track simulation
- 2 Direction independent resolution
- 3 Direction dependent resolution



Grid-like track simulation

- We use Garfield++ for track simulation
 - Primary relativistic particle simulated using HEED program [1]
 - Secondary ionization electrons simulated using microscopic tracking (uses equation of motion)
- Microscopic tracks were simulated on MetaCentrum to test the Runge-Kutta padded reconstruction
 - 2000 + 9702 jobs, each 20-160 hours runtime, 20 GB RAM allocated
 - 5 sets of 9702 tracks were simulated (electron vs positron, 21 theta angles, 21 phi angles, 11 energies 3-13 MeV)
 - Reconstruction lasts around 0.3 seconds per track



Simulation ranges

Spherical angles (θ, φ) with respect to z
 θ taken from the equatorial plane xy




Diagram of the batch simulation parameters:
 $\theta \in [-17.1^\circ, 17.1^\circ]$, $\varphi \in [-16.3^\circ, 16.3^\circ]$, $E_{\text{kin.}} \in [3, 13] \text{ MeV}$.



Outline


- 1 Grid-like track simulation
- 2 Direction independent resolution
- 3 Direction dependent resolution



Energy resolution for all tracks


Electrons

Energy resolution of Runge-Kutta reconstruction (with pads)



Positrons


Energy resolution of Runge-Kutta reconstruction (with pads)



Energy resolution dependence on simulated energy


Electrons

Energy resolution dependence on energy



Positrons

Energy resolution dependence on energy



Outline


- 1 Grid-like track simulation
- 2 Direction independent resolution
- 3 Direction dependent resolution



Energy resolution dependence on phi


Electrons

Energy resolution dependence on phi




Positrons

Energy resolution dependence on phi




Energy resolution dependence on theta

Electrons



Positrons



Thank you for your attention.



[1] I. B. Smirnov.

Modeling of ionization produced by fast charged particles in gases.
Nucl. Instr. Meth. A, 554:474–493, 2005.

