

RESURGENCE OF THE AIRY FUNCTION AND OTHER EXPONENTIAL INTEGRALS

VERONICA FANTINI AND AARON FENYES

1. HYPERGEOMETRIC FUNCTIONS AS BOREL TRANSFORM OF SECOND ORDER ODE (series normales de 1er ordre)

Let us consider the following linear second order ODE

$$(1.1) \quad \left[P\left(\frac{\partial}{\partial z}\right) + \frac{1}{z}Q\left(\frac{\partial}{\partial z}\right) + \frac{1}{z}R\left(\frac{1}{z}\right) \right] f(z) = 0$$

with $\deg P = 2$, $\deg Q = 1$ and $R = O(\frac{1}{z})$. We denote by α_1, α_2 the roots of $P(\lambda)$ and we assume they are distinct. Furthermore we assume $\tau_j := \frac{Q(-\alpha_j)}{P'(-\alpha_j)} \neq 0$. The latter assumption guarantees the formal solution \tilde{f} being slight, while the former assumption implies there will be two independent solutions.

Under the previous assumptions we prove that the Borel transformed solution $\hat{f}(\zeta_j)$ is a Gauss hypergeometric function, $\zeta_j = \zeta - \alpha_j$.

Proposition 1.1. Let $P(\lambda) = \lambda^2 + a_1\lambda + a_0$, $Q(\lambda) = b_1\lambda + b_0$ and $R(\frac{1}{z}) = \frac{c_1}{z}$ satisfying the previous assumptions. Then

$$(1.2) \quad \hat{f}(\zeta_1) = \left(1 - \frac{\zeta_1}{\alpha_2 - \alpha_1}\right)^{c-1} {}_2F_1\left(a, b; c; 1 - \frac{\zeta_1}{\alpha_2 - \alpha_1}\right)$$

$$(1.3) \quad \hat{f}(\zeta_2) = \left(1 - \frac{\zeta_2}{\alpha_1 - \alpha_2}\right)^{c-1} {}_2F_1\left(a, b; c; 1 - \frac{\zeta_2}{\alpha_1 - \alpha_2}\right)$$

where the coefficients a, b, c depend on the parameter of P, Q, R .

Proof. We start by taking the Borel transform of (1.1):

$$(1.4) \quad (\zeta^2 - a_1\zeta + a_0)\hat{f}(\zeta) + \int_0^\zeta b_1(-\zeta')\hat{f}(\zeta')d\zeta' + b_0 \int_0^\zeta \hat{f}(\zeta')d\zeta' + c_1 \int_0^\zeta (\zeta - \zeta')\hat{f}(\zeta')d\zeta' = 0$$

then we differentiate twice in order to have a differential equation which can be easier recognized as a hypergeometric equation. Since \tilde{F} is slight and locally integrable at 0 by assumption, Proposition 1 Resurgent Airy doc by Aaron tells we are not loosing information taking derivatives, and that $\hat{f}(\zeta)$ is a solution of (1.4) if and only

if it is a solution of (1.5)

$$(1.5) \quad [(\zeta^2 - a_1\zeta + a_0)\partial_\zeta^2 + (4\zeta - b_1\zeta - 2a_1 + b_0)\partial_\zeta + (c_1 + 2 - b_1)]\hat{f}(\zeta) = 0$$

We introduce some notation to simplify the computations, we denote by $\beta_1 = 4 - b_1$, $\beta_0 = b_0 - 2a_1$, $\gamma = c_1 + 2 - b_1$ so (1.5) turns into

$$[(\zeta - \alpha_1)(\zeta - \alpha_2)\partial_\zeta^2 + (\beta_1\zeta + \beta_0)\partial_\zeta + \gamma]\hat{f}(\zeta) = 0$$

We consider the following change of coordinates $\zeta = \alpha_2 - (\alpha_2 - \alpha_1)\xi$ ¹

$$[(\alpha_2 - (\alpha_2 - \alpha_1)\xi - \alpha_1)(\alpha_2 - (\alpha_2 - \alpha_1)\xi - \alpha_2)(\alpha_1 - \alpha_2)^{-2}\partial_\xi^2 + (\beta_1(\alpha_2 - (\alpha_2 - \alpha_1)\xi) + \beta_0)(\alpha_1 - \alpha_2)^{-1}\partial_\xi + \gamma]\hat{f}(\xi) = 0$$

$$[(\alpha_2 - \alpha_1)(1 - \xi)(\alpha_1 - \alpha_2)\xi(\alpha_1 - \alpha_2)^{-2}\partial_\xi^2 + (\beta_1\alpha_2 - \beta_1(\alpha_2 - \alpha_1)\xi + \beta_0)(\alpha_1 - \alpha_2)^{-1}\partial_\xi + \gamma]\hat{f}(\xi) = 0$$

$$[-(1 - \xi)\xi\partial_\xi^2 + ((\beta_1\alpha_2 + \beta_0)(\alpha_1 - \alpha_2)^{-1} + \beta_1\xi)\partial_\xi + \gamma]\hat{f}(\xi) = 0$$

$$[(1 - \xi)\xi\partial_\xi^2 - ((\beta_1\alpha_2 + \beta_0)(\alpha_1 - \alpha_2)^{-1} + \beta_1\xi)\partial_\xi - \gamma]\hat{f}(\xi) = 0$$

The latter equation is an hypergeometric equation of parameters

$$C = (\beta_1\alpha_2 + \beta_0)(\alpha_2 - \alpha_1)^{-1}$$

$$A + B + 1 = \beta_1 = 4 - b_1 \Rightarrow A + B = 3 - b_1$$

$$AB = \gamma = c_1 + 2 - b_1$$

and a solution is given by

$$\begin{aligned} \hat{f}(\xi) &= \xi^{1-C} {}_2F_1(A - C + 1, B - C + 1; 2 - C; \xi) \\ &= \left(\frac{\alpha_2 - \xi}{\alpha_2 - \alpha_1}\right)^{1-C} {}_2F_1\left(A - C + 1, B - C + 1; 2 - C; \frac{\alpha_2 - \xi}{\alpha_2 - \alpha_1}\right) \\ &= \left(1 - \frac{\xi_1}{\alpha_2 - \alpha_1}\right)^{1-C} {}_2F_1\left(A - C + 1, B - C + 1; 2 - C; 1 - \frac{\xi_1}{\alpha_2 - \alpha_1}\right) \end{aligned}$$

□

¹ $\partial_\zeta = (\alpha_1 - \alpha_2)^{-1}\partial_\xi$ and $\partial_\zeta^2 = (\alpha_1 - \alpha_2)^{-2}\partial_\xi^2$