

# Resurgence of a parabolic cylinder function

Aaron Fenyes

May 19, 2022

## 1 Hypergeometric functions in trigonometry

### 1.1 Chebyshev sine formulas

By definition,

$$\begin{aligned}\cos(n\phi) &= T_n(\cos(\phi)) \\ \sin(n\phi) &= U_{n-1}(\cos(\phi)) \sin(\phi).\end{aligned}$$

Furthermore,

$$\begin{aligned}\cos(n\phi - \frac{n}{2}\pi) &= T_n(\cos(\phi - \frac{\pi}{2})) \\ \cos(n\phi) \cos(\frac{n}{2}\pi) + \sin(n\phi) \sin(\frac{n}{2}\pi) &= T_n(\sin(\phi)).\end{aligned}$$

Thus, for  $n = 2k + 1$ , we have

$$(-1)^k \sin(n\phi) = T_n(\sin(\phi)),$$

implying also that

$$\begin{aligned}(-1)^k n \cos(n\phi) &= T'_n(\sin(\phi)) \cos(\phi) \\ (-1)^k n T_n(\cos(\phi)) &= n U_{n-1}(\sin(\phi)) \cos(\phi) \\ (-1)^k T_n(\cos(\phi)) &= U_{n-1}(\sin(\phi)) \cos(\phi).\end{aligned}$$

### 1.2 A hypergeometric identity

Let  $a = \frac{1}{2} - \frac{1}{2n}$ , giving  $1 - 2a = \frac{1}{n}$ .

#### 1.2.1 All orders

From [?, equation 15.4.16],

$$F(a, 1 - a, \frac{3}{2}; \sin(\theta)^2) = \frac{\sin((1 - 2a)\theta)}{(1 - 2a) \sin(\theta)}.$$

so we have

$$\begin{aligned}
F\left(\frac{1}{2} - \frac{m}{2n}, \frac{1}{2} + \frac{m}{2n}, \frac{3}{2}; \sin(\theta)^2\right) &= \frac{\sin(\frac{m}{n}\theta)}{\frac{m}{n}\sin(\theta)} \\
&= \frac{n}{m} \cdot \frac{\sin(m\theta/n)}{\sin(\theta/n)} \cdot \frac{\sin(\theta/n)}{\sin(\theta)} \\
&= \frac{n}{m} \cdot \frac{U_{m-1}(\cos(\theta/n))}{U_{n-1}(\cos(\theta/n))}.
\end{aligned}$$

Letting  $u = \cos(\theta/n)$ , we get the identity

$$F\left(\frac{1}{2} - \frac{m}{2n}, \frac{1}{2} + \frac{m}{2n}, \frac{3}{2}; 1 - T_n(u)^2\right) = \frac{n}{m} \cdot \frac{U_{m-1}(u)}{U_{n-1}(u)}.$$

From our reasoning so far, we can only conclude the identity holds when  $\pm T_n(u)$  is in the right half-plane.

Identity 15.10.17 [or, better, 15.8.4] from [?] splits the left-hand side above into

$$\begin{aligned}
&\frac{\Gamma(\frac{1}{2})\Gamma(\frac{3}{2})}{\Gamma(1 - \frac{m}{2n})\Gamma(1 + \frac{m}{2n})} F\left(\frac{1}{2} - \frac{m}{2n}, \frac{1}{2} + \frac{m}{2n}, \frac{1}{2}; T_n(u)^2\right) \\
&\pm \frac{\Gamma(-\frac{1}{2})\Gamma(\frac{3}{2})}{\Gamma(\frac{1}{2} - \frac{m}{2n})\Gamma(\frac{1}{2} + \frac{m}{2n})} T_n(u) F\left(1 - \frac{m}{2n}, 1 + \frac{m}{2n}, \frac{3}{2}; T_n(u)^2\right),
\end{aligned}$$

where the sign must be chosen so that  $\pm T_n(u)$  is in the right half-plane (?). Applying identities 15.8.27–28 from [?] turns this into

$$\begin{aligned}
&\frac{1}{2}F\left(1 - \frac{m}{n}, 1 + \frac{m}{n}, \frac{3}{2}, \frac{1}{2} \mp \frac{1}{2}T_n(u)\right) + \frac{1}{2}F\left(1 - \frac{m}{n}, 1 + \frac{m}{n}, \frac{3}{2}, \frac{1}{2} \pm \frac{1}{2}T_n(u)\right) \\
&+ \frac{1}{2}F\left(1 - \frac{m}{n}, 1 + \frac{m}{n}, \frac{3}{2}, \frac{1}{2} \mp \frac{1}{2}T_n(u)\right) - \frac{1}{2}F\left(1 - \frac{m}{n}, 1 + \frac{m}{n}, \frac{3}{2}, \frac{1}{2} \pm \frac{1}{2}T_n(u)\right).
\end{aligned}$$

After cancellation, we conclude that

$$F\left(1 - \frac{m}{n}, 1 + \frac{m}{n}, \frac{3}{2}, \frac{1}{2} \mp \frac{1}{2}T_n(u)\right) = \frac{n}{m} \cdot \frac{U_{m-1}(u)}{U_{n-1}(u)}$$

when  $\pm T_n(u)$  is in the right half-plane (?).

### 1.2.2 Odd order

From [?, equation 15.4.14],

$$F(a, 1 - a, \frac{1}{2}; \sin(\theta)^2) = \frac{\cos((1 - 2a)\theta)}{\cos(\theta)}.$$

so we have

$$\begin{aligned}
F\left(\frac{1}{2} - \frac{1}{2n}, \frac{1}{2} + \frac{1}{2n}, \frac{1}{2}; \sin(\theta)^2\right) &= \frac{\cos(\theta/n)}{\cos(\theta)} \\
&= \frac{\cos(\theta/n)}{T_n(\cos(\theta/n))}.
\end{aligned}$$

Let  $u = \sin(\theta/n)$ . If  $n = 2k + 1$ , we get the identity

$$F\left(\frac{1}{2} - \frac{1}{2n}, \frac{1}{2} + \frac{1}{2n}, \frac{1}{2}; T_n(u)^2\right) = \frac{(-1)^k}{U_{n-1}(u)}.$$

## 2 The Weber equation

### 2.1 Basics

The Weber equation is

$$\left[ \left( \frac{\partial}{\partial y} \right)^2 - \left( \frac{1}{4} y^2 + a \right) \right] \psi = 0.$$

Setting  $a$  to zero, we get

$$\left[ \left( \frac{\partial}{\partial y} \right)^2 - \frac{1}{4} y^2 \right] \psi = 0. \quad (1)$$

One solution is given by the parabolic cylinder function [?, equation 12.7.10]

$$U(0, y) = \frac{1}{\sqrt{2\pi}} y^{1/2} K_{1/4}(\tfrac{1}{4} y^2).$$

Combining the identity [?, equation 10.27.4]

$$K_\nu(z) = \frac{\pi}{2 \sin(\nu\pi)} (I_{-\nu}(z) - I_\nu(z))$$

and the integral [?, equation 10.32.12]

$$I_\nu(z) = \frac{1}{2\pi i} \int_{\mathcal{H}} e^{z \cosh(\phi)} e^{-\nu\phi} d\phi,$$

where  $\mathcal{H}$  runs clockwise around the rectangle

$$0 < \operatorname{Re}(\phi) \qquad |\operatorname{Im}(\phi)| < \pi,$$

we learn that

$$\begin{aligned} I_{-\nu}(z) - I_\nu(z) &= \frac{1}{\pi i} \int_{\mathcal{H}} e^{z \cosh(\phi)} \sinh(\nu\phi) d\phi \\ K_\nu(z) &= \frac{1}{2i \sin(\nu\pi)} \int_{\mathcal{H}} e^{z \cosh(\phi)} \sinh(\nu\phi) d\phi. \end{aligned}$$

In particular, we have

$$K_{1/4}(z) = \frac{1}{i\sqrt{2}} \int_{\mathcal{H}} e^{z \cosh(\phi)} \sinh(\phi/4) d\phi.$$

Setting  $u = \cosh(\phi/4)$  and getting  $\cosh(\phi) = 8u^4 - 8u^2 + 1$  from a table of Chebyshev polynomials, we can write

$$K_{1/4}(z) = \frac{2\sqrt{2}}{i} \int_{\Gamma} \exp[z(8u^4 - 8u^2 + 1)] du, \quad (2)$$

where  $\Gamma$  is a path that comes from  $\infty$  at  $-45^\circ$  and goes to  $\infty$  at  $45^\circ$ . **[This feels dubious; check using definition of  $\mathcal{H}$ .]**

## 2.2 Contour argument

We can recast integral 2 into  $\hat{\mathbb{C}}$  by setting  $-\zeta = 8u^4 - 8u^2 + 1$ . Projecting  $\Gamma$  to a contour  $\gamma$  in  $\hat{\mathbb{C}}$  and choosing the branch of  $u$  that lifts  $\gamma$  back to  $\Gamma$ , we have

$$K_{1/4} = \frac{i}{\sqrt{2}} \int_{\gamma} e^{-z\zeta} \frac{d\zeta}{8u^3 - 4u}. \quad (3)$$

The integrand has poles at  $u = \pm \frac{1}{\sqrt{2}}$ , and  $\gamma$  runs counterclockwise around  $[\frac{1}{\sqrt{2}}, \infty)$ .

Using the identity from Section 1.2.1, we can rewrite integral 3 as

$$K_{1/4} = \frac{i}{\sqrt{2}} \int_{\gamma_z} e^{-z\zeta} F\left(\frac{3}{8}, \frac{5}{8}; \frac{3}{2}; 1 - \zeta^2\right) d\zeta.$$