

Resurgence of the Airy function and other exponential integrals

Veronica Fantini and Aaron Fenyes

February 25, 2022

1 Introduction

1.1 Why does Borel resummation work?

Borel resummation is a way of turning a formal power series

$$\varphi_{\bullet} = z^{\sigma} \left(\frac{\varphi_0}{z} + \frac{\varphi_1}{z^2} + \frac{\varphi_2}{z^3} + \frac{\varphi_3}{z^4} + \dots \right),$$

with $\sigma \in [0, 1)$, into a function which is asymptotic to φ_{\bullet} as $z \rightarrow \infty$. Different functions can be asymptotic to the same power series, and Borel resummation picks one of them, performing an implicit regularization [[arXiv:1705.03071](#), or maybe [arXiv:1412.6614](#)]. When a function matches its Borel sum, we'll say it's *Borel regular*. Several familiar kinds of regularity imply Borel regularity, and shed light on why it occurs.

- **Having a good asymptotic approximation**

Let R_N be the difference between a function and the partial sum

$$\frac{\varphi_0}{z} + \frac{\varphi_1}{z^2} + \frac{\varphi_2}{z^3} + \dots + \frac{\varphi_{N-2}}{z^{N-1}}$$

of its asymptotic series. Watson showed a century ago that the function is Borel regular whenever there's a constant $c \in (0, \infty)$ with

$$|R_N| \leq \frac{c^{N+1} N!}{|z|^N}$$

over all orders N and all z in a wide enough wedge around infinity.

- **Satisfying a singular differential equation**

- **Being a thimble integral**

Let X be a translation surface—a Riemann surface carrying a holomorphic 1-form ν . Suppose X is of *meromorphic type*, meaning that we got it by puncturing a compact Riemann surface \bar{X} at finitely many points, and ν has a pole at each puncture. A *translation coordinate* on X is a local coordinate whose derivative is ν .

Take another meromorphic-type translation surface B and a holomorphic map $f: \bar{X} \rightarrow \bar{B}$ that sends punctures to punctures. Suppose the critical points of f are all simple. For each critical point p , let Γ_p be the ray going rightward from $f(p)$, and let ζ_p be the translation coordinate around Γ_p which vanishes at $f(p)$. These are well-defined as long as Γ_p misses the singularities of B . The preimage $f^{-1}(\Gamma_p)$ is a bunch of disjoint curves, as long as Γ_p misses the other critical values of f . [Is B the Borel plane? Can we assume all its singularities are critical values of f ?] The Lefschetz thimble Λ_p is the component of $f^{-1}(\Gamma_p)$ that goes through p , oriented so that shifting it to its left would make its projection run clockwise around Γ_p . The thimble integral

$$I_p = \int_{\Lambda_p} e^{-zf^*\zeta_p} \nu$$

is a holomorphic function on the right half-plane parameterized by z and it turns out [we hope] to be Borel regular.

[Talk about exponential integrals and their decomposition into thimble integrals.]

In higher-dimensional complex manifolds, integrals over Lefschetz thimbles are still Borel regular [“Exponential integrals, Lefschetz thimbles and linear resurgence”][“Exponential Integral” lectures?]. This fact plays an important technical role in quantum mechanics, where infinite-dimensional exponential integrals are supposed to give the expectation values of observable quantities. Physicists often use Borel summation and related techniques to assign values to these integrals [Costin & Kruskal, “On optimal truncation...”].

Choose a path $\gamma: \mathbb{R} \rightarrow X$ whose projection $f \circ \gamma$ starts out going leftward out of a puncture, ends up going rightward into a puncture, and never touches a critical value of f . [How do we get jumps at the critical values?] Choose a translation coordinate ζ on B and continue it along $f \circ \gamma$, noting that it may become multi-valued if $f \circ \gamma$ intersects itself. This data defines the exponential integral

$$I = \int_{\gamma} e^{-zf^*\zeta} \nu,$$

a holomorphic function on the right half-plane parameterized by z . It turns out [we hope] that we can get I by summing $e^{-\alpha_p z} I_p$ over various critical points. The constants α_p are values of ζ , continued to the critical points along certain paths.

- Each resummation method for asymptotic series makes some implicit assumption that allows us to reconstruct a holomorphic function from its asymptotic behavior.
- The resummation method works correctly for functions which satisfy that assumption.
- For the modified Bessel function $K_{1/3}$, Borel resummation works because the asymptotic series encodes a second-order differential equation.
 - Different aspects of this example appear in various places (Mariño, Kawai–Takei, Sauzin). We give a detailed, unified treatment.

- We can generalize this argument to all $K_{1/n}$ and their limit K_0 .
- We can also generalize to all third-order exponential integrals.
 - Most of them are equivalent to the $K_{1/3}$ integral, but there's also an interesting degeneration.

1.2 Fractional derivative formula

- Theorem ?? says that for a certain class of exponential integrals

$$I(z) = \int_{\Gamma} e^{-zf} \nu,$$

the inverse Laplace [better to say Borel?] transform is the $\frac{3}{2}$ derivative of $d\zeta/df$, where $f^*d\zeta = \nu$ [check].

- the asymptotic expansion of $I(z)$ is a resurgent function.
- Is it always a *simple* resurgent function?
 - Maxim believes it is in general, and indeed in our examples we get simple resurgent functions. But how to prove it in general?

1.3 Stokes phenomenon

- For Bessel functions, we can see explicitly how solutions jump when the Laplace transform angle crosses a critical value.
- The jump comes from the branch cut difference identity for hypergeometric functions.
- Possible interpretation of the Stokes factors as intersection numbers in Morse–Novikov theory [ask Maxim]

2 The Laplace and Borel transforms

2.1 The Laplace transform

- Action on differential equations.
 - Can we find a way to prove this when the differential operator spits out a function that's not integrable around zero?
- Global picture?

2.2 The Borel transform

- Action on differential equations.
 - No inhomogeneous terms! How is this consistent with the Laplace transform's action? Is there always an inhomogeneous solution with subexponential asymptotics?

3 Third-order exponential integrals

- Reduce to

$$I(z) = \int \exp [-z(u^3 + pu + q)] \, du$$

using change of coordinate.

- When $p \neq 0$, can reduce further to

$$I(z) = p^{1/2} e^{-qz} K_{1/3}(p^{3/2}z).$$

- As p goes to zero, $I(z)$ degenerates to

$$\left(\frac{1}{2}\right)^{2/3} e^{-qz} \Gamma\left(\frac{1}{3}\right) z^{-1/3} = \left(\frac{1}{2}\right)^{2/3} e^{-qz} \mathcal{L}_{\zeta,0}(\zeta^{-2/3}) = \left(\frac{1}{2}\right)^{2/3} \mathcal{L}_{\zeta-q,q}(\zeta^{-2/3}).$$