

Additional Information Table

Patient Information

Name:	Jitendra Bangare	Age:	40
Ref Dr:	Dr. Tina Pramanik	Patient ID:	6072026
Test By:	Sambodhi Awchar	Test Date:	06-Jul-2026 17:50

Sample Information - #6072026

Sample Type:	MASTRUBATION	Volume(ml):	1
Liquefaction Duration:	15	Viscosity:	EQUIVOCAL
pH:	7.5	Sperm conc(raw):	50

Parameter	Result (%)	Reference Chart
DNA Fragmentation Index SCD - Sperm Chromatin Dispersion	69.5%	A bar chart with four bars representing different levels of DNA fragmentation. The first bar, labeled 'Pt. Score', is blue and reaches 69.5%. The second bar, labeled 'Excellent', is green and is below 15%. The third bar, labeled 'Good', is orange and is at 25%. The fourth bar, labeled 'Poor', is red and is above 25%.

Parameter	Count	Percentage (%)	Analyzed Chart
Non Fragmented	61	30.5%	A donut chart showing the distribution of sperm DNA fragmentation. The blue segment represents 'Non Fragmented' sperm at 30.5%, and the red segment represents 'Fragmented' sperm at 69.5%.
Fragmented	139	69.5%	
Total	200	100%	

DNA Fragmentation Index References

Sperm DNA Fragmentation (SDF) is a part of Semen Analysis patient profile and is the optimal approach to obtain a comprehensive clinical assessment. Coupled with a personal history and physical assessment, the physician can more readily diagnose a problem and establish a highly effective strategy to treat male factor infertility. DNA Fragmentation Index is a useful parameter to identify male subfertility as the etiological factor for repeated miscarriage and embryonic abnormalities.

DNA fragmentation is based on the Sperm Chromatin Dispersion (SCD) test.

(Fernandez et al, J Androl 24:59–66, 2003; Fertil Steril 84:833–842)

Infertile patients usually have DFI > 25%

Clinical achievement: IUI – 7%, IVF – 16%, ICSI – 33%


The DFI value must correspond with all the other clinical and laboratory findings of the respective semen sample.



The test offers presumable quality information of DNA fragmentation of sperm.

This test was done with QwikCheck DFI

Additional Information Table

DNA Fragmentation Halo Classification

Parameter		Count	Percentage (%)	Analyzed Chart
	LARGE HALO	22	11%	
	MEDIUM HALO	39	19.5%	
	SMALL HALO	80	40%	
	NO HALO	45	22.5%	
	DEGRADED	14	7%	
Total		200	100%	



DNA Fragmentation Index References

Sperm DNA Fragmentation (SDF) is a part of Semen Analysis patient profile and is the optimal approach to obtain a comprehensive clinical assessment. Coupled with a personal history and physical assessment, the physician can more readily diagnose a problem and establish a highly effective strategy to treat male factor infertility. DNA Fragmentation Index is a useful parameter to identify male subfertility as the etiological factor for repeated miscarriage and embryonic abnormalities.

DNA fragmentation is based on the Sperm Chromatin Dispersion (SCD) test.

(Fernandez et al, J Androl 24:59–66, 2003; Fertil Steril 84:833–842)

Infertile patients usually have DFI > 25%

Clinical achievement: IUI – 7%, IVF – 16%, ICSI – 33%

The DFI value must correspond with all the other clinical and laboratory findings of the respective semen sample.

The test offers presumable quality information of DNA fragmentation of sperm.

This test was done with QwikCheck DFI