

Travel health Action Sheet

Peru · Lagos, Nigeria → Lima, Peru · 90 days · Leisure
May 11, 2026

SECTION 1: CRITICAL ACTIONS BEFORE DEPARTURE

- Book a pre travel health consultation with a travel medicine physician or travel health clinic immediately. Departure is 21 May 2026 and multiple vaccine series require weeks to complete.
- Begin rabies pre exposure prophylaxis series (3 doses: days 0, 7, and 21) as soon as possible. This is time critical before departure.
- Receive yellow fever vaccination at a designated yellow fever vaccination centre at least 10 days before departure. Confirm Peru entry requirements for travellers arriving from Nigeria.
- Receive hepatitis A vaccination as soon as possible. Accelerated schedule available if needed.

SECTION 2: TRIP SNAPSHOT

Alone, 90 days, Leisure, overall risk HIGH

SECTION 3: VACCINES

No qualifying vaccines listed in dossier.

SECTION 4: WHAT TO PACK AND DAILY ROUTINE

- Malaria prophylaxis: Atovaquone proguanil (Malarone) or doxycycline, subject to clinician review. Malaria is present in rural and jungle regions of Peru, including areas accessible from Lima. Both *Plasmodium vivax* and *P. falciparum* are present. The traveller is staying in a rural environment for 90 days with planned hiking and outdoor activities, representing sustained high risk exposure. Atovaquone proguanil is well tolerated and suitable for a 90 day course, though cost over this duration should be discussed with your clinician. Doxycycline is a cost effective alternative but requires strict sun protection (photosensitivity risk) and consistent daily dosing. Mefloquine is an option but is not preferred given the lack of prior use history and the availability of better tolerated alternatives. Chloroquine is not effective in Peru due to resistance. Chemoprophylaxis must begin before arrival in a malaria risk area, continue throughout, and for 7 days (atovaquone proguanil) or 4 weeks (doxycycline or mefloquine) after leaving the risk area. Your clinician will confirm the most appropriate agent based on your full health profile.
- Pre travel review and vaccination records: Book a pre travel health consultation as soon as possible given the 21 May 2026 departure date. Multiple vaccines are required, some needing 3 doses over 21 days (rabies pre exposure prophylaxis) and others requiring at least 10 days before travel (yellow fever). Bring all existing vaccination records to the consultation. Confirm hepatitis A, hepatitis B, typhoid, yellow fever, and rabies vaccination schedules with your clinician immediately. Obtain a physician letter documenting your health status and any medications for use at border crossings and in medical emergencies.
- Food and water hygiene: In rural Peru, treat all water as potentially contaminated. Drink only bottled water with intact seals, or water that has been boiled or treated with iodine or chlorine tablets. Avoid ice in drinks unless you are certain it was made from safe water. Eat only thoroughly cooked food served hot. Avoid raw or undercooked meat, seafood, and eggs. Avoid salads, raw vegetables, and fruit that cannot be peeled. Be cautious with street food. Wash hands thoroughly with soap and water before eating and after using the toilet. Carry alcohol based hand sanitiser (at least 60 percent alcohol) for situations where

handwashing is not possible. Discuss standby antibiotic therapy (for traveller's diarrhoea) with your clinician before departure. Azithromycin is typically recommended for travellers from Nigeria to Peru.

- Vector bite prevention: Malaria, dengue, yellow fever, and other vector borne diseases are present in rural Peru. Apply DEET repellent (30 to 50 percent) to all exposed skin every 4 to 6 hours. Wear permethrin treated long clothing during outdoor activities. Use a treated bed net every night. Dengue has no prophylactic medication: bite prevention is your only protection. Be aware that dengue transmitting Aedes mosquitoes are active during the day, unlike malaria transmitting Anopheles mosquitoes which are most active at dusk and dawn. Both require consistent protection throughout the day and night.
- Sun, heat, and environmental precautions: Peru's UV index is very high, particularly at altitude and during outdoor activities. Apply SPF50 plus broad spectrum sunscreen to all exposed skin and reapply every 2 hours and after swimming. Wear a wide brimmed hat and UV protective sunglasses. Stay hydrated with safe water, especially during hiking. Avoid prolonged sun exposure between 10:00 and 14:00. If your hiking itinerary takes you to Andean regions above 2,500 metres, ascend gradually to allow acclimatisation. Symptoms of acute mountain sickness include headache, nausea, dizziness, and fatigue. Discuss acetazolamide (Diamox) prophylaxis with your clinician if high altitude travel is planned. Lima's air quality can be poor: if you have any respiratory sensitivity, carry a suitable mask.
- Injury and road safety: Peru has a high rate of road traffic accidents. Avoid travelling by road at night where possible. Use reputable transport providers. Always wear a seatbelt. Do not ride motorcycles or motorbike taxis without a helmet. During hiking, use appropriate footwear, stay on marked trails, and inform someone of your planned route and expected return time. When swimming, only swim in designated safe areas, be aware of currents, and never swim alone in remote locations. Carry a basic first aid kit including wound care supplies, antiseptic, and blister treatment. Ensure your travel insurance covers emergency medical evacuation from rural areas.
- Sexual health and blood exposure: You have indicated you anticipate new sexual partners during this trip. Use condoms consistently and correctly for every sexual encounter to reduce the risk of HIV, hepatitis B, gonorrhoea, chlamydia, syphilis, and other STIs. Condom use significantly reduces but does not eliminate all STI risk. Confirm your hepatitis B vaccination status before departure and complete the series if not immune. Discuss HIV pre exposure prophylaxis (PrEP) with your clinician before travel: PrEP is highly effective at preventing HIV when taken correctly. Know how to access HIV post exposure prophylaxis (PEP) in Lima in the event of a potential exposure: PEP must be started within 72 hours. Avoid tattooing, piercing, or any cosmetic procedures abroad unless you can verify sterile professional facilities. Avoid sharing needles or any equipment that may contact blood.
- Jet lag, sleep, and mental wellbeing: Lagos (WAT, UTC plus 1) to Lima (PET, UTC minus 5) represents a 6 hour time difference. Expect significant jet lag on arrival. Adjust sleep and meal times gradually in the days before departure. On arrival, expose yourself to natural daylight to help reset your circadian rhythm. Maintain regular sleep patterns as quickly as possible. A 90 day solo trip in a rural environment can be isolating. Plan regular contact with family and friends at home. Be aware of signs of culture shock, loneliness, or low mood and have a plan for accessing support if needed. Carry any personal mental health resources or contacts you rely on.

SECTION 5: RED FLAGS AND EMERGENCY CONTACTS

URGENT: Seek emergency medical care immediately for any of the following after returning from Peru: high fever with chills or rigors (possible malaria), severe headache with neck stiffness or rash (possible meningitis), difficulty breathing or chest pain, jaundice (yellow skin or

eyes), altered consciousness or confusion, severe diarrhoea with blood or signs of dehydration, or any animal bite that was not assessed during the trip. Always tell the treating doctor you have returned from rural Peru.

CLINICAL FACILITIES

Clinica Internacional, Jr. Washington 1471, Lima 15001, Peru

Clinica Anglo Americana, Av. Salazar, San Isidro, Lima 15073, Peru

LOCAL EMERGENCY NUMBER

+51 1 460 1060

INSURANCE

Insurance: Yes

Full clinical dossier attached. Share with your travel medicine clinician at your pre-travel appointment.