

Pre-Travel Health Advisory

United States · 7 days · Leisure

Departing 5 December 2026

Traveller: Magid Cleos

Low risk

COMPLETED

Apr 24, 2026

TRIP AT A GLANCE	
DURATION	7 days
PURPOSE	Leisure tourism
TRAVELLING	Alone
ACCOMMODATION	Hotel (urban)
INSURANCE	Not purchased

ITINERARY-SPECIFIC GUIDANCE		
Trip type: ONE_WAY		
Magid is travelling one-way from Lagos, Nigeria to Manhattan, United States, departing 5 December 2026 for a 7-day leisure trip. The trip involves a 6-hour longest flight leg and 10 hours total flying time. Accommodation is hotel-based in an urban setting. Planned activities include hiking and outdoor pursuits.		
ROUTE GUIDANCE BY STOP		
STOP	COUNTRY	GUIDANCE
Manhattan	United States	Manhattan in December is cold (1 to 8 degrees Celsius). Pack warm, layered clothing and waterproof footwear. December is peak respiratory virus season — influenza and COVID-19 vaccines should be current before arrival. Healthcare in the United States is extremely expensive without insurance; travel insurance with comprehensive medical cover is essential and must be purchased before departure. Planned hiking activities should only proceed if surgical clearance has been obtained from Magid's treating physician following his recent appendectomy. Emergency services are reached by dialling 911. The Nigerian Consulate General is located at 828 Second Avenue, New York, and can provide consular assistance if needed.

HEALTH RISK OVERVIEW		
CATEGORY	LEVEL	SUMMARY
Food and water safety	LOW	Manhattan has a treated municipal water supply and regulated food safety standards. Standard hygiene precautions apply, but the risk of food or waterborne illness is low for a hotel-based urban stay.
Vector-borne diseases	LOW	Manhattan in December presents negligible vector-borne disease risk. Mosquito and tick activity is minimal in winter months in New York City. No malaria, dengue, or yellow fever risk at this destination.

Respiratory infections	MODERATE	December is peak respiratory virus season in the northeastern United States. Influenza, COVID-19, RSV, and other respiratory pathogens circulate actively. Crowded indoor environments such as transit hubs, hotels, and tourist attractions elevate exposure risk. Travelling from Nigeria adds a transition from a different seasonal pattern.
Environmental health (heat, sun, air quality)	LOW	December in Manhattan is cold, with average temperatures between 1 and 8 degrees Celsius. Heat illness is not a concern. Cold weather exposure, particularly for a traveller from Lagos, warrants appropriate clothing. Air quality in New York City is generally acceptable.
Injuries and road traffic safety	MODERATE	New York City road traffic is dense. Pedestrian safety requires vigilance, particularly given left-hand traffic patterns differing from Nigeria. Planned hiking and outdoor activities carry a low-to-moderate injury risk depending on terrain and conditions.
Rabies and animal contact	LOW	The United States is not rabies-free, but urban Manhattan presents very low practical risk of rabies exposure. Bat contact is the primary urban rabies vector. No pre-exposure prophylaxis is routinely recommended for a short urban leisure trip. Standard advice: avoid contact with wild or stray animals.
Blood-borne and sexual health	MODERATE	Magid has a documented STI history. Although no risk behaviours are anticipated for this trip, the STI history warrants a pre-travel sexual health review and confirmation of hepatitis B immune status. Standard precautions apply.
Altitude-related illness	LOW	Manhattan is at sea level. No altitude-related illness risk applies to this destination. This category is not clinically relevant for this trip.

FLIGHT & JOURNEY HEALTH

VTE Risk Level: **MEDIUM**

PREVENTION MEASURES

- Longest flight leg is 6 hours and total flying time is 10 hours, placing this in the medium VTE (venous thromboembolism, blood clot) risk category
- Wear graduated compression stockings (15 to 30 mmHg) for all flight legs of 4 hours or more
- Perform regular in-seat leg exercises: ankle circles, calf raises, and knee lifts every 30 to 60 minutes
- Walk the aisle for 5 minutes every 1 to 2 hours when the seatbelt sign is off
- Stay well hydrated throughout the flight; avoid excessive alcohol and caffeine which promote dehydration
- Avoid sleeping in positions that compress the legs for extended periods
- Recent appendix surgery is a potential additional VTE risk factor — confirm with surgeon whether LMWH (low molecular weight heparin, a blood-thinning injection) is recommended for this flight given the surgical history

AIRLINE MEDICAL CLEARANCE REQUIRED

Contact your airline to arrange medical clearance before travel.

MEDICATION TIMING

No current medications reported. If any medications are commenced before travel (e.g. vaccines, prophylaxis), carry all in original packaging with a physician letter.

VACCINATIONS

VACCINE	STATUS	GUIDANCE
Routine immunizations (e.g. MMR, varicella, dTdap, polio /IPV)	Believed up to date (self-reported)	Recommended — verify documentation before travel Magid reports believing his routine vaccinations are up to date but has no documented travel-related vaccines. Confirm MMR (2-dose series), varicella, dTdap (tetanus-diphtheria-pertussis booster if more than 10 years since last dose), and polio/IPV status with his primary care physician before departure. Hiking and outdoor activities increase the relevance of a current tetanus booster.
Influenza	Unknown	Strongly recommended December is peak influenza season in the United States. A current-season influenza vaccine is strongly recommended before departure. This is particularly relevant given travel from Nigeria, where seasonal patterns differ, and planned exposure to crowded urban environments.
COVID-19	Unknown	Recommended Ensure COVID-19 vaccination is up to date with the most current booster available before travel. Check US entry requirements at the time of travel, as these may be updated. Respiratory virus season in December makes current vaccination especially relevant.
Hepatitis A	Unknown — no travel vaccines previously received	Recommended Hepatitis A vaccination is recommended for travellers from Nigeria to the United States as a general travel health measure, and to ensure protection is in place for future travel. A single dose provides protection within 2 to 4 weeks; a second dose at 6 to 12 months provides long-term immunity. Discuss with your physician or travel clinic.
Hepatitis B	Unknown — no travel vaccines previously received	Recommended — elevated priority given STI history Hepatitis B vaccination is recommended. Given Magid's documented STI history, confirming hepatitis B immune status (via blood test) and completing the vaccine series if not immune is a clinical priority. An accelerated schedule (doses at day 0, day 7, and day 21 with a booster at 12 months) is available if the pre-travel window is short.
Typhoid	Not indicated	Not recommended for this destination Manhattan has a treated municipal water supply and regulated food safety infrastructure. Typhoid vaccination is not recommended for this trip per CDC and WHO guidelines. No action required.
Yellow fever	Not indicated	Not required or recommended for the United States Yellow fever is not present in the United States and is not an entry requirement. No action required for this destination. Note: if Magid plans

		future travel to yellow fever endemic regions, vaccination should be arranged in advance.
Japanese encephalitis	Not indicated	Not recommended for this destination Japanese encephalitis is not present in the United States. No action required for this trip.
Meningococcal	Unknown	Consider — discuss with physician Meningococcal vaccination (MenACWY) is not a routine requirement for leisure travel to Manhattan. However, if Magid's routine vaccination records are incomplete or if he plans to visit university campuses or crowded venues, discuss with his physician whether a dose is appropriate. MenACWY is part of routine adolescent schedules in many countries.
Rabies pre-exposure	Not indicated	Not recommended for this trip Pre-exposure rabies prophylaxis is not recommended for a short urban leisure trip to Manhattan. The practical risk of rabies exposure in this setting is very low. Standard advice applies: avoid contact with wild or unfamiliar animals, including bats.
Cholera (oral vaccine)	Not indicated	Not recommended for this destination Cholera is not a risk in Manhattan. The oral cholera vaccine is not recommended for this trip. No action required.

MEDICAL ADVICE & RECOMMENDATIONS	
TOPIC	DETAILS
Pre-travel review and vaccination records	Magid has not received any documented travel-related vaccines and reports believing his routine vaccinations are up to date without formal verification. Before departure on 5 December 2026, he should visit his primary care physician or a travel clinic in Lagos to: (1) verify and document routine vaccine status (MMR, varicella, dTdap, polio/IPV); (2) receive influenza and COVID-19 vaccines; (3) discuss hepatitis A and hepatitis B vaccination given his travel history and STI history; and (4) obtain a physician letter documenting his recent appendix surgery (within the last 12 months) confirming fitness to travel. Travel insurance has not been purchased — this must be addressed urgently before departure (see below).
Food and water hygiene	Manhattan has a safe, treated municipal water supply. Tap water is safe to drink. Standard food hygiene applies: wash hands before eating, choose reputable food establishments, and avoid raw or undercooked foods if there is any doubt about preparation standards. The risk of traveller's diarrhoea in Manhattan is low. No standby antibiotics are indicated for this destination.
Vector bite prevention	December in Manhattan presents negligible mosquito or tick activity. No specific vector bite prevention measures are required for this trip. If Magid undertakes hiking or outdoor activities in green spaces or parks, he should check for ticks after outdoor activity as a general precaution, though the risk in December is very low.

Sun, heat, and environmental precautions	December in Manhattan is cold, with temperatures typically between 1 and 8 degrees Celsius and potential for wind chill, rain, and occasional snow. Magid is travelling from Lagos, where temperatures are significantly warmer. He should pack appropriate cold-weather clothing including a warm coat, layers, waterproof footwear, and gloves. Cold exposure can be a genuine health risk for travellers unaccustomed to winter conditions. UV index is low in December in New York, but SPF protection is still advisable on bright days, particularly during outdoor activities.
Injury and road safety	New York City has dense road traffic. Pedestrians should use designated crossings and be aware that traffic patterns and road rules differ from Nigeria. Planned hiking and outdoor activities should be undertaken on marked trails with appropriate footwear. Given Magid's recent appendix surgery within the last 12 months, he should confirm with his surgeon that strenuous hiking is safe before departure. Carry a basic first aid kit for outdoor activities. Emergency services in the United States are accessed by dialling 911.
Sexual health and blood exposure	Magid has a documented STI history. Although no risk behaviours are anticipated for this trip, a pre-travel sexual health review is recommended, including confirmation of hepatitis B immune status. If sexual activity occurs during travel, consistent use of barrier protection (condoms) is strongly advised to reduce the risk of STIs including HIV, gonorrhoea, chlamydia, syphilis, and hepatitis B. PrEP (pre-exposure prophylaxis for HIV) is widely available in the United States and can be discussed with a physician if relevant. Avoid tattooing, piercing, or any invasive cosmetic procedures unless sterile professional facilities are confirmed.
Jet lag, sleep, and mental wellbeing	The time difference between Lagos (WAT, UTC+1) and New York (EST, UTC-5) is 6 hours. Magid will be travelling westward, which generally results in difficulty falling asleep at the local bedtime. Strategies to minimise jet lag include: adjusting sleep and meal times gradually before departure, seeking natural light exposure on arrival, staying hydrated during the flight, and avoiding excessive alcohol or caffeine during transit. Travelling alone to a new country can be disorienting; having a clear plan for accommodation, local transport, and emergency contacts will support wellbeing.
Malaria and other chemoprophylaxis (state if not indicated)	Malaria prophylaxis is NOT indicated for this trip. Manhattan, United States, is not a malaria-endemic area. No antimalarial medication is required. Note: Magid is travelling from Lagos, Nigeria, which is a malaria-endemic region. If he has had any recent fever or flu-like symptoms before departure, he should seek medical review in Lagos before travelling, as malaria should be excluded.
Traveller-specific considerations (from health context)	Several specific factors require attention for this traveller: (1) Recent appendix surgery within the last 12 months — Magid must obtain a physician letter confirming fitness to travel and fitness for planned hiking activities before departure. Depending on the timing and nature of the surgery, there may be restrictions on strenuous physical activity or long-haul flying. (2) Travel insurance — Magid has not purchased travel insurance for this trip. This is a significant gap. Medical care in the United States is among the most expensive in the world. A single emergency department visit can cost thousands of US dollars. Comprehensive travel insurance with medical evacuation cover is strongly recommended and should be purchased before departure. (3) STI history — pre-travel sexual health review and hepatitis B status confirmation are recommended. (4)

Travelling alone — Magid should ensure he has emergency contact details, knows how to access medical care in Manhattan, and has a clear plan for what to do in a medical emergency.

MEDICAL CONDITIONS & PRECAUTIONS

CONDITION	PRECAUTIONS
Recent appendix surgery (appendectomy within last 12 months)	Magid reports hospitalisation for appendix surgery within the last 12 months. The timing and nature of this surgery are not specified in the questionnaire. Before travel, he must obtain written clearance from his treating surgeon or physician confirming: (1) fitness to fly on a 6-hour flight leg with 10 hours total flying time; (2) fitness to undertake hiking and outdoor activities; and (3) that there are no post-operative complications (such as hernia, adhesions, or wound issues) that could be exacerbated by travel or physical activity. Post-surgical patients may have elevated VTE risk — the surgeon should advise on whether compression stockings alone are sufficient or whether LMWH prophylaxis is warranted for the flight. A physician letter should be carried throughout the trip in case medical attention is required in the United States.

MEDICATION LOGISTICS

PACKAGING	Carry all medications in original packaging with labels intact. Obtain a physician letter listing all medications by generic and brand name if any are commenced before travel.
SUPPLY RULE	Carry sufficient supply for the full trip duration plus a minimum 3-day buffer for delays. Split supply between carry-on and checked luggage where possible.
DESTINATION LEGALITY VERIFIED	Yes — confirm all medications are legal at destination.

CLINICAL DECISION FLAGS

- TREE_3_MEDIUM_RISK_DESTINATION
- TREE_4_DURATION
- TREE_5_FLIGHT
- Recent serious illness or surgery within 12 months (appendectomy) — fitness to travel and fitness for hiking activities must be confirmed by treating physician before departure
- No travel insurance purchased — critical gap for travel to the United States where healthcare costs are extremely high
- STI history documented — pre-travel sexual health review and hepatitis B immune status confirmation recommended
- No travel-related vaccines previously received — full pre-travel vaccine review required
- Substance use and adherence risk: prefer not to say — flag for clinician to explore medication adherence and risk behaviour counselling in consultation

SPECIALIST REFERRALS

CONDITION	SPECIALIST	URGENCY
Recent appendix surgery within 12 months	Treating surgeon or general practitioner	BEFORE_TRAVEL
STI history and unknown hepatitis B immune status	General practitioner or sexual health clinic	BEFORE_TRAVEL
No travel vaccines received and routine vaccine status unverified	Travel medicine clinic or general practitioner	BEFORE_TRAVEL

SEXUAL HEALTH & RISK BEHAVIOURS

Risk Level: MODERATE

PREVENTION ADVICE

- Magid has a documented STI history. A pre-travel sexual health review with his physician or a sexual health clinic in Lagos is recommended before departure.
- Confirm hepatitis B immune status via blood test. If not immune, commence hepatitis B vaccination series before travel.
- If sexual activity occurs during travel, use barrier protection (condoms) consistently and correctly for every encounter to reduce the risk of HIV, gonorrhoea, chlamydia, syphilis, and hepatitis B.
- Be aware that STI rates, including antibiotic-resistant gonorrhoea, are present in urban US settings. Symptoms such as unusual discharge, sores, or pain should prompt prompt medical review.
- Post-travel STI screening is recommended on return to Nigeria if any sexual activity occurred during the trip.
- PrEP (pre-exposure prophylaxis for HIV prevention) is widely available in the United States and can be discussed with a physician if considered relevant.

PREP/PEP

PrEP/PEP availability discussed for high-risk destination.

AFTER YOU RETURN

RED FLAGS — SEEK IMMEDIATE CARE IF YOU EXPERIENCE

Seek immediate medical attention if any of the following occur after return: high fever (above 38 degrees Celsius), severe abdominal pain, chest pain or difficulty breathing, signs of deep vein thrombosis (leg swelling, redness, or pain), or any neurological symptoms. Always inform the treating clinician of recent international travel.

WITHIN 1 WEEK

- Monitor for any signs of post-surgical complications that may have been aggravated by travel or physical activity, including abdominal pain, swelling, or fever. Seek medical review promptly if any of these occur.
- If any respiratory symptoms develop (fever, cough, sore throat), seek medical review and mention recent travel to the United States during peak respiratory virus season.
- Check for any skin changes, rashes, or insect bites noted during outdoor activities.

WITHIN 4 WEEKS

- If any sexual activity occurred during travel, attend a sexual health clinic for STI screening within 2 to 4 weeks of return.

If fever develops within 4 weeks of return, seek urgent medical attention and inform the clinician of recent travel. Although malaria is not a risk at the destination, Magid is returning to Lagos from a trip and any febrile illness should be assessed.
Review travel insurance claim if any medical expenses were incurred during the trip.
BEYOND 4 WEEKS
Complete any commenced vaccine series (hepatitis A second dose at 6 to 12 months, hepatitis B booster at 12 months if accelerated schedule was used).
Schedule a routine follow-up with his primary care physician to review overall health and travel experience.
If any persistent symptoms developed during or after travel, ensure these are fully investigated.

MEDICAL CARE & CONTACTS		
CLINICS & FACILITIES		
NAME		LOCATION & NOTES
NYU Langone Health Urgent Care — Midtown Manhattan		317 East 34th Street, New York, NY 10016 +1 646 929 7800 · Midtown Manhattan
		Walk-in urgent care available. Full emergency services at NYU Langone Medical Center nearby.
Mount Sinai Hospital Emergency Department		1 Gustave L. Levy Place, New York, NY 10029 +1 212 241 6500 · Upper East Side, Manhattan
		Major academic medical centre with full emergency and specialist services.
NewYork-Presbyterian Hospital Weill Cornell Medical Center		525 East 68th Street, New York, NY 10065 +1 212 746 5454 · Upper East Side, Manhattan
		Major tertiary hospital. Emergency department available 24 hours.
EMBASSY & CONSULAR		
NAME	DETAILS	
Nigerian Consulate General, New York	828 Second Avenue, New York, NY 10017. Phone: +1 212 808 0301. Provides consular assistance to Nigerian nationals in the United States.	
EMERGENCY NUMBERS		
LABEL		CONTACT
US Emergency Services (Police, Fire, Ambulance)		911
NYC Health and Hospitals General Information		+1 212 788 3321
Nigerian Consulate General New York (Emergency Line)		+1 212 808 0301

NEXT STEPS
URGENT: Purchase comprehensive travel insurance with medical evacuation cover before departure on 5 December 2026. Medical care in the United States without insurance can cost tens of thousands of US dollars.

- **URGENT:** Obtain written fitness-to-travel and fitness-for-hiking clearance from your treating surgeon or physician regarding your recent appendectomy. Ask specifically whether LMWH (blood-thinning injection) is recommended for your 6-hour flight given your surgical history.
- Visit a travel clinic or your primary care physician in Lagos as soon as possible before departure to: verify and document routine vaccine status; receive influenza and COVID-19 vaccines; discuss hepatitis A and hepatitis B vaccination; and obtain a physician letter for travel.
- Attend a sexual health clinic or discuss with your physician: confirm hepatitis B immune status and commence vaccination if not immune; consider pre-travel STI screening given your STI history.
- Pack appropriate cold-weather clothing for Manhattan in December. Temperatures will be significantly colder than Lagos.
- Carry a copy of all medical documents including your physician fitness-to-travel letter, vaccine records, and any insurance documents in both your carry-on bag and as a digital copy accessible on your phone.
- Wear graduated compression stockings (15 to 30 mmHg) for all flight legs of 4 hours or more to reduce VTE risk.
- Save emergency contact numbers before departure: US emergency services 911, Nigerian Consulate General New York +1 212 808 0301.

This advisory has been generated by the TMAG (Travel Medicine Advisory Global) AI Clinical Decision Support System and is intended to inform a pre-travel health consultation with a qualified clinician. It does not constitute medical advice and must not replace a consultation with a licensed physician, travel medicine specialist, or pharmacist. All vaccine and medication recommendations require clinician review and a valid prescription where applicable. Information is based on CDC and WHO travel health guidelines current at the time of generation; destination risk profiles may change. Always check for the latest alerts before departure. Travellers with complex medical histories, chronic conditions, or pregnancy are strongly advised to seek specialist travel medicine review in person before travelling. Medication guidance in this report is advisory only and must not be used to alter, substitute, or discontinue any medication currently prescribed to you. Please consult your physician or pharmacist before acting on any medication recommendation. Copyright 2026 Travel Medicine Advisory Global. travelmedicine.global

Travel Medicine Advisory Global — not a substitute for professional medical advice.

DOCTOR VERIFICATION & DIGITAL SIGNATURE

This travel health plan has been reviewed and approved by a licensed physician on the TMAG Doctor Network.



PHYSICIAN NAME	Mat Smith
LICENSE NUMBER	MT898320084
VALIDATION DATE	24 April 2026
STATUS	TMAG Verified & Approved

Digital Signature

TMAG Verified Seal

This document was generated by Travel Medicine Advisory Global and approved by a licensed physician.