

FYS3150 - Project 2

Vegard Falmår and Sigurd Sørli Rustad
Universitetet i Oslo*
(Dated: September 29, 2020)

I. INTRODUCTION

The harmonic oscillator (HO) potential appears in many areas within quantum physics. This is because many potentials can be approximated, at least in some intervals, with a HO potential. The most famous example of this might be the model for a hydrogen atom, where the electron is bound to the proton by a HO potential. That is why, in this report, we are going to study both one and multiple electrons in a HO potential. We are also going to account for electrical forces between the particles.

When we try to solve Schrodinger's equation for both one and two particles, we end up with a eigenvalue problem (see theory section). This can be represented with a linear set of equations written on matrix-form (from project 1 [3]). Using Jacobi's rotation algorithm, also described in the theory section, we can diagonalize the matrix to find these eigenvalues and eigenvectors. Then plot the results, comparing them to experimental values.

When implementing algorithms it is important to test your code and make sure it performs as expected. Therefore we perform unit-tests to make sure sections of our program works as expected. We also test the code with a test scenario (buckling beam), which have analytical eigenvalues and eigenvectors. The testing is described in the appendix (VII).

For our studies we have used c++ for heavy computation, python for visualization and bash for automation. All the code along with instructions on how to run it, can be cloned from our GitHub repository here [4].

II. THEORY

A. Unitary transformation

The transposed of a unitary matrix (U) is its inverse.

$$U^T = U^{-1}$$

From this we can prove that a unitary transformation preserves the orthonormality of vectors. Consider the set of orthonormal vectors $\{\mathbf{v}_i\}_i$ and the unitary trans-

formation $\{U\mathbf{v}_i\}_i = \{\mathbf{w}_i\}_i$.

$$\begin{aligned}\mathbf{w}_i^T \mathbf{w}_j &= (U\mathbf{v}_i)^T U\mathbf{v}_j \\ &= \mathbf{v}_i^T U^T U\mathbf{v}_j = \mathbf{v}_i^T \mathbf{v}_j \\ &= \delta_{i,j}\end{aligned}$$

We notice that orthonormality is perserved.

B. Jacobi's rotation algorithm

Jacobi's rotation algorithm uses unitary transformations to diagonalize a matrix. A detailed description of the algorithm can be found here [1], however we will describe it briefly. In order to diagonalize a given matrix A , as mentioned over, we perform a series of unitary transformations.

$$B = U_n^T U_{n-1}^T \dots U_0^T A U_0 \dots U_{n-1} U_n$$

Here U_i are the unitary matrices and B the resulting diagonal matrix. An example of how U_i can look is given under.

$$\begin{bmatrix} 1 & 0 & \dots & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & 0 & \dots \\ 0 & 0 & \dots & \cos \theta & 0 & \dots & 0 & \sin \theta \\ 0 & 0 & \dots & 0 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & 1 & \dots \\ 0 & 0 & \dots & -\sin \theta & 0 & \dots & 0 & \cos \theta \end{bmatrix}$$

The geometric interpretation is that U_i performs a rotation on T in order to zero out non-diagonal elements. It turns out that the fastest way to do this, is to zero out the largest non-diagonal matrix-element. First We define:

$$\cot(\theta) = \tau = \frac{a_{ll} - a_{kk}}{2a_{kl}},$$

Where a_{kl} is the largest non-diagonal element in A . Now to shorten notation we use $\tan(\theta) = t = s/c$, where $s = \sin(\theta) \wedge c = \cos(\theta)$. By defining θ such that a_{kl} becomes zero we get the quadratic equation

$$t^2 + 2\tau t - 1 = 0 \implies t = -\tau \pm \sqrt{1 + \tau^2},$$

and can also obtain c and s

$$c = \frac{1}{1 + t^2} \wedge s = tc.$$

* vegardfa@uio.no; sigurdsr@gmail.com

The actual transformation is defined by the set of equations

$$\begin{aligned} b_{ik} &= a_{ik}c - a_{il}s, i \neq k, i \neq l \\ b_{il} &= a_{il}c + a_{ik}s, i \neq k, i \neq l \\ b_{kk} &= a_{kk}c^2 - 2a_{kl}cs + a_{ll}s^2 \\ b_{ll} &= a_{ll}c^2 + 2a_{kl}cs + a_{kk}s^2 \\ b_{kl} &= (a_{kk} - a_{ll})cs + a_{kl}(c^2 - s^2) \end{aligned} \quad (1)$$

Where b_{ij} are elements in B , again see [1] for a more detailed description of the algorithm.

An interesting question is; what happens to the eigenvectors and eigenvalues? Because B is diagonal the eigenvectors are just vectors with a 1 in one element and 0 everywhere else. We consider an arbitrary eigenvector \mathbf{v} and eigenvalue λ that satisfy

$$B\mathbf{v} = \lambda\mathbf{v}.$$

Applying the unitary transforms we get

$$\begin{aligned} B\mathbf{v} &= U_n^T U_{n-1}^T \dots U_0^T A U_0 \dots U_{n-1} U_n \mathbf{v} \\ &= \lambda\mathbf{v} \end{aligned}$$

We notice that this is true if

$$\mathbf{v}' = U_0 \dots U_{n-1} U_n \mathbf{v} \quad (2)$$

is an eigenvector with eigenvalue λ that satisfy

$$A\mathbf{v}' = \lambda\mathbf{v}'$$

Therefore, if you have the eigenvector \mathbf{v} and matrices U_i you can find \mathbf{v}' with equation (2).

C. Electrons in a harmonic oscillator potential

It turns out that if you try to solve Schrodinger's equation you end up with a eigenvalue problem. We will derive the math here, but everything is taken from here [2]. First we consider the Schrodinger's equation for one electron. Because of spherical symmetry we only need to look at the radial part.

$$-\frac{\hbar^2}{2m} \left(\frac{1}{r^2} \frac{d}{dr} r^2 \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) R(r) + V(r)R(r) = ER(r) \quad (3)$$

Where $R(r)$ is the radial wave equation, $V(r)$ the potential, l orbital momentum and r distance from the origin. Now because we only have one electron we end up with the HO potential $V(r) = (1/2)kr^2$. Substituting $R(r) = (1/r)u(r)$ and $\rho = (1/\alpha)r$ (where α is a constant with dimension length) we obtain

$$-\frac{\hbar^2}{2m\alpha^2} \frac{d^2}{d\rho^2} u(\rho) + \left(V(\rho) + \frac{l(l+1)}{\rho^2} \frac{\hbar^2}{2m\alpha^2} \right) u(\rho) = Eu(\rho).$$

Setting the orbital momentum $l = 0$, inserting $V(\rho) = (1/2)k\alpha^2\rho^2$ and defining

$$\alpha \equiv \left(\frac{\hbar^2}{mk} \right)^{1/4} \wedge \lambda \equiv \frac{2m\alpha^2}{\hbar} E$$

Schrodinger's equation can then be described by the simple equation (4).

$$-\frac{d^2}{d\rho^2} u(\rho) + \rho^2 u(\rho) = \lambda u(\rho) \quad (4)$$

Now we can expand our model to two particles with Coulomb potential between them. First introducing the relative position \mathbf{r} and center-of-mass coordinate \mathbf{R}

$$\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2 \wedge \mathbf{R} = \frac{1}{2(\mathbf{r}_1 + \mathbf{r}_2)}.$$

Where \mathbf{r}_1 and \mathbf{r}_2 are the particles positions. Then the radial Schrodinger's becomes

$$\left(-\frac{\hbar^2}{m} \frac{d^2}{dr^2} - \frac{\hbar^2}{4m} \frac{d^2}{dR^2} + \frac{1}{4}kr^2 + kR^2 \right) u(r, R) = E^{(2)}u(r, R).$$

$E^{(2)}$ denotes the fact that we are looking at the total energy of two particles. From an ansatz for the wave equation, we can perform separation of variable on r and R .

$$u(r, R) = \psi(r)\phi(R) \wedge E^{(2)} = E_r + E_R$$

Adding the Coulomb potential

$$\frac{\beta e^2}{r}$$

between the two electrons we get

$$\left(-\frac{\hbar^2}{m} \frac{d^2}{dr^2} + \frac{1}{4}kr^2 + \frac{\beta e^2}{r} \right) \psi(r) = E_r \psi(r).$$

Again introducing $\rho = r/\alpha$ and defining a few variables

$$\omega_r^2 \equiv \frac{1}{4} \frac{mk}{\hbar^2} \alpha^4 \wedge \alpha \equiv \frac{\hbar^2}{m\beta e^2} \wedge \lambda \equiv \frac{m\alpha^2}{\hbar^2} E$$

We arrive at the Schrodinger's equation (5)

$$-\frac{d^2}{d\rho^2} \psi(\rho) + \omega_r^2 \rho^2 \psi(\rho) + \frac{1}{\rho} \psi(\rho) = \lambda \psi(\rho) \quad (5)$$

III. METHOD

As mentioned in the theory section, when you study one or more particles in a HO potential, you get a eigenvalue problem. For one particle this is given by equation (4). From Project 1 (see the theory section in [3]) we

know this type of differential equation creates a set of equations, that can be represented as follows.

$$\begin{bmatrix} d_1 & e_1 & 0 & 0 & \dots & 0 & 0 \\ e_1 & d_2 & e_2 & 0 & \dots & 0 & 0 \\ 0 & e_2 & d_3 & e_3 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots & e_{N-2} & e_{N-2} \\ 0 & \dots & \dots & \dots & \dots & e_{N-2} & d_{N-1} \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \dots \\ \dots \\ \dots \\ u_{N-1} \end{bmatrix} = \lambda \begin{bmatrix} u_1 \\ u_2 \\ \dots \\ \dots \\ \dots \\ u_{N-1} \end{bmatrix}$$

Where $e_i = -1/h^2$, $d_i = 2/h^2 + V_i$ and $h = (\rho_N - \rho_0)/N$. Now we can solve this with Jacobi's rotation algorithm described in the theory section. We do this with the sets of equations (1). To make sure algorithm was implemented properly we performed several tests. First unit-tests to make sure small chunks run as expected, then with a buckling beam problem that has analytical eigenvalues and eigenvectors. Further information on the tests can be found in the appendix (VII).

IV. RESULTS




Figure 1.

V. DISCUSSION

VI. CONCLUSION

VII. APPENDIX

A. Unit-tests

B. Buckling beam

-
- [1] http://compphysics.github.io/ComputationalPhysics/doc/pub/eigvalues/html/_eigvalues-bs011.html
 - [2] OPPGAVETEKST REF
 - [3] REF PROJECT 1
 - [4] REF GITHUB




Figure 2.




Figure 3.