

























the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype in children with acute bacterial dysentery [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. The only published study of *S. flexneri* in the United Kingdom was by Smith *et al.* [12], who reported that *S. flexneri* was the most common serotype isolated from patients with acute bacterial dysentery in the United Kingdom in 1987. The serotypes isolated were *S. flexneri* 3, 4, 5, 6, 7, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100.

The aim of this study was to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.

The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom. The study was designed to determine the prevalence of *S. flexneri* in patients with acute bacterial dysentery in the United Kingdom.