TECH WITH SHIVANSH
Learn | Apply | Grow

MODULE 10
STATISTICAL TESTS FOR CATEGORICAL DATA
From Concepts to Real-World Business Decisions

"Better Analysts Make a Smarter Tomorrow..."

MODULE 10
CHI-SQUARE TESTS

From Data to Business Decisions



"Statistics turns data into opportunity."

- Simple Explanations
- Business Examples
- Excel Implementation
- Interview Preparation


> "Chi-square tests help us analyze categorical data and find meaningful patterns, so we can make better data-driven business decisions."

Topics We Will Cover (8 Topics)

Total: 9 Pages

1. Chi-Square Test: Overview — Page 2
2. Chi-Square Goodness of Fit Test — Page 3
3. Chi-Square Test of Independence — Page 4
4. Expected Frequency — Page 5
5. Assumptions — Page 6
6. Steps in Chi-Square Test — Page 7
7. Excel Implementation — Page 8
8. Real Life Business Applications & Interview Questions — Page 9

Chi-Square Distribution (Visual Overview)



The chi-square distribution is always right-skewed and used for analyzing categorical data.

Real-World Business Use Cases

- **Marketing** (Customer Preferences)
- **Operations** (Process Effectiveness)
- **HR Analytics** (Attrition & Satisfaction)
- **Finance** (Fraud Detection)
- **Healthcare** (Treatment Outcomes)
- **Product** (User Segments)

- Concepts in Simple Words
- Step-by-Step with Visuals
- Business Examples from Multiple Industries
- Excel Implementation with Formulas
- Interview Questions with Detailed Answers

TECH WITH SHIVANSH
Learn | Apply | Grow

MODULE 10

STATISTICAL TESTS FOR CATEGORICAL DATA
From Concepts to Real-World Business Decisions

"Better Analysts Make a Smarter Tomorrow..."



1 Chi-Square Test: Overview

Analyzing Categorical Data to Find Meaningful Patterns

What is a Chi-Square Test?

The Chi-Square test is a statistical method used to analyze **categorical data** and check whether the observed frequencies differ significantly from the expected frequencies.

> It helps us find if the difference between **what we observed and what we expected** is large enough to be statistically significant.

When to Use Chi-Square Test?

- When your data is categorical (e.g., gender, product type, region).
- To check if observed frequencies match expected frequencies.
- To test if two categorical variables are independent.
- Commonly used in marketing, operations, HR, healthcare, and finance.

Types of Chi-Square Tests


1. **Chi-Square Goodness of Fit Test** (Check if distribution matches expected distribution)
2. **Chi-Square Test of Independence** (Check if two categorical variables are related or independent)
3. **Chi-Square Test of Homogeneity** (Check if multiple groups have same distribution)

Chi-Square Test Statistic Formula

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

χ^2	Chi-square test statistic
O	Observed frequency (actual data)
E	Expected frequency (what we expect if H_0 is true)
Σ	Sum across all categories

Chi-Square Distribution (Right-Skewed)



The chi-square distribution is always right-skewed and takes only positive values ($\chi^2 \geq 0$).

Real-World Examples

Marketing
- Check if customer preferences (e.g., product categories) are different from expected distribution.

Are product preferences as expected?

Operations
- Check if the proportion of defective products differs from the target across multiple production lines.

Is defect rate consistent across lines?

HR Analytics
- Check if gender distribution is the same across departments.

Is gender distribution independent of department?

Healthcare
- Check if treatment outcomes (e.g., recovery, no change, worsening) differ between two groups.

Are treatment outcomes different across groups?

Key Takeaways

- Chi-Square test is used for categorical data.
- Compares observed and expected frequencies.
- Used to check goodness of fit or independence.
- Always uses a right-skewed distribution ($\chi^2 \geq 0$).
- Widely used in business, healthcare, marketing, and more.

Common Mistakes to Avoid

- Using Chi-Square test for continuous data.
- Ignoring expected frequency conditions (should be ≥ 5).
- Misinterpreting significance as practical importance.
- Not checking assumptions.
- Incorrectly calculating expected frequencies.



2 Chi-Square Goodness of Fit Test

Checking If Observed Frequencies Match an Expected Distribution

What is Chi-Square Goodness of Fit Test?

The Chi-Square Goodness of Fit Test is used to check whether the observed frequencies in a single categorical variable match a specified expected distribution.

> It helps us determine if the data we collected follows a particular distribution (e.g., uniform, given percentages, or a known pattern).

When to Use?

- When you have one categorical variable.
- When you want to see if the observed frequencies match a known or expected distribution.
- Commonly used for product preferences, customer feedback, survey results, or quality control.
- Used in marketing, operations, retail, HR, and healthcare.

Hypotheses

H_0 The observed frequencies match the expected distribution. (data follows the given pattern)

H_1 The observed frequencies do not match the expected distribution. (data does not follow the given pattern)

Test Statistic Formula

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

χ^2	Chi-square test statistic
O	Observed frequency (from data)
E	Expected frequency (from hypothesis)
$\sum$	Sum across all categories
df	Degrees of freedom = $k - 1$ (where k = number of categories)

Example: Product Preference Survey

Problem:
A company wants to check if customer preferences for its 4 products are equally distributed (25% each). A survey of 200 customers gives the following results.

Product	Observed (O)	Expected (E)	$(O - E)^2$	$\frac{(O - E)^2}{E}$
A	50	50	0	0.00
B	70	50	400	8.00
C	45	50	25	0.50
D	35	50	225	4.50
Total	200	200	-	13.00

Chi-square statistic: $\chi^2 = 13.00$
Degrees of freedom: $df = 4 - 1 = 3$


Decision Using Critical Value

Significance level (α)	0.05
Degrees of freedom (df)	3
Critical value ($\chi^2_{critical}$)	7.815
Calculated χ^2 value	13.00

Decision:
Since 13.00 > 7.815, we reject H_0.

Interpretation:
The observed product preferences do not match an equal distribution. Some products are significantly more or less preferred than others.

Chi-Square Distribution (Right-Skewed)



The chi-square distribution is right-skewed and only takes positive values ($\chi^2 \geq 0$).

Excel Implementation

1. Enter observed frequencies in B2:B5
2. Enter expected frequencies in C2:C5
3. Calculate $(O - E)^2 / E$ in D2:D5
   `=(B2-C2)^2/C2`
4. Sum the values in D2:D5
   `=SUM(D2:D5)`
5. Find critical value ($\alpha = 0.05$, df = 3)
   `=CHISQ.INV.RT(0.05,3)` Result: 7.815

More Business Examples

Marketing
Check if customer preferences across product categories follow a specific distribution.

Operations
Check if defects are evenly distributed across different production lines.

HR Analytics
Check if employee distribution across departments matches the expected plan.

Healthcare
Check if patient outcomes follow a known distribution (e.g., equal rates).

Key Takeaways

- Used for one categorical variable.
- Compares observed and expected frequencies.
- Right-skewed distribution ($\chi^2 \geq 0$).
- Degrees of freedom = number of categories – 1.
- Reject H_0 if $\chi^2 > \chi^2$ critical.
- Helpful for marketing, operations, HR, and healthcare.

Common Mistakes to Avoid

- Using Chi-Square test for continuous data.
- Expected frequencies less than 5.
- Incorrect degrees of freedom.
- Not checking if data is categorical.
- Misinterpreting the result in business context.
- Forgetting to compare with critical value.

Quick Summary

- Tests if observed frequencies follow a specified distribution.
- Use χ^2 formula.
- Compare with critical value.
- If $\chi^2 > \chi^2$ critical → Reject H_0.
- Widely used in real business scenarios.



3 Chi-Square Test of Independence

Checking If Two Categorical Variables Are Related or Independent

What is Chi-Square Test of Independence?

The Chi-Square Test of Independence is used to check whether two categorical variables are related or independent.

> It helps us find out if there is a significant association between two categorical variables (e.g., gender and product preference).

When to Use?

- When you want to test if two categorical variables are related.
- When you have data in a contingency table (two or more categories for each variable).
- Commonly used in marketing, operations, HR, healthcare, and product analytics.
- Both variables should be categorical (nominal).

Hypotheses

H_0 The two categorical variables are independent. (No relationship between the variables.)


H_1 The two categorical variables are dependent. (There is a significant relationship between the variables.)

Test Statistic Formula

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

χ^2	Chi-square test statistic
O	Observed frequency (from data)
E	Expected frequency (under assumption of independence)
$\sum$	Sum across all cells
df	Degrees of freedom = $(r-1)(c-1)$ where r = number of rows, c = number of columns

Chi-Square Distribution (Right-Skewed)



The chi-square distribution is right-skewed and only takes positive values ($\chi^2 \geq 0$).

Example: Gender vs Product Preference

Problem:
A company wants to check whether gender and product preference are independent. A survey of 200 customers gives the following data.

Observed Frequencies (O)

	Product A	Product B	Product C	Total
Male	30	50	20	100
Female	20	30	50	100
Total	50	80	70	200

Calculation of χ^2

Cell	O	E	$(O-E)^2$	$(O-E)^2/E$
Male-A	30	25	25	1.00
Male-B	50	40	100	2.50
Male-C	20	35	225	6.43
Female-A	20	25	25	1.00
Female-B	30	40	100	2.50
Female-C	50	35	225	6.43
Total χ^2	-	-	-	19.86

Expected Frequencies (E)

$$E = \frac{(\text{Row Total} \times \text{Column Total})}{\text{Grand Total}}$$

	Product A	Product B	Product C
Male	25	40	35
Female	25	40	35

Degrees of Freedom

$$df = (r-1)(c-1) = (2-1)(3-1) = 1 \times 2 = 2$$

Critical Value and Decision

Significance level (α) = 0.05
degrees of freedom (df) = 2
Critical value ($\chi^2_{critical}$) = 5.991
Calculated χ^2 = 19.86

Since 19.86 > 5.991, we reject H_0.

There is a significant relationship between gender and product preference.

Excel Implementation

1. Enter observed data in B3:D4.
2. Calculate expected frequency for B3: `=(B5*B2)/$E$5`
3. Copy the formula for all cells.
4. Calculate $(O-E)^2 / E$ for each cell.
5. Sum all values to get χ^2 statistic.
6. Find critical value: `=CHISQ.INV.RT(0.05,2)` **Result: 5.991**

Key Takeaways

- Used to test the relationship between two categorical variables.
- Requires a contingency table.
- Compare observed and expected frequencies.
- Degrees of freedom = $(r-1)(c-1)$.
- Right-skewed distribution ($\chi^2 \geq 0$).
- Reject H_0 if $\chi^2 > \chi^2_{critical}$.
- Helps find associations, not causation.

Real-World Business Use Cases

Marketing
Check if product preference is related to gender, age group, or region.

Retail
Check if store location and purchase category are related.

HR Analytics
Check if department and attrition (Yes/No) are related.

Healthcare
Check if treatment type and recovery outcome are related.



4 Expected Frequency

How to Calculate and Interpret Expected Frequencies in Chi-Square Tests

What is Expected Frequency?

Expected frequency (E) is the number of observations we expect to see in each category if the null hypothesis is true (i.e., no difference or no relationship).

> It represents the theoretical distribution of data based on the assumption of independence or a specified distribution.

Why is it Important?

- Used to compare with observed frequency (O) in Chi-Square Test.
- Helps determine if the difference between observed and expected is statistically significant.
- Ensures a fair comparison across categories.
- If expected frequencies are very small (< 5), the Chi-Square test may not be valid.

Formula for Expected Frequency

$$E = \frac{(\text{Row Total} \times \text{Column Total})}{\text{Grand Total}}$$

E	Expected frequency
Row Total	Total of the row
Column Total	Total of the column
Grand Total	Total of all observations

Example: Customer Preference by Region (Marketing Use Case)

A company conducted a survey of 300 customers to study the preference for three product categories (A, B, C) across two regions (North and South). Test whether product preference is independent of the region.

Observed Frequency Table (O)

Region	Product A	Product B	Product C	Total
North	50	30	40	120
South	30	70	80	180
Total	80	100	120	300

Expected Frequency Calculation (E)

$$E = \frac{(\text{Row Total} \times \text{Column Total})}{\text{Grand Total}}$$


Example: Expected frequency for North - Product A

$$E = \frac{120 \times 80}{300} = 32$$

Expected Frequency Table (E)

Region	Product A	Product B	Product C	Total
North	32	40	48	120
South	48	60	72	180
Total	80	100	120	300

Chi-Square Distribution (Right-Skewed)



The chi-square distribution is right-skewed and only takes positive values ($\chi^2 \geq 0$).

Calculation of $(O - E)^2 / E$

Cell	O	E	O – E	$(O - E)^2$	$(O - E)^2/E$
N - A	50	32	18	324	10.13
N - B	30	40	−10	100	2.50
N - C	40	48	−8	64	1.33
S - A	30	48	−18	324	6.75
S - B	70	60	10	100	1.67
S - C	80	72	8	64	0.89
Total χ^2		-	-	-	23.27

Key Takeaways

- Expected frequency is based on the assumption of independence or a specified distribution.
- Calculated using row and column totals.
- All expected frequencies should ideally be ≥ 5.
- Used in Chi-Square goodness of fit and independence tests.
- Helps compare observed vs. expected counts.

Excel Implementation

1. Enter observed frequencies in a table.
2. Calculate row totals and column totals using SUM().
3. Use the formula = (Row Total * Column Total) / Grand Total.
4. Fill the formula for all cells.
5. Verify that all expected frequencies are ≥ 5.
6. Then calculate $(O - E)^2 / E$ and sum to get χ^2.

Common Mistakes to Avoid

- Incorrect calculation of row or column totals.
- Using observed instead of expected frequencies.
- Expected frequencies less than 5.
- Rounding errors in calculations.
- Not checking the assumption of independence.

Real-World Business Applications

Marketing
Product preference across regions, age groups, or channels

Operations
Defect types across production lines

HR Analytics
Employee satisfaction across departments

Healthcare
Treatment outcomes across patient groups



5 Assumptions of the Chi-Square Test

Conditions for Valid Results in Chi-Square Tests

Why Assumptions are Important?

The Chi-Square test is based on certain assumptions. If these assumptions are not met, the test results may not be valid and can lead to incorrect conclusions.

> Always check the assumptions before applying the Chi-Square test to ensure the results are reliable and meaningful.

Key Assumptions

1. Data must be categorical (nominal variables).
2. Observations must be independent.
3. Sample should be randomly selected.
4. Expected frequency in each cell should generally be ≥ 5.
5. Categories should be mutually exclusive and collectively exhaustive.
6. Data should be in the form of frequency counts (not percentages).

What Do These Mean?

1. Variables should be categories (e.g., gender, product type, Yes/No).
2. Each observation should belong to only one category and should not influence another.
3. Data should represent the population fairly (no selection bias).
5. Most expected frequencies should be 5 or more (rule of thumb).
5. All possible categories should be included and should not overlap.
6. Use actual counts of observations, not percentages or proportions.

Example 1: Independence Test (Valid Assumptions)

A company surveys 200 customers to check if gender and product preference are related.

Gender	Product Preference			Total
	A	B	C	
Male	30	50	20	100
Female	20	30	50	100
Total	50	80	70	200

Assumptions Check:

- Data is categorical → Yes
- Randomly selected customers → Yes
- Observations are independent → Yes
- All expected frequencies ≥ 5 → Yes
- Categories are mutually exclusive → Yes
- Counts are used (not percentages) → Yes

Example 2: Goodness of Fit Test

A dice is rolled 60 times. Test if it follows a uniform distribution (each number 1/6).

Face	Observed (O)	Expected (E)
1	8	10
2	12	10
3	9	10
4	11	10
5	10	10
6	10	10
Total	60	60

Assumptions Check:

- Data is categorical → Yes
- Random sample → Yes
- Independent rolls → Yes
- Expected frequencies = 10 (≥ 5) → Yes

Example 3: Violation of Assumption

A small survey of 12 customers to check brand preference (3 brands).

Brand	Observed (O)	Expected (E)
A	8	4
B	2	4
C	2	4
Total	12	12

Assumptions Check:


- Sample size is very small (n = 12).
- Expected frequency for Brand B and C is 4 (< 5) → Violates assumption.
- Chi-Square test may not be valid.

In such cases, use Fisher's Exact Test (instead of Chi-Square Test).

Common Issues and Solutions

Issue	Explanation	Solution
Expected frequency < 5	Small sample size or rare categories.	Combine categories or collect more data. Use Fisher's Exact Test if needed.
Non-categorical data	Data is continuous (e.g., age, salary).	Convert into categories or use other tests (e.g., t-test, ANOVA).
Dependent observations	Observations are related (e.g., same person counted twice).	Ensure independent sampling.
Missing categories	Not all possible categories are included.	Include all categories that can occur.
Using percentages	Using % instead of counts for calculations.	Always use frequency counts.

Visual Understanding



If expected frequencies are very small (E < 5), the Chi-Square approximation may not be accurate, leading to incorrect conclusions.

Key Takeaways

- Check all assumptions before applying the Chi-Square test.
- Use categorical and independent data.
- Expected frequencies should be ≥ 5 for most cells.
- Use Fisher's Exact Test for small sample sizes.
- Ensure categories are mutually exclusive and collectively exhaustive.
- Violating assumptions can lead to misleading results.

Interview Questions

1. What are the assumptions of the Chi-Square test?
3. Why is an expected frequency of 5 used?
3. What happens if expected frequency is less than 5?
4. When would you use Fisher's Exact Test?



6 Steps in a Chi-Square Test

A Step-by-Step Approach with Complete Example

Overview

The Chi-Square test follows a structured process. By following these steps, we can determine whether the observed frequencies differ significantly from the expected frequencies.

> A systematic approach helps in accurate calculation, better interpretation, and correct business conclusions.

Steps in a Chi-Square Test

1. Define the problem and state hypotheses.
2. Create a contingency table with observed frequencies (O).
3. Calculate the expected frequencies (E).
4. Compute the chi-square test statisitc (χ^2).
5. Determine the degrees of freedom (df).
6. Find the critical value from chi-square table.
7. Make a decision (Reject or Fail to Reject H_0).
8. Interpret the result in business context.

Key Questions at Each Step

1. What is the business problem?
2. What are H_0 and H_1?
3. Are my observed counts correct?
4. How do I calculate expected counts?
5. What is the degrees of freedom?
6. What is the critical value at my significance level?
7. Is $\chi^2 > \chi^2$ critical?
8. What does the result mean for the business?

Step-by-Step Example (Chi-Square Test of Independence)

A company wants to check if there is a relationship between customer age group and product preference. A survey of 300 customers gives the following data:

Observed Frequencies (O)

Age Group	Product A	Product B	Product C	Total
18–25	40	50	30	120
26–40	30	60	30	120
41+	20	40	20	80
Total	90	150	80	300

Step 1: State Hypotheses

H_0: Age group and product preference are independent (no relationship).

H_1: Age group and product preference are dependent (there is a relationship).

Step 2: Calculate Expected Frequencies (E)

$$E = \frac{(\text{Row Total} \times \text{Column Total})}{\text{Grand Total}}$$

Example: Expected frequency for age group 18–25, Product A

$$E = \frac{120 \times 90}{300} = 36$$

Expected Frequency Table (E)

Age Group	Product A	Product B	Product C	Total
18–25	36	60	24	120
26–40	36	60	24	120
41+	24	40	16	80
Total	90	150	80	300

Step 3: Calculate χ^2 Statistic

Cell	O	E	$(O-E)^2$	$(O-E)^2/E$
18–25, A	40	36	16	0.44
18–25, B	50	60	100	1.67
41–25, C	30	24	36	1.50
... (other cells)		24	16	1.80
Total χ^2	-	-		13.56

Step 4: Degrees of Freedom

$$df = (r-1)(c-1)$$

r = number of rows = 3

c = number of columns = 3

$$df = (3-1)(3-1) = 2 \times 2 = 4$$

Step 5: Find Critical Value

Significance level (α) = 0.05

Degrees of freedom (df) = 4

From chi-square table:

χ^2 critical = **9.488**


Step 5: Make a Decision

Since 13.56 > 9.488, we reject H_0.

Step 7: Interpretation

There is a significant relationship between age group and product preference.

Chi-Square Distribution (Right-Skewed)



The chi-square distribution is right-skewed and only takes positive values ($\chi^2 \geq 0$).

Excel Implementation

1. Enter observed frequencies in B3:D5.
2. Calculate row totals and column totals using SUMI).
3. Calculate expected frequencies using:
   = (Row Total × Column Total) / Grand Total
4. Compute $(O-E)^2 / E$ for each cell.
5. Sum all values to get the chi-square statistic.
6. Find critical value using:
   =CHISQ.INV.RT(0.05,4)

Result: 9.488

Key Takeaways

- Follow a systematic 8-step process.
- State clear hypotheses.
- Calculate expected frequencies correctly.
- Use the correct degrees of freedom.
- Compare χ^2 with critical value.
- Interpret the result in business terms.
- Widely used in marketing, operations, HR, and healthcare.

Common Mistakes to Avoid

- Incorrect row or column totals.
- Expected frequency < 5.
- Using percentages instead of counts.
- Wrong degrees of freedom.
- Not checking assumptions.
- Misinterpreting the result.
- Using wrong significance level.
- Calculation errors in Excel.

Real-World Business Use Cases

Marketing
Customer preferences across age groups, regions or channels.

Retail
Check if product categories are associated with store locations.

HR Analytics
Employee satisfaction across departments.

Healthcare
Treatment type is related to recovery outcome.



7 Excel Implementation of the Chi-Square Test

Step-by-Step Guide with Formulas, Functions and Practical Example

Why Use Excel?

Excel makes it easy to perform Chi-Square tests using built-in functions. We can quickly calculate expected frequencies, test statistic, critical value and p-value, and draw conclusions.

> Excel is widely used in business, making it a practical skill for data analysts, business analysts, and MIS professionals.

Key Excel Functions

Function	Purpose
SUM()	Calculate row/column totals
CHISQ.TEST()	Returns p-value for the test
CHISQ.INV.RT()	Critical value (right tail)
CHISQ.DIST.RT()	p-value from χ^2 statistic
SUMPRODUCT()	Calculate χ^2 statistic manually
IF()	Make decision based on result

Steps in Excel

1. Enter observed frequencies in a table.
2. Calculate row, column and grand totals.
3. Calculate expected frequencies (E).
4. Compute $(O - E)^2 / E$ for each cell.
5. Sum all values to get χ^2 statistic.
6. Find critical value using CHISQ.INV.RT().
7. Find p-value using CHISQ.TEST() or CHISQ.DIST.RT().
8. Compare and interpret the result.

Example: Marketing Campaign (Channel vs Purchase)

A company surveys 200 customers to check if purchase decision is independent of marketing channel (Online, TV, In-Store).

Observed Frequency Table (O)

A	Online	TV	In-Store	Total
Purchased	50	30	40	120
Not Purchased	30	20	30	80
Total	80	50	70	200

Step 1: Calculate Expected Frequencies (E)

A	B	C	D	E
Category	Online	TV	In-Store	Total
Purchased	48	30	42	120
Not Purchased	32	20	28	80
Total	80	50	70	200

Formula in Excel:
= (Row Total × Column Total) / Grand Total

Example for cell B2:
=($E2*B$4)/$E$4
= (120*80)/200 = 48

Step 2: Calculate $(O - E)^2 / E$ and χ^2 Statistic

A	B	C	D	E
Category	Online	TV	In-Store	Total
Purchased	0.083	0.000	0.095	0.179
Not Purchased	0.125	0.000	0.143	0.268
Total χ^2	0.208	0.000	0.238	0.446

Excel Formula for each cell: =(B2-C2)^2/C2

Chi-square statistic: =SUM(B2:D3)

Result: 0.446

Step 3: Critical Value and p-Value

Degrees of freedom:

$$df = (r - 1)(c - 1)$$
$$= (2 - 1)(3 - 1) = 2$$

Significance level (α)	0.05
Degrees of freedom (df)	2
Critical value ($\chi^2_{crittical}$)	5.991

Excel formula for critical value: =CHISQ.INV.RT(0.05,2)

Result: 5.991

Step 4: p-Value Using Excel


Excel Function:
=CHISQ.TEST(B2:D3,B7:D8)

Result (p-value): 0.800

Step 5: Decision and Interpretation

- Since p-value (0.800) > 0.05, we fail to reject H_0.
- There is no significant relationship between marketing channel and purchase decision.

Chi-Square Distribution in Excel



The chi-square distribution is right-skewed and only takes positive values ($\chi^2 \geq 0$).

Alternative: Get p-Value from χ^2 Statistic

If you already have χ^2 statistic (e.g., 0.446):

=CHISQ.DIST.RT(0.446,2)

Result: 0.800

Quick Excel Workflow (Formulas Summary)

Row totals:	=SUM(B2:D2)
Column totals:	=SUM(B2:B3)
Grand total:	=SUM(B4:D4)
Expected frequency:	=($E2*B$4)/$E$4
Chi-square statistic:	=SUMPRODUCT((B2:D3-C2:D3)^2/C2:D3)
p-value:	=CHISQ.TEST(B2:D3,B7:D8)
Critical value:	=CHISQ.INV.RT(0.05,2)

Real-World Business Applications

Marketing
Check if purchase behavior is related to channel, region, or customer segment.

Retail
Test if product category is independent of store location.

HR Analytics
Check if employee department is related to attrition (Yes/No).

Healthcare
Test if treatment type is independent of recovery outcome.

Key Takeaways

- Excel can quickly perform Chi-Square tests.
- CHISQ.TEST() gives p-value directly.
- CHISQ.INV.RT() gives critical value.
- Always check assumptions before using Excel.
- Interpret the result in business context.



8 Real-Life Business Applications

Use Cases, Insights and Interview Questions with Answers

1 Marketing Analytics

Are customer preferences different across regions?

Region	A	B	C	Total
North	50	30	20	100
South	30	40	30	100
East	20	30	50	100
Total	100	100	100	300

Business Insight

If the test is significant, product preference varies by region → use region-specific marketing strategies.

2 Retail & E-commerce

Is product category independent of store type?

Store Type	Electronics	Fashion	Home	Total
Online	60	50	40	150
Offline	40	70	40	150
Total	100	120	80	300

Business Insight

A significant result means product category and store type are related → adjust inventory and promotions accordingly.

3 HR Analytics

Is employee satisfaction related to department?

Department	Satisfied	Neutral	Not Satisfied
Onles	40	30	30
IT	50	30	20
HR	30	40	30
Total	120	100	80

Business Insight

A significant result shows satisfaction levels differ by department → take targeted actions to improve engagement.

4 Healthcare

Is treatment type related to recovery outcome?

Treatment	Recovered	Not Recovered	Total
Type A	70	30	100
Type B	50	50	100
Type C	40	60	100
Total	160	140	300

Business Insight

A significant result indicates recovery rate depends on treatment type → helps in decision-making and resource allocation.

5 Customer Behavior

Is purchase decision related to customer segment?

Segment	Purchased	Not Purchased	Total
Students	40	60	100
Working	70	30	100
Retired	20	80	100
Total	130	170	300

Business Insight

If significant, purchase behavior varies by customer segment → create targeted offers and campaigns.

6 Operations & Quality

Is defect type related to production line?

Production Line	Type A	Type B	Type C	Total
Line 1	30	40	30	100
Line 2	20	50	30	100
Line 3	50	20	30	100
Total	100	110	90	300

Business Insight

A significant result shows defect types vary across production lines → focus on quality control for specific lines.

7 Finance & Risk

Is loan default related to customer category?

Customer Type	Default	No Default	Total
Salaried	20	80	100
Self-Employed	40	60	100
Business	50	50	100
Total	110	190	300

Business Insight

A significant result indicates default risk varies by customer type → helps in credit risk assessment and policy making.

8 Education / Academia

Is course preference related to gender?

Gender	Analytics	Finance	Marketing	Total
Male	50	30	20	100
Female	30	40	30	100
Total	80	70	50	200

Business Insight

A significant result shows course preference depends on gender → helps in designing targeted programs and resources.

Key Takeaways

1. Chi-Square tests are used for categorical data.
2. We compare observed vs. expected frequencies.
3. Expected frequency should generally be ≥ 5.
4. Choose the correct test (Goodness of Fit or Independence).
5. Use Excel functions for quick and accurate calculations.
6. Results help in business decision-making, showing associations, not causation.

Common Interview Questions & Answers

Q1. What is a Chi-Square test?
A: It is a non-parametric test used to analyze categorical data to determine if there is a significant difference or relationship between variables.

Q2. What are the types of Chi-Square tests?
A: Chi-Square Goodness of Fit Test and Chi-Square Test of Independence.

Q3. What is the null hypothesis in Chi-Square test?
A: For Goodness of Fit: Observed distribution matches expected distribution.
For Independence: Two variables are independent.

Q4. What is the expected frequency?
A: It is the number of observations we expect in a category if the null hypothesis is true, calculated as (Row Total × Column Total) / Grand Total.

Q5. What is the minimum expected frequency required?
A: Generally, each expected frequency should be ≥ 5.

Q6. What does a significant Chi-Square result mean?
A: It means there is a significant difference (Goodness of Fit) or a significant relationship (Independence) between the categorical variables.

Q7. How do you calculate degrees of freedom?
A: For Goodness of Fit: $df = k - 1$
For Independence: $df = (r - 1)(c - 1)$

Q8. Which Excel functions are used for Chi-Square test?
A: CHISQ.TEST(), CHISQ.INV.RT(), CHISQ.DIST.RT()

Q9. When do you use Chi-Square Goodness of Fit Test?
A: When you want to check if the observed frequency matches a specified expected distribution.

Q10. When do you use Chi-Square Test of Independence?
A: When you want to check if there is a relationship between two categorical variables.

Q11. What are the assumptions of Chi-Square test?
A: Data should be categorical, observations independent, expected frequency ≥ 5, and categories mutually exclusive and collectively exhaustive.

Q12. Does Chi-Square test tell us causation?
A: No, it only tells us if there is an association or relationship, not causation.