Centre No.					Pape	er Refer	ence			Surname	Initial(s)
Candidate No.			6	6	6	8	/	0	1	Signature	

Paper Reference(s)

6668/01

Edexcel GCE

Further Pure Mathematics FP2 Advanced/Advanced Subsidiary

Thursday 23 June 2011 – Morning

Time: 1 hour 30 minutes

Materials required for examination	Items included with question paper
Mathematical Formulae (Pink)	Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance wit Edexcel Limited copyright policy. ©2011 Edexcel Limited.

Printer's Log. No. P35413A



Examiner's use only

Team Leader's use only

Turn over

Total



	$\frac{3}{x+3} > \frac{x-4}{x}$	(7)
		()

2.

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} = \mathrm{e}^x \left(2y \frac{\mathrm{d}y}{\mathrm{d}x} + y^2 + 1 \right)$$

(a) Show that

$$\frac{d^{3}y}{dx^{3}} = e^{x} \left[2y \frac{d^{2}y}{dx^{2}} + 2\left(\frac{dy}{dx}\right)^{2} + ky \frac{dy}{dx} + y^{2} + 1 \right],$$

where k is a constant to be found.

(3)

Given that, at x = 0, y = 1 and $\frac{dy}{dx} = 2$,

(b) find a series solution for y in ascending powers of x, up to and including the term in x^3 .

(4)

3.	Find the	general	solution	of the	differential	equation
-----------	----------	---------	----------	--------	--------------	----------

$$x\frac{\mathrm{d}y}{\mathrm{d}x} + 5y = \frac{\ln x}{x}, \qquad x > 0$$

giving your answer in the form $y = f(x)$.	
	(8)

4. Given that

$$(2r+1)^3 = Ar^3 + Br^2 + Cr + 1,$$

(a) find the values of the constants A, B and C.

(2)

(2)

(b) Show that

$$(2r+1)^3 - (2r-1)^3 = 24r^2 + 2$$

(c) Using the result in part (b) and the method of differences, show that

$$\sum_{r=1}^{n} r^2 = \frac{1}{6} n(n+1)(2n+1) \tag{5}$$

Question 4 continued	blank
Question 4 continued	



5. The point P represents the complex number z on an Argand diagram, where

$$|z - \mathbf{i}| = 2$$

The locus of P as z varies is the curve C.

(a) Find a cartesian equation of *C*.

(2)

(b) Sketch the curve *C*.

(2)

A transformation T from the z-plane to the w-plane is given by

$$w = \frac{z+i}{3+iz}, \quad z \neq 3i$$

The point Q is mapped by T onto the point R. Given that R lies on the real axis,


(c) show that Q lies on C.

(5)

stion 5 continued		



6.



The curve C shown in Figure 1 has polar equation

$$r = 2 + \cos \theta$$
, $0 \leqslant \theta \leqslant \frac{\pi}{2}$

At the point A on C, the value of r is $\frac{5}{2}$.

The point N lies on the initial line and AN is perpendicular to the initial line.

The finite region R, shown shaded in Figure 1, is bounded by the curve C, the initial line and the line AN.

Find the exact area of the shaded region R.

	1	•	١,	
- (ľ		,	١
- 1	١.	,	•	ı

Question 6 continued	blank
Question o constitue	

_eave	
1001-	
hlank	

7.	(a)	Use	de	Moivre'	S	theorem	to	show	that
/ •	(a)	USC	uc	IVIOIVIC	S	uicorciii	w	SHOW	mai

$$\sin 5\theta = 16\sin^5\theta - 20\sin^3\theta + 5\sin\theta$$

(5)

Hence, given also that $\sin 3\theta = 3\sin\theta - 4\sin^3\theta$,

(b) find all the solutions of

$$\sin 5\theta = 5\sin 3\theta,$$

in the interval $0 \le \theta < 2\pi$. Give your answers to 3 decimal places.

(6)

Question 7 continued	blanl

8. The differential equation

$$\frac{\mathrm{d}^2 x}{\mathrm{d}t^2} + 6\frac{\mathrm{d}x}{\mathrm{d}t} + 9x = \cos 3t, \quad t \geqslant 0$$

describes the motion of a particle along the *x*-axis.

(a) Find the general solution of this differential equation.

(8)

(b) Find the particular solution of this differential equation for which, at t = 0,

$$x = \frac{1}{2} \text{ and } \frac{\mathrm{d}x}{\mathrm{d}t} = 0.$$
 (5)

On the graph of the particular solution defined in part (b), the first turning point for t > 30 is the point A.

(c) Find approximate values for the coordinates of A.

(2)

		Leave blank
Question 8 continued		
		Q8
	(Total 15 marks)	
Т	OTAL FOR PAPER: 75 MARKS	
END		