

Mark Scheme (Results)

Summer 2013 - Home

GCE Mechanics M2 6678/01 Original Paper

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2013
Publications Code N/A
All the material in this publication is copyright
© Pearson Education Ltd 2013

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

1306



June 2013 M2 6678/01 - breach paper Mark Scheme

Question Number	Scheme	Marks	Notes
1. (a)	$2\binom{a}{3} + 3\binom{3}{-1} + m\binom{-2}{4} = (m+5)\binom{0}{2}$		
	$6-3+4m=(m+5)\times 2$	M1	Moments equation for <i>y</i> coordinate. Terms of correct structure but condone sign slips
		A2	-1 each error
	(7=2m), m=3.5	A1 (4)	
(b)	$2a+9-2m=0\big(\times(m+5)\big)$	M1	Moments equation for <i>x</i> coordinate. Terms of correct structure but condone sign slips
		A1	
	2a+9-7=0	M1	Substitute their <i>m</i> and solve for <i>a</i> .
	use their m		
	a = -1	A1 (4)	
		[8]	
2. (a)	Use of $\frac{P}{V} = F$	M1	
	Use of $\frac{P}{V} = F$ $\frac{32000}{V} = 800$	A1	
	V = 40	A1 (3)	
(b)	$F = \frac{32000}{20} = 1600$	M1	Find the driving force
	$1600 - 800 + 1200g\sin\theta = 1200a$	M1	Equation of motion for their <i>F</i> . Requires all 4 terms. Condone sign errors and trig confusion
	$800 + \frac{1200g}{40} = 1200a$	A1	Correct equation (allow with $\sin \theta$)
	$a = 0.91 \mathrm{(m s^{-2})},$	A1	Accept 0.912
	\ //	(4)	*

Question Number	Scheme	Marks	Notes
		[7]	
3. (a)	Momentum = $0.25 \times ((2-12) \mathbf{i} + (9+6) \mathbf{j})$	M1	Substitution of $m = 0.25$ and $t = 3$
	$= -2.5\mathbf{i} + 3.75\mathbf{j}$	A1	or simplified equivalent $-\frac{5}{2}\mathbf{i} + \frac{15}{4}\mathbf{j}$
		(2)	
(b)	$\mathbf{a} = -4\mathbf{i} + (2\mathbf{t} + 2)\mathbf{j}$	M1	Differentiate v
(-)	79	A2	A1 for each term
	Use of $\mathbf{F} = m\mathbf{a}$	M1	
	Use of Pythagoras on F or a	M1	
	$ F = 0.25 \times \sqrt{16 + (6+2)^2} = \sqrt{5}$	A1	2.24 or better
		(6)	
(c)	$\mathbf{s} = \left(2t - 2t^2\right)\mathbf{i} + \left(\frac{t^3}{3} + t^2\right)\mathbf{j} \ (+ \mathbf{r_0})$	M1	Integrate to find displacement
		A2	A1 for each component. \mathbf{r}_0 not required.
	Use of $\mathbf{r_0}$ and $t = 3$	M1	
	Position vector of A is $-10\mathbf{i} + 14\mathbf{j}$	A1	
		(5)	
		[13]	

Question Number	Scheme	Marks	Notes
4. (a)	Energy at A = Energy at B	M1	Equation requires all 3 terms. Condone sign errors
	$\frac{1}{2}m \times 15^2 = \frac{1}{2}m \times 12^2 + mgh$	A2	-1 each error
	$h = \frac{27 \times 3}{2g} \approx 4.1$	A1	Allow 4.13 $\frac{405}{98} \text{ is A0}$
		(4)	
(b)	Use of $s = ut + \frac{1}{2}gt^2$	M1	or equivalent complete method to an equation in α
	$4 = 12\sin\alpha \times 1.5 - \frac{g}{2} \times 1.5^{2}$	A1	Correct unsimplied equation
	$(\sin \alpha = 0.834) \qquad \alpha = 56.6^{\circ}$	A1	Or 57°
		(3)	
(c)	$\cos\theta = \frac{12\cos\alpha}{15}$	M1 A1ft	Use of horizontal component and speed to find direction, or equivalent. follow their α 6.606, 13.47, 15
	$\theta = 64^{\circ}$	All Al	63.9 from calculator values, 64.2 from 57
	<i>U</i> – 0 1	(3)	03.7 Hom calculator values, 04.2 Hom 37
		[10]	
		[10]	

Question Number	Scheme	Marks	Notes
5. (a)	Moments about <i>OE</i> : $\frac{a}{2} \times 1 + (a+2) \times 1 = 2\overline{x}$	M1	Moments equation. Needs to include both rectangles and the whole shape.
	$2\bar{x} = \frac{3a}{2} + 2$, $\bar{x} = \frac{3a}{4} + 1$	A2 A1	-1 each error. or equivalent Or simplified equivalent.
	2 4	(4)	
(b)	Moments about <i>OA</i> : $2 \times 1 + \frac{a}{2} \times 1 = 2\overline{y}$ o.e.	M1	Moments equation. Needs to include both rectangles and the whole shape.
	$\overline{y} = 1 + \frac{a}{4}$ o.e.	A2 A1	-1 each error. or equivalent Or simplified equivalent.
		(4)	
(c)	Hanging in Equilibrium: $\tan \alpha = \frac{\overline{x}}{\overline{y}} = \frac{\frac{3a}{4} + 1}{1 + \frac{a}{4}}$	M1	Condone reciprocal
	$\frac{1+-}{4}$	A1ft	For their \bar{x} , \bar{y}
	$\frac{4}{3} = \frac{\frac{3a}{4} + 1}{1 + \frac{a}{4}} = \frac{3a + 4}{4 + a}, 4(4 + a) = 3(3a + 4)$	DM1	Their $\tan \alpha = \frac{4}{3}$ and solve for a
	16 + 4a = 9a + 12, $4 = 5a$,		
	$16 + 4a = 9a + 12, \ 4 = 5a,$ $a = \frac{4}{5}$	A1	Or 0.8
		(4)	
		[12]	

Question Number	Scheme	Marks	Notes
6. (a)	N B R d m P 45 cm		
	Moments about A: $30 \times 1.5 \cos 15 = d \times N$	M1	Requires correct terms of correct structure. Condone with value of <i>d</i> not yet found.
	$30 \times 1.5 \cos 15 = \frac{0.45}{\sin 15} N$	A2	-1 each error. Requires a value for d . Accept unsimplified. $d = 1.738$.
	$N = 100\cos 15\sin 15 = 25 \text{ (N)}$ *AG*	A1	Watch out for given answer
		(4)	
(b)	Moments about P: $R \times d \cos 15 = 30 \times (d-1.5) \cos 15$	M1	Terms of correct structure. Condone sign errors & trig confusion.
		A2ft	their d. Correct unsimplified1 each error
	$R = \frac{30(d-1.5)}{d} = 4.118095$	A1	4.1 or better
		(4)	
(c)	Resolving horizontally at P : $N \cos 75 = F \cos 15$	M1 A1	Condone trig confusion
	$F = \frac{N\cos 75}{\cos 15}$		
	$\mu \ge \frac{F}{N}$: $\mu \ge \frac{\cos 75}{\cos 15}$	M1	Use of $F \leq \mu N$
	$\mu \ge 0.268$	A1	0.27 or better
		(4)	
		[12]	

Question Number	Scheme	Marks	Notes
7. (a)	$\stackrel{2u}{\longrightarrow}$ $\stackrel{3u}{\longrightarrow}$		
	Q $2m$ $3u$		
	$\stackrel{2u}{\longleftarrow}$		
	$v \longrightarrow w$		
	Q collides with wall, $y' = \frac{1}{y^2y} = y$ towards R	M1	Correct use of impact law
	Q collides with wall: $v' = \frac{1}{3} \times 3u = u$ towards P	A1	
	P and Q collide.	M1	Requires all 4 terms. Condone sign errors
	Conservation of momentum:		
	$m \times 2u - 2m \times u = 2m \times w - m \times v$	A1	XXI A X
	$0 = 2m \times w - m \times v, 2w = v *AG*$	A1 (=	Watch out
(1)	Y	(5)	
(b)	Impact law: $v + w = e(2u + u)$	M1	Correct use of impact law to find w
	3w = 3eu, $w = eu$	A1	
	Kinetic energy:	M1	Requires all terms. Each term of correct structure
	$\frac{1}{2} \left[\frac{1}{2} m.4u^2 + \frac{1}{2} 2m.u^2 \right] = \frac{1}{2} m.v^2 + \frac{1}{2} 2m.w^2$	A2	-1 each error
	$\frac{1}{2}3u^2 = \frac{1}{2}4w^2 + \frac{1}{2}2w^2 = 3w^2$	M1	Solve for <i>w</i>
	$w = \frac{1}{\sqrt{2}}u$	A1	
	$e = \frac{1}{\sqrt{2}}$	A1	0.71 or better
	·	(8)	
		[13]	

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email <u>publication.orders@edexcel.com</u>
Order Code N/A Summer 2013

For more information on Edexcel qualifications, please visit our website $\underline{www.edexcel.com}$

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE





