Centre No.					Pape	r Refer	ence			Surname	Initial(s)
Candidate No.			6	6	7	9	/	0	1	Signature	

Paper Reference(s)

6679/01

Edexcel GCE

Mechanics M3

Advanced/Advanced Subsidiary

Friday 6 June 2008 – Afternoon

Time: 1 hour 30 minutes

Materials required for examination
Mathematical Formulae (Green)Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions. Write your answers in the spaces provided in this question paper.

If you need more space to complete your answer to any question, use additional answer sheets.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 6 questions in this question paper. The total mark for this question paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2008 Edexcel Limited.

 $\stackrel{\text{Printer's Log. No.}}{N29501}A$

W850/R6679/57570 3/3/3/2/



Turn over

Total



Examiner's use only

Team Leader's use only

Ouestion

1

Leave

3 4 5

Leave

1.




Figure 1

A light elastic spring, of natural length L and modulus of elasticity λ , has a particle P of mass m attached to one end. The other end of the spring is fixed to a point O on the closed end of a fixed smooth hollow tube of length L.

The tube is placed horizontally and P is held inside the tube with $OP = \frac{1}{2}L$, as shown in Figure 1. The particle P is released and passes through the open end of the tube with speed $\sqrt{(2gL)}$.

(a) Show that $\lambda = 8mg$. (4)

The tube is now fixed vertically and P is held inside the tube with $OP = \frac{1}{2}L$ and P above O. The particle P is released and passes through the open top of the tube with speed u.

(5)

(b) Find *u*.

		Leave
2.	A particle P moves with simple harmonic motion and comes to rest at two points A and B which are 0.24 m apart on a horizontal line. The time for P to travel from A to B is 1.5 s. The midpoint of AB is O . At time $t = 0$, P is moving through O , towards A , with speed u m s ⁻¹ .	blank
	(a) Find the value of u.	
	(4)	
	(b) Find the distance of P from B when $t = 2$ s. (5)	
	(c) Find the speed of P when $t = 2$ s. (2)	
_		
_		

Question 2 continued	blank
	_
	_
	_
	-
	-
	-
	-
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	-
	-
	-
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
	1



bla

Leave blank




Figure 2

Figure 2 shows a particle B, of mass m, attached to one end of a light elastic string. The other end of the string is attached to a fixed point A, at a distance h vertically above a smooth horizontal table. The particle moves on the table in a horizontal circle with centre O, where O is vertically below A. The string makes a constant angle θ with the downward vertical and B moves with constant angular speed ω about OA.

(a) Show that
$$\omega^2 \leqslant \frac{g}{h}$$
. (8)

The elastic string has natural length h and modulus of elasticity 2 mg.

Given that $\tan \theta = \frac{3}{4}$,

3.

(b) find ω in terms of g and h.

(5)

Question 3 continued		blank



Leave blank

4.




Figure 3

A uniform solid hemisphere, of radius 6a and centre O, has a solid hemisphere of radius 2a, and centre O, removed to form a bowl B as shown in Figure 3.

(a) Show that the centre of mass of B is $\frac{30}{13}a$ from O. (5)




Figure 4

The bowl B is fixed to a plane face of a uniform solid cylinder made from the same material as B. The cylinder has radius 2a and height 6a and the combined solid S has an axis of symmetry which passes through O, as shown in Figure 4.

(b) Show that the centre of mass of S is
$$\frac{201}{61}a$$
 from O. (4)

The plane surface of the cylindrical base of S is placed on a rough plane inclined at 12° to the horizontal. The plane is sufficiently rough to prevent slipping.

(c) Determine whether or not S will topple.

(4)

	Leave
Question 4 continued	blank
Question 4 continued	



Question 4 continued	bla

Leave

- 5. A particle *P* of mass *m* is attached to one end of a light inextensible string of length *a*. The other end of the string is attached to a fixed point *O*. The particle is released from rest with the string taut and *OP* horizontal.
 - (a) Find the tension in the string when *OP* makes an angle of 60° with the downward vertical.

(6)

A particle Q of mass 3m is at rest at a distance a vertically below O. When P strikes Q the particles join together and the combined particle of mass 4m starts to move in a vertical circle with initial speed u.

(b) Show that
$$u = \sqrt{\left(\frac{ga}{8}\right)}$$
.

The combined particle comes to instantaneous rest at A.

- (c) Find
 - (i) the angle that the string makes with the downward vertical when the combined particle is at A,
 - (ii) the tension in the string when the combined particle is at A.

(6)



Question 5 continued	Lea bla
	_
	_
	_
	_
	_
	_
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	-
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_



Leave blank

6. A particle P of mass 0.5 kg moves along the positive x-axis. It moves away from the origin O under the action of a single force directed away from O. When OP = x metres,

the magnitude of the force is $\frac{3}{(x+1)^3}$ N and the speed of P is v m s⁻¹.

Initially *P* is at rest at *O*.

(a) Show that $v^2 = 6\left(1 - \frac{1}{(x+1)^2}\right)$. (6)

(b) Show that the speed of P never reaches $\sqrt{6}$ m s⁻¹.

(1)

(c) Find x when P has been moving for 2 seconds.

(7)

	Leave blank
Question 6 continued	

