

6678/01

Edexcel GCE

Mechanics M2

Advanced/Advanced Subsidiary

Tuesday 9 June 2015 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Mathematical Formulae (Pink)

 $\overline{\text{Nil}}$

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either two significant figures or three significant figures.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 32 pages in this question paper. Any blank pages are indicated.

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2015 Pearson Education Ltd

Printer's Log. No.

Printer's Log. No.
P44837A

W850/R6678/57570 5/1/1/1/



Turn over

PEARSON

- At the instant when the van is moving down the road at 5 m s^{-1} , the acceleration of the van is $a \text{ m s}^{-2}$.

Find the value of a .

(5)



The diagram shows a square $ABCD$ with side length $2a$. The vertices are labeled D (top-left), A (top-right), C (bottom-left), and B (bottom-right). A point O is located inside the square, forming a V-shape with vertices D and A . The segments DO and AO are drawn, and the angle $\angle DOA$ is labeled as α . The distance from O to the bottom side CB is labeled as x .

The uniform lamina $OABCD$, shown in Figure 1, is formed by removing the triangle OAD from the square $ABCD$ with centre O . The square has sides of length $2a$.

- The mass of the lamina is M . A particle of mass kM is attached to the lamina at D to form the system S . The system S is freely suspended from A and hangs in equilibrium with AO vertical.

-
- This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper appears to be a standard notebook page or a sheet of stationery.

Leave
blank

Question 2 continued

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- Find

- (a) the value of θ ,
- (b) the kinetic energy gained by P as a result of the impulse.

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Leave
blank

Question 3 continued

Lined area for writing the answer to Question 3.



4. A ladder AB , of weight W and length $2l$, has one end A resting on rough horizontal ground. The other end B rests against a rough vertical wall. The coefficient of friction between the ladder and the wall is $\frac{1}{3}$. The coefficient of friction between the ladder and the ground is μ . Friction is limiting at both A and B . The ladder is at an angle θ to the ground, where $\tan \theta = \frac{5}{3}$. The ladder is modelled as a uniform rod which lies in a vertical plane perpendicular to the wall.

Find the value of μ .

(9)



Leave
blank

Question 4 continued



Diagram showing a cable AB of length 6.5 m suspended between two points A and B on a slope. The cable is divided into three equal segments of 2.17 m each by points C and D. The angle of the slope is α .

A particle P of mass 10 kg is projected from a point A up a line of greatest slope AB of a fixed rough plane. The plane is inclined at angle α to the horizontal, where $\tan \alpha = \frac{5}{12}$ and $AB = 6.5$ m, as shown in Figure 2. The coefficient of friction between P and the plane is μ . The work done against friction as P moves from A to B is 245 J.

- The particle is projected from A with speed 11.5 m s^{-1} . By using the work-energy principle,

-
- This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Leave
blank

Question 5 continued

[illegible]

- Find

- (b) the acceleration of P when $t = 5$ (3)

- (c) the total distance travelled by P in the interval $0 \leq t \leq 5$ (5)

Leave
blank

Question 6 continued

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Diagram illustrating the motion of a particle from point O to point C . The path is a semi-circle. The initial velocity at O is $u \text{ m s}^{-1}$ at an angle θ° to the horizontal. At point A , the velocity is 15 m s^{-1} at an angle of 20° to the horizontal. The horizontal distance from O to C is 10 m .

At time $t = 0$, a particle is projected from a fixed point O on horizontal ground with speed $u \text{ m s}^{-1}$ at an angle θ° to the horizontal. The particle moves freely under gravity and passes through the point A when $t = 4 \text{ s}$. As it passes through A , the particle is moving upwards at 20° to the horizontal with speed 15 m s^{-1} , as shown in Figure 3.

- (c) Find the distance OC . (3)

Leave
blank

Question 7 continued

This image shows a full page of blank, lined paper. It features approximately 28 horizontal grey lines spaced evenly apart, typical of standard notebook paper. The lines extend across the entire width of the page, leaving small margins at the top and bottom. There are no vertical lines, text, or other markings present.

Leave
blank

Question 8 continued

