

Mark Scheme (Results) January 2011

GCE

GCE Mechanics M2 (6678) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

January 2011

Publications Code UA026580

All the material in this publication is copyright

© Edexcel Ltd 2011

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M marks:** method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A marks:** Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B marks** are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations



These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod - benefit of doubt
- ft - follow through
- the symbol \checkmark will be used for correct ft
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw - ignore subsequent working
- awrt - answers which round to
- SC: special case
- oe - or equivalent (and appropriate)
- dep - dependent
- indep - independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- \square The second mark is dependent on gaining the first mark


January 2011
Mechanics M2 6678
Mark Scheme


Question Number	Scheme	Marks
1. (a)	Constant speed \Rightarrow Driving force = resistance , $F = 32$. $P = F \times v = 32v = 384$ $v = 12 \text{ (ms}^{-1}\text{)}$	B1 M1 A1 (3)
(b)	$P = F \times v \Rightarrow 384 = F \times 9, F = \frac{384}{9}$ Their $F - 32 = 120a$, $a = 0.089 \text{ (ms}^{-2}\text{)}$	M1 M1 A1 (3) [6]
2.	$\mathbf{I} = (-6\mathbf{i} + 8\mathbf{j}) = 2(\mathbf{v} - (5\mathbf{i} + \mathbf{j}))$ $-3\mathbf{i} + 4\mathbf{j} = \mathbf{v} - 5\mathbf{i} - \mathbf{j}$ $\mathbf{v} = 2\mathbf{i} + 5\mathbf{j}$ $\text{KE} = \frac{1}{2} \times 2 \times \mathbf{v} ^2 = (\sqrt{2^2 + 5^2})^2 = 29 \text{ (J)}$	M1A1 A1 M1 A1 (5)
3. (a)	$a = 4t^3 - 12t$ Convincing attempt to integrate $v = t^4 - 6t^2 (+c)$ Use initial condition to get $v = t^4 - 6t^2 + 8 \text{ (ms}^{-1}\text{)}$.	M1 A1 A1 (3)
(b)	Convincing attempt to integrate $s = \frac{t^5}{5} - 2t^3 + 8t (+0)$ Integral of their v	M1 A1ft (2)
(c)	Set their $v = 0$ Solve a quadratic in t^2 $(t^2 - 2)(t^2 - 4) = 0 \Rightarrow$ at rest when $t = \sqrt{2}, t = 2$	M1 DM1 A1 (3) [8]

Question Number	Scheme	Marks
4. (a)	<p>Work done against friction = $50 \times \mu R$ $= 50 \times \frac{1}{4} \times 30 \cos 20^\circ \times 9.8$</p> <p>Gain in GPE = $30 \times 9.8 \times 50 \sin 20^\circ$</p> <p>Total work done = WD against Friction + gain in GPE $= 8480(\text{J}), 8500(\text{J})$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>DM1 A1</p> <p>(6)</p>
(b)	<p>Loss in GPE = WD against friction + gain in KE 3 terms</p> <p>$30 \times 9.8 \times 50 \sin 20^\circ = 50 \times \frac{1}{4} \times 30 \times 9.8 \times \cos 20^\circ + \frac{1}{2} \times 30 \times v^2$ -1 ee</p> <p>$\frac{1}{2} v^2 = 50 \times 9.8 \times (\sin 20^\circ - \frac{1}{4} \cos 20^\circ),$</p> <p>$v = 10.2 \text{ m s}^{-1}.$</p>	<p>M1</p> <p>A2, 1, 0</p> <p>DM1 A1</p> <p>(5) [11]</p>

<p>5.</p> <p>(a)</p>	 <p>Divide the shape into usable areas, e.g.:</p> <table border="1" data-bbox="301 734 968 927"> <thead> <tr> <th>Shape</th><th>C of mass</th><th>Units of mass</th></tr> </thead> <tbody> <tr> <td>Rectangle 27 x 9</td><td>(13.5,4.5)</td><td>243 (6)</td></tr> <tr> <td>Right hand triangle</td><td>(30,3)</td><td>40.5 (1)</td></tr> <tr> <td>Top triangle</td><td>(3,30)</td><td>40.5 (1)</td></tr> <tr> <td>Rectangle 9 x 18</td><td>(4.5,18)</td><td>162 (4)</td></tr> </tbody> </table> <p>Mass ratios Centres of mass</p> <p>Take moments about AB:</p> $6 \times 13.5 + 1 \times 30 + 4 \times 4.5 + 1 \times 3 = 132 = 12\bar{x},$ <p>$\bar{x} = 11(\text{cm})$ solve for x (or y) co-ord $\bar{y} = 11(\text{cm})$ using the symmetry</p>	Shape	C of mass	Units of mass	Rectangle 27 x 9	(13.5,4.5)	243 (6)	Right hand triangle	(30,3)	40.5 (1)	Top triangle	(3,30)	40.5 (1)	Rectangle 9 x 18	(4.5,18)	162 (4)	<p>B1 B1</p> <p>M1 A(2,1,0)</p> <p>A1 B1ft</p>
Shape	C of mass	Units of mass															
Rectangle 27 x 9	(13.5,4.5)	243 (6)															
Right hand triangle	(30,3)	40.5 (1)															
Top triangle	(3,30)	40.5 (1)															
Rectangle 9 x 18	(4.5,18)	162 (4)															
	<p>Alternative:</p> <table border="1" data-bbox="301 1301 968 1420"> <thead> <tr> <th>Shape</th><th>C of mass</th><th>Units of mass</th></tr> </thead> <tbody> <tr> <td>Small triangle</td><td>(12,12)</td><td>.5 x 18x 18</td></tr> <tr> <td>Large triangle</td><td>(15,15)</td><td>.5 x 36 x 36</td></tr> </tbody> </table> $\frac{1}{2} \times 36 \times 36 \times 12 - \frac{1}{2} \times 18 \times 18 \times 15 = \frac{1}{2} (36 \times 36 - 18 \times 18) \bar{x} \text{ etc.}$	Shape	C of mass	Units of mass	Small triangle	(12,12)	.5 x 18x 18	Large triangle	(15,15)	.5 x 36 x 36	<p>(7)</p>						
Shape	C of mass	Units of mass															
Small triangle	(12,12)	.5 x 18x 18															
Large triangle	(15,15)	.5 x 36 x 36															
<p>(b)</p>	 $\tan \theta = \frac{\bar{x}}{36 - \bar{y}}$ $\tan \theta = \frac{11}{25} = 0.44$ $\theta = 24^\circ$	<p>M1</p> <p>A1ft</p> <p>A1</p> <p>(3) [10]</p>															

6.	(a)	Using $s = ut + \frac{1}{2}at^2$ clear $\mathbf{r} = (3t)\mathbf{i} + (10 + 5t - 4.9t^2)\mathbf{j}$	Method must be Answer given	M1 A1 A1 (3)
	(b)	\mathbf{j} component = 0: $10 + 5t - 4.9t^2$ quadratic formula: $t = \frac{5 \pm \sqrt{25 + 196}}{9.8} = \frac{5 \pm \sqrt{221}}{9.8}$ $T = 2.03(\text{s}), 2.0(\text{s})$ positive solution only.		M1 DM1 A1 (3)
	(c)	Differentiating the position vector (or working from first principles) $\mathbf{v} = 3\mathbf{i} + (5 - 9.8t)\mathbf{j} \text{ (ms}^{-1}\text{)}$		M1 A1 (2)
	(d)	At B the \mathbf{j} component of the velocity is the negative of the \mathbf{i} component: $5 - 9.8t = -3, 8 = 9.8t,$ $t = 0.82$		M1 A1 (2)
	(e)	$\mathbf{v} = 3\mathbf{i} - 3\mathbf{j}$, speed = $\sqrt{3^2 + 3^2} = \sqrt{18} = 4.24 \text{ (m s}^{-1}\text{)}$		M1A1 (2) [12]

Question Number	Scheme	Marks
7.	 <p>Taking moments about A:</p> $3S = 100 \times 2 \times \cos \alpha$ <p>Resolving vertically:</p> $R + S \cos \alpha = 100$ <p>Resolving horizontally:</p> $S \sin \alpha = F$ <p>(Most alternative methods need 3 independent equations, each one worth M1A1. Can be done in 2 e.g. if they resolve horizontally and take moments about X then $R \times 2 \times \cos \alpha = S \times (3 - 2 \times \cos^2 \alpha)$ scores M2A2)</p> <p>Substitute trig values to obtain correct values for F and R (exact or decimal equivalent).</p> $\left(S = \frac{200\sqrt{8}}{9} \right), R = 100 - \frac{1600}{27} = \frac{1100}{27} \approx 40.74, F = \frac{200\sqrt{8}}{27} \approx 20.95...$ $F \leq \mu R, 200\sqrt{8} \leq \mu \times 1100, \mu \geq \frac{200\sqrt{8}}{1100} = \frac{2\sqrt{8}}{11}.$ <p>Least possible μ is 0.514 (3sf), or exact.</p>	<p>M1 A1</p> <p>M1 A1</p> <p>M1 A1</p> <p>DM1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[10]</p>

Question Number	Scheme	Marks
8. (a)	<p>KE lost : $\frac{1}{2} \times m \times 36 - \frac{1}{2} \times m \times v^2 = 64$</p> <p>Restitution: $v = 1/3 \times 6 = 2$</p> <p>Substitute and solve for m: $\frac{1}{2} \times m \times 36 - \frac{1}{2} \times m \times 4 = 64 = 16m$</p> <p style="text-align: right;">$m = 4$ answer given</p>	<p>M1A1</p> <p>M1A1</p> <p>DM1</p> <p>A1</p> <p style="text-align: right;">(6)</p>
(b)	 <p>Conservation of momentum: $6 - 8 = 4w - 2v$ their "2"</p> <p>Restitution: $v + w = \frac{1}{3}(2 + 3)$ their "2"</p> <p>$v = \frac{5}{3} - w$</p> <p>Solve for w: $-2 = 4w - 2(\frac{5}{3} - w) = 6w - \frac{10}{3}$</p> <p>$\frac{4}{3} = 6w$</p> <p>$(w = 4/18 = 2/9 \text{ m s}^{-1})$</p> <p>$w > 0 \Rightarrow$ will collide with the wall again</p>	<p>M1A1ft</p> <p>M1A1ft</p> <p>DM1</p> <p>A1</p> <p>A1</p> <p style="text-align: right;">(7) [13]</p>

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code UA026580 January 2011

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH