

Static Analysis and Verification

SAV 2022/2023

Tomáš Vojnar

`vojnar@fit.vutbr.cz`

**Brno University of Technology
Faculty of Information Technology
Božetěchova 2, 612 66 Brno**

Lattices and Fixpoints

A Brief Introduction

Partial Orders

- ❖ A tuple $(A, \leq_A)$ is a **poset** (partially-ordered set) iff A is a **set** and $\leq_A \subseteq A \times A$ is a **partial order** (i.e., a reflexive, transitive, and antisymmetric binary relation) on A .
- ❖ An example: Given a set S , $(2^S, \subseteq)$ is a poset.

Partial Orders

- ❖ A tuple $(A, \leq_A)$ is a **poset** (partially-ordered set) iff A is a **set** and $\leq_A \subseteq A \times A$ is a **partial order** (i.e., a reflexive, transitive, and antisymmetric binary relation) on A .
- ❖ An example: Given a set S , $(2^S, \subseteq)$ is a poset.
- ❖ Given a poset $(A, \leq_A)$ and a set $B \subseteq A$,
 - an element $a \in A$ is the **greatest lower bound** of B (glb/infimum/meet of B , $\sqcap B$) iff
 1. $\forall b \in B. a \leq_A b$ (“lower bound”) and
 2. $\forall a' \in A. (\forall b \in B. a' \leq_A b) \implies a' \leq_A a$ (“greatest”),

Partial Orders

- ❖ A tuple $(A, \leq_A)$ is a **poset** (partially-ordered set) iff A is a **set** and $\leq_A \subseteq A \times A$ is a **partial order** (i.e., a reflexive, transitive, and antisymmetric binary relation) on A .
- ❖ An example: Given a set S , $(2^S, \subseteq)$ is a poset.
- ❖ Given a poset $(A, \leq_A)$ and a set $B \subseteq A$,
 - an element $a \in A$ is the **greatest lower bound** of B (glb/infimum/meet of B , $\sqcap B$) iff
 1. $\forall b \in B. a \leq_A b$ (“lower bound”) and
 2. $\forall a' \in A. (\forall b \in B. a' \leq_A b) \implies a' \leq_A a$ (“greatest”),
 - an element $a \in A$ is the **least upper bound** of B (lub/supremum/join of B , $\sqcup B$) iff
 1. $\forall b \in B. b \leq_A a$ (“upper bound”) and
 2. $\forall a' \in A. (\forall b \in B. b \leq_A a') \implies a \leq_A a'$ (“least”),

Partial Orders

- ❖ A tuple $(A, \leq_A)$ is a **poset** (partially-ordered set) iff A is a **set** and $\leq_A \subseteq A \times A$ is a **partial order** (i.e., a reflexive, transitive, and antisymmetric binary relation) on A .
- ❖ An example: Given a set S , $(2^S, \subseteq)$ is a poset.
- ❖ Given a poset $(A, \leq_A)$ and a set $B \subseteq A$,
 - an element $a \in A$ is the **greatest lower bound** of B (glb/infimum/meet of B , $\sqcap B$) iff
 1. $\forall b \in B. a \leq_A b$ (“lower bound”) and
 2. $\forall a' \in A. (\forall b \in B. a' \leq_A b) \implies a' \leq_A a$ (“greatest”),
 - an element $a \in A$ is the **least upper bound** of B (lub/supremum/join of B , $\sqcup B$) iff
 1. $\forall b \in B. b \leq_A a$ (“upper bound”) and
 2. $\forall a' \in A. (\forall b \in B. b \leq_A a') \implies a \leq_A a'$ (“least”),
- ❖ An example: For $(2^{\{a,b,c\}}, \subseteq)$, $\sqcup\{\emptyset, \{a\}, \{b\}\} = \{a, b\}$, and $\{a\} \sqcap \{b, c\} = \emptyset$.

Lattices

- ❖ A poset $(A, \leq_A)$ is a **lattice** iff each *non-empty, finite* subset B of A has a **lub** as well as a **glb** in A .
- ❖ A poset $(A, \leq_A)$ is a **complete lattice** iff each subset B of A has a **lub** as well as a **glb** in A .

Lattices

- ❖ A poset $(A, \leq_A)$ is a **lattice** iff each *non-empty, finite* subset B of A has a **lub** as well as a **glb** in A .
- ❖ A poset $(A, \leq_A)$ is a **complete lattice** iff each subset B of A has a **lub** as well as a **glb** in A .
 - $\perp_A = \bigcap A$ and $\top_A = \bigcup A$ are the least and greatest elements of a complete lattice, respectively.

Lattices

- ❖ A poset $(A, \leq_A)$ is a **lattice** iff each *non-empty, finite* subset B of A has a **lub** as well as a **glb** in A .
- ❖ A poset $(A, \leq_A)$ is a **complete lattice** iff each subset B of A has a **lub** as well as a **glb** in A .
 - $\perp_A = \sqcap A$ and $\top_A = \sqcup A$ are the least and greatest elements of a complete lattice, respectively.
- ❖ Examples:
 - $(2^{\{a,b,c\}}, \subseteq)$ is a complete lattice, $\sqcap$ corresponds to $\cap$, $\sqcup$ to $\cup$, $\perp$ to $\emptyset$, and $\top$ to $\{a, b, c\}$.

Lattices

- ❖ A poset $(A, \leq_A)$ is a **lattice** iff each *non-empty, finite* subset B of A has a **lub** as well as a **glb** in A .
- ❖ A poset $(A, \leq_A)$ is a **complete lattice** iff each subset B of A has a **lub** as well as a **glb** in A .
 - $\perp_A = \sqcap A$ and $\top_A = \sqcup A$ are the least and greatest elements of a complete lattice, respectively.
- ❖ Examples:
 - $(2^{\{a,b,c\}}, \subseteq)$ is a complete lattice, $\sqcap$ corresponds to $\cap$, $\sqcup$ to $\cup$, $\perp$ to $\emptyset$, and $\top$ to $\{a, b, c\}$.
 - $(\mathbb{N}, \leq)$ is a lattice with $\sqcup$ being *max* and $\sqcap$ being *min*, but not a complete lattice since $\sqcup \mathbb{N}$ does not exist in $\mathbb{N}$.

Lattices

- ❖ A poset $(A, \leq_A)$ is a **lattice** iff each *non-empty, finite* subset B of A has a **lub** as well as a **glb** in A .
- ❖ A poset $(A, \leq_A)$ is a **complete lattice** iff each subset B of A has a **lub** as well as a **glb** in A .
 - $\perp_A = \sqcap A$ and $\top_A = \sqcup A$ are the least and greatest elements of a complete lattice, respectively.
- ❖ Examples:
 - $(2^{\{a,b,c\}}, \subseteq)$ is a complete lattice, $\sqcap$ corresponds to $\cap$, $\sqcup$ to $\cup$, $\perp$ to $\emptyset$, and $\top$ to $\{a, b, c\}$.
 - $(\mathbb{N}, \leq)$ is a lattice with $\sqcup$ being *max* and $\sqcap$ being *min*, but not a complete lattice since $\sqcup \mathbb{N}$ does not exist in $\mathbb{N}$.
 - $(\mathbb{N}_\infty, \leq)$, where $\mathbb{N}_\infty = \mathbb{N} \cup \{\infty\}$ and $\forall n \in \mathbb{N}. n \leq \infty$, is a complete lattice.

Lattices

- ❖ A poset $(A, \leq_A)$ is a **lattice** iff each *non-empty, finite* subset B of A has a **lub** as well as a **glb** in A .
- ❖ A poset $(A, \leq_A)$ is a **complete lattice** iff each subset B of A has a **lub** as well as a **glb** in A .
 - $\perp_A = \sqcap A$ and $\top_A = \sqcup A$ are the least and greatest elements of a complete lattice, respectively.
- ❖ Examples:
 - $(2^{\{a,b,c\}}, \subseteq)$ is a complete lattice, $\sqcap$ corresponds to $\cap$, $\sqcup$ to $\cup$, $\perp$ to $\emptyset$, and $\top$ to $\{a, b, c\}$.
 - $(\mathbb{N}, \leq)$ is a lattice with $\sqcup$ being *max* and $\sqcap$ being *min*, but not a complete lattice since $\sqcup \mathbb{N}$ does not exist in $\mathbb{N}$.
 - $(\mathbb{N}_\infty, \leq)$, where $\mathbb{N}_\infty = \mathbb{N} \cup \{\infty\}$ and $\forall n \in \mathbb{N}. n \leq \infty$, is a complete lattice.
- ❖ Given a poset $(A, \leq_A)$, a set $B \subseteq A$ is a **chain** iff $\forall b, b' \in B. b \leq_A b' \vee b' \leq_A b$.
 - E.g., $\{\emptyset, \{a\}, \{a, b\}, \{a, b, c\}\}$ is a chain wrt. $(2^{\{a,b,c\}}, \subseteq)$.

Functions on Lattices

❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.

❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.

Functions on Lattices

- ❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.
- ❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.
- ❖ A function $f : A \longrightarrow B$ is **$\sqcup$ -continuous** iff for every chain $C \subseteq A$, we have $f(\sqcup C) = \sqcup \{f(c) \mid c \in C\}$. Analogously, one can define **$\sqcap$ -continuous** functions.

Functions on Lattices

- ❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.
- ❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.
- ❖ A function $f : A \longrightarrow B$ is **$\sqcup$ -continuous** iff for every chain $C \subseteq A$, we have $f(\sqcup C) = \sqcup \{f(c) \mid c \in C\}$. Analogously, one can define **$\sqcap$ -continuous** functions.
- ❖ An element $a \in A$ is a **fixpoint** of a function $f : A \longrightarrow A$ iff $f(a) = a$.

Functions on Lattices

- ❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.
- ❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.
- ❖ A function $f : A \longrightarrow B$ is **$\sqcup$ -continuous** iff for every chain $C \subseteq A$, we have $f(\sqcup C) = \sqcup \{f(c) \mid c \in C\}$. Analogously, one can define **$\sqcap$ -continuous** functions.
- ❖ An element $a \in A$ is a **fixpoint** of a function $f : A \longrightarrow A$ iff $f(a) = a$.
- ❖ An example: Consider the function $f : \mathbb{N}_\infty \longrightarrow \mathbb{N}_\infty$ defined such that $\forall n \in \mathbb{N}. f(n) = 0$ and $f(\infty) = \infty$.

Functions on Lattices

❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.

❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.

❖ A function $f : A \longrightarrow B$ is **$\sqcup$ -continuous** iff for every chain $C \subseteq A$, we have $f(\sqcup C) = \sqcup \{f(c) \mid c \in C\}$. Analogously, one can define **$\sqcap$ -continuous** functions.

❖ An element $a \in A$ is a **fixpoint** of a function $f : A \longrightarrow A$ iff $f(a) = a$.

❖ An example: Consider the function $f : \mathbb{N}_\infty \longrightarrow \mathbb{N}_\infty$ defined such that $\forall n \in \mathbb{N}. f(n) = 0$ and $f(\infty) = \infty$.

- f is **monotonic** since (1) $\forall n_1, n_2 \in \mathbb{N}. f(n_1) = 0 \leq 0 = f(n_2)$ and (2) $\forall n \in \mathbb{N}. f(n) = 0 \leq \infty = f(\infty)$.

Functions on Lattices

- ❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.
- ❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.
- ❖ A function $f : A \longrightarrow B$ is **$\sqcup$ -continuous** iff for every chain $C \subseteq A$, we have $f(\sqcup C) = \sqcup \{f(c) \mid c \in C\}$. Analogously, one can define **$\sqcap$ -continuous** functions.
- ❖ An element $a \in A$ is a **fixpoint** of a function $f : A \longrightarrow A$ iff $f(a) = a$.
- ❖ An example: Consider the function $f : \mathbb{N}_\infty \longrightarrow \mathbb{N}_\infty$ defined such that $\forall n \in \mathbb{N}. f(n) = 0$ and $f(\infty) = \infty$.
 - f is **monotonic** since (1) $\forall n_1, n_2 \in \mathbb{N}. f(n_1) = 0 \leq 0 = f(n_2)$ and (2) $\forall n \in \mathbb{N}. f(n) = 0 \leq \infty = f(\infty)$.
 - f is **not $\sqcup$ -continuous** since $\mathbb{N}$ is a chain and $f(\sqcup \mathbb{N}) = f(\infty) = \infty$, but $\sqcup \{f(n) \mid n \in \mathbb{N}\} = \sqcup \{0\} = 0$.

Functions on Lattices

- ❖ Let $(A, \leq_A)$ and $(B, \leq_B)$ be lattices.
- ❖ A function $f : A \longrightarrow B$ is **monotonic** iff $\forall a, a' \in A. a \leq_A a' \implies f(a) \leq_B f(a')$.
- ❖ A function $f : A \longrightarrow B$ is **$\sqcup$ -continuous** iff for every chain $C \subseteq A$, we have $f(\sqcup C) = \sqcup \{f(c) \mid c \in C\}$. Analogously, one can define **$\sqcap$ -continuous** functions.
- ❖ An element $a \in A$ is a **fixpoint** of a function $f : A \longrightarrow A$ iff $f(a) = a$.
- ❖ An example: Consider the function $f : \mathbb{N}_\infty \longrightarrow \mathbb{N}_\infty$ defined such that $\forall n \in \mathbb{N}. f(n) = 0$ and $f(\infty) = \infty$.
 - f is **monotonic** since (1) $\forall n_1, n_2 \in \mathbb{N}. f(n_1) = 0 \leq 0 = f(n_2)$ and (2) $\forall n \in \mathbb{N}. f(n) = 0 \leq \infty = f(\infty)$.
 - f is **not $\sqcup$ -continuous** since $\mathbb{N}$ is a chain and $f(\sqcup \mathbb{N}) = f(\infty) = \infty$, but $\sqcup \{f(n) \mid n \in \mathbb{N}\} = \sqcup \{0\} = 0$.
 - 0 is the **least fixpoint** of f and ∞ is the **greatest fixpoint** of f .

Knaster–Tarski Theorem

- ❖ **Knaster–Tarski Theorem.** Let $(A, \leq_A)$ be a complete lattice and let $f : A \longrightarrow A$ be a monotonic function. Then the set of fixpoints of f in $(A, \leq_A)$ is also a complete lattice.
- ❖ Since complete lattices have the least and the greatest element, the theorem in particular guarantees the existence of a **least** and **greatest fixpoint** of f in A .

Knaster–Tarski Theorem

❖ **Knaster–Tarski Theorem.** Let $(A, \leq_A)$ be a complete lattice and let $f : A \longrightarrow A$ be a monotonic function. Then the set of fixpoints of f in $(A, \leq_A)$ is also a complete lattice.

❖ Since complete lattices have the least and the greatest element, the theorem in particular guarantees the existence of a **least** and **greatest fixpoint** of f in A .

For the more curious:

❖ In more constructive terms, the least fixpoint of f is the stationary limit of $f^\alpha(\perp_A)$ for α ranging over the ordinals.

- An **ordinal** is the order type of a well-ordered set.

Knaster–Tarski Theorem

❖ **Knaster–Tarski Theorem.** Let $(A, \leq_A)$ be a complete lattice and let $f : A \longrightarrow A$ be a monotonic function. Then the set of fixpoints of f in $(A, \leq_A)$ is also a complete lattice.

❖ Since complete lattices have the least and the greatest element, the theorem in particular guarantees the existence of a **least** and **greatest fixpoint** of f in A .

For the more curious:

❖ In more constructive terms, the least fixpoint of f is the stationary limit of $f^\alpha(\perp_A)$ for α ranging over the ordinals.

- An **ordinal** is the order type of a well-ordered set.
- Every ordinal can be represented as the set of all smaller ordinals. There is the **zero ordinal**, **successor ordinals**, and **limit ordinals**. Natural numbers correspond to the so called finite ordinals (ordering types of finite sets), the set of natural numbers is the first infinite ordinal, and so on.

Knaster–Tarski Theorem

❖ **Knaster–Tarski Theorem.** Let $(A, \leq_A)$ be a complete lattice and let $f : A \longrightarrow A$ be a monotonic function. Then the set of fixpoints of f in $(A, \leq_A)$ is also a complete lattice.

❖ Since complete lattices have the least and the greatest element, the theorem in particular guarantees the existence of a **least** and **greatest fixpoint** of f in A .

For the more curious:

❖ In more constructive terms, the least fixpoint of f is the stationary limit of $f^\alpha(\perp_A)$ for α ranging over the ordinals.

- An **ordinal** is the order type of a well-ordered set.
- Every ordinal can be represented as the set of all smaller ordinals. There is the **zero ordinal**, **successor ordinals**, and **limit ordinals**. Natural numbers correspond to the so called finite ordinals (ordering types of finite sets), the set of natural numbers is the first infinite ordinal, and so on.
- f^α is defined by **transfinite induction**: $f^{\alpha+1} = f(f^\alpha)$ and f^γ for a limit ordinal γ is the least upper bound of f^β for all ordinals β smaller than γ .

Knaster–Tarski Theorem

❖ **Knaster–Tarski Theorem.** Let $(A, \leq_A)$ be a complete lattice and let $f : A \longrightarrow A$ be a monotonic function. Then the set of fixpoints of f in $(A, \leq_A)$ is also a complete lattice.

❖ Since complete lattices have the least and the greatest element, the theorem in particular guarantees the existence of a **least** and **greatest fixpoint** of f in A .

For the more curious:

❖ In more constructive terms, the least fixpoint of f is the stationary limit of $f^\alpha(\perp_A)$ for α ranging over the ordinals.

- An **ordinal** is the order type of a well-ordered set.
- Every ordinal can be represented as the set of all smaller ordinals. There is the **zero ordinal**, **successor ordinals**, and **limit ordinals**. Natural numbers correspond to the so called finite ordinals (ordering types of finite sets), the set of natural numbers is the first infinite ordinal, and so on.
- f^α is defined by **transfinite induction**: $f^{\alpha+1} = f(f^\alpha)$ and f^γ for a limit ordinal γ is the least upper bound of f^β for all ordinals β smaller than γ .

❖ A dual result holds for the **greatest fixpoint**.

Kleene Fixpoint Theorem

- ❖ **Kleene Fixpoint Theorem.** Let $(A, \leq_A)$ be a complete lattice and $f : A \longrightarrow A$ a function.
- If f is $\sqcup$ -continuous, the least fixpoint of f is $\mu f = \sqcup \{f^i(\perp_A) \mid i \geq 0\}$.

Kleene Fixpoint Theorem

❖ **Kleene Fixpoint Theorem.** Let $(A, \leq_A)$ be a complete lattice and $f : A \longrightarrow A$ a function.

- If f is $\sqcup$ -continuous, the least fixpoint of f is $\mu f = \sqcup \{f^i(\perp_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcup$ -continuous function is monotone, and hence one in fact computes the supremum of the ascending chain
 $\perp_A \leq_A f(\perp_A) \leq_A f(f(\perp_A)) \leq \dots$

Kleene Fixpoint Theorem

❖ **Kleene Fixpoint Theorem.** Let $(A, \leq_A)$ be a complete lattice and $f : A \longrightarrow A$ a function.

- If f is $\sqcup$ -continuous, the least fixpoint of f is $\mu f = \sqcup \{f^i(\perp_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcup$ -continuous function is monotone, and hence one in fact computes the supremum of the ascending chain
$$\perp_A \leq_A f(\perp_A) \leq_A f(f(\perp_A)) \leq \dots$$
- If f is $\sqcap$ -continuous, the greatest fixpoint of f is $\nu f = \sqcap \{f^i(\top_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcap$ -continuous function is monotone, and hence one in fact computes the infimum of the descending chain
$$\top_A \geq_A f(\top_A) \geq_A f(f(\top_A)) \geq \dots$$

Kleene Fixpoint Theorem

❖ **Kleene Fixpoint Theorem.** Let $(A, \leq_A)$ be a complete lattice and $f : A \longrightarrow A$ a function.

- If f is $\sqcup$ -continuous, the least fixpoint of f is $\mu f = \sqcup \{f^i(\perp_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcup$ -continuous function is monotone, and hence one in fact computes the supremum of the ascending chain
$$\perp_A \leq_A f(\perp_A) \leq_A f(f(\perp_A)) \leq \dots$$
- If f is $\sqcap$ -continuous, the greatest fixpoint of f is $\nu f = \sqcap \{f^i(\top_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcap$ -continuous function is monotone, and hence one in fact computes the infimum of the descending chain
$$\top_A \geq_A f(\top_A) \geq_A f(f(\top_A)) \geq \dots$$

❖ **Theorem.** For finite complete lattices, every monotonic function is $\sqcap$ - and $\sqcup$ -continuous.

Kleene Fixpoint Theorem

❖ **Kleene Fixpoint Theorem.** Let $(A, \leq_A)$ be a complete lattice and $f : A \longrightarrow A$ a function.

- If f is $\sqcup$ -continuous, the least fixpoint of f is $\mu f = \sqcup\{f^i(\perp_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcup$ -continuous function is monotone, and hence one in fact computes the supremum of the ascending chain
$$\perp_A \leq_A f(\perp_A) \leq_A f(f(\perp_A)) \leq \dots$$
- If f is $\sqcap$ -continuous, the greatest fixpoint of f is $\nu f = \sqcap\{f^i(\top_A) \mid i \geq 0\}$.
 - Moreover, a $\sqcap$ -continuous function is monotone, and hence one in fact computes the infimum of the descending chain
$$\top_A \geq_A f(\top_A) \geq_A f(f(\top_A)) \geq \dots$$

❖ **Theorem.** For finite complete lattices, every monotonic function is $\sqcap$ - and $\sqcup$ -continuous.

❖ **Corollary.** On finite lattices, the Kleene fixpoint theorem is applicable, hence,

- to compute the least fixpoint, start with $\perp_A$ and iteratively apply f till
$$f^i(\perp_A) = f^{i+1}(\perp_A) = \mu f,$$
- to compute the greatest fixpoint, start with $\top_A$ and iteratively apply f till
$$f^i(\top_A) = f^{i+1}(\top_A) = \nu f.$$