

SATELLITE ALIGNMENTS IN GALAXY PAIRS

VERÓNICA ARIAS ¹, JAIME E. FORERO-ROMERO ²

¹Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio Ip, Bogotá, Colombia

Submitted for publication in ApJ

ABSTRACT

Alineaciones.

Subject headings: Galaxies: halos — Galaxies: high-redshift — Galaxies: statistics — Dark Matter
 — Methods: numerical

1. INTRODUCTION

Since the early works of (el que libeskind dice que habl de eso primero) and Lynden-Bell (1976), the observed anisotropic distribution of satellite galaxies in the Local Group has been a key point in the discussion of galaxy formation models within Λ CDM cosmology. These pioneer papers described that the then known satellite galaxies of the Milky Way (MW) were distributed roughly following the plane that contained the Magellanic clouds and their stellar tails. This anisotropic distribution was further studied in the work of Pawlowski et. al. (2012), where they included all the known MW satellite galaxies (XXX more than in the original Lynden-Bell paper) as well as some MW globular clusters, and found that they are all contained in what they call a "Vast Plane Of Satellites" (VPOS). This structure is around 30kpc thick and is perpendicular to the stellar disc of the MW galaxy. This plane of satellites was considered a rarity until the PAndAS survey of the Andromeda (M31) galaxy and its halo (missing reference on the survey) provided a complete sample of the satellite galaxy population (above a XXXX magnitude) of our neighboring galaxy, allowing their consistent distance estimations (Conn et al 2012). In a groundbreaking work, Ibata et. al. (2013) found that 15 out of the 30 satellites are in a planar structure that is 15kpc thick and has an extension of 400kpc. Additionally, from line of sight velocity measurements, they found that the structure has an apparent coherent rotation, with the satellites south of M31 moving away from us and those north of M31 coming towards us. This apparent organized rotation of satellite galaxies was also statistically found to occur in diametrically opposed pairs of satellites from the SDSS (Ibata et al 2014), although this results were contested in a follow up paper by Cuatun et al (2014).

These discoveries of the planes in M31 and the MW were followed by the remark from Shaya et al (2013) that the other satellite galaxies in M31 could be part of a second plane, further reinforcing the idea that satellites in the local group are not isotropically distributed. Despite the challenges in estimating distances to the satellites galaxies, Tully et al. managed to go beyond the local group and found two planes of satellites galaxies in Centaurus A, making a clearer case for the planes of satellites; in all the galaxies where distance measurements could be performed evidence of satellite planar structures have been found.

These observed planes present a challenge for current galaxy formation models (Pawlowski et. al 2014)

A comparison of the distribution of satellite galaxies around Andromeda and the results of Λ CDM simulations (Bahl and Baumgardt 2013) Encuentran planos en Millenium II.

A thousand shadows of Andromeda: rotating planes of satellites in the Millennium-II cosmological simulation (Ibata et al 2013) dicen que el paper de Bahl y Baumgardt está mal y que no hay planos en millenium II

Co-orbiting satellite galaxy structures are still in conflict with the distribution of primordial dwarf galaxies (Pawlowski et al 2014)

Co-orbiting planes of sub-halos are similarly unlikely around paired and isolated hosts (Pawlowski et al 2014) Vast planes of satellites in a high resolution simulation of the Local Group: comparison to Andromeda (Gillet et al 2014). finds a plnes similar to that of M31 in Clues simulations.

Planes of satellite galaxies: when exceptions are the rule (Cautun et al 2015) encuentra que 10 por ciento de los halos tienen planos iguales o más prominentes que los del LG.

1.1. Posibles orígenes de los planos:

Preferential accretion (Libeskind et al 20??)

Alignments with the cosmic web (Tempel et al 20??)

The vast thin plane of M31 co-rotating dwarfs: an additional fossil signature of the M31 merger and of its considerable impact in the whole Local Group (Hammer et al 2013) Major merger in M31-MW system plane galaxies are tidal dwarfs (n-body simulations)

Kroupa tiene todo un carretazo de que las dwarfs son todas tidal dwarfs

MIRAR lo que está haciendo Pierre Alan Duc con tidal dwarfs porque en una conferencia este a;o habló de un posible escenario intermedio...

Evidence for Early Filamentary Accretion from the Andromeda Galaxy's Thin Plane of Satellites (Buck et al 2015)

Alignments between galaxies, satellite systems and haloes (Cautun et al 2016) NO LO HE LEIDO...

1.2. Problemas con esas explicaciones:

The Vast Polar Structure of the Milky Way and Filamentary Accretion of Sub-Halos (Pawlowski et al