

## RESEARCH ARTICLE

# Complications of biliary stenting versus T-tube insertion after common bile duct exploration: A systematic review and meta-analysis

Reno Rudiman<sup>1\*</sup>, Ricarhdo Valentino Hanafi<sup>2</sup>, Almawijaya<sup>1</sup>, Freda Halim<sup>3</sup>

**1** Division of Digestive Surgery, Department of General Surgery, School of Medicine, Universitas Padjadjaran, Hasan Sadikin General Hospital, Bandung, Indonesia, **2** Faculty of Medicine, Universitas Pelita Harapan, Tangerang, Indonesia, **3** Department of General Surgery, Faculty of Medicine, Universitas Pelita Harapan, Tangerang, Indonesia

\* [rudiman@unpad.ac.id](mailto:rudiman@unpad.ac.id)



## Abstract

### Background

Complications following the insertion T-tube or stent after common bile duct exploration (CBDE) remain problematic in nowadays surgical era. Based on our knowledge, we did not find any meta-analysis intentionally evaluating the complications between both groups. At this moment, we aimed to analyze and compare both procedures' complications, efficacy, efficiency, and feasibility.

### Methods

We searched literature from four databases (EuroPMC, PubMed, Scopus, and Clinical-Trials.gov) up to June 2022 to compile the randomized controlled trials and pro-/retrospective cohort studies. Review Manager 5.4 was used to statistically analyze each outcome measured between biliary stenting and T-tube insertion.

### Results

Sixteen studies with 1,080 patients (534 biliary stents and 546 T-tube) were included for qualitative and quantitative analysis. The pooled risk ratio (RR) of the overall postoperative complications rate was significantly lower in the biliary stent group compared to the T-tube group 0.43 [95% confidence interval (CI) 0.23–0.80,  $p = 0.007$ ]. In terms of the operation time, length of hospital stay, and readmission rate was also decreased in stenting as biliary drainage over T-tube placement 1.02 minutes [95% CI -1.53, -0.52,  $p < 0.0001$ ], 1.96 days [95% CI -2.63, -1.29,  $p < 0.00001$ ], and RR 0.39 [95% CI 0.15–0.97,  $p = 0.04$ ], respectively.

### Conclusions

Stenting as biliary drainage after CBDE was superior to T-tube insertion. A shorter operation time and hospital stay in biliary drainage resulted in a lower overall postoperative complication rate. Other influences, including the complexity and shorter learning curve, might also affect the superiority of biliary stenting.

## OPEN ACCESS

**Citation:** Rudiman R, Hanafi RV, Almawijaya, Halim F (2023) Complications of biliary stenting versus T-tube insertion after common bile duct exploration: A systematic review and meta-analysis. PLoS ONE 18(1): e0280755. <https://doi.org/10.1371/journal.pone.0280755>

**Editor:** Wenguo Cui, Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital, CHINA

**Received:** October 29, 2022

**Accepted:** January 6, 2023

**Published:** January 20, 2023

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0280755>

**Copyright:** © 2023 Rudiman et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data are available from the following repository: <https://doi.org/10.5281/zenodo.7263085>.

**Funding:** The author(s) received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Common bile duct exploration (CBDE), either open or laparoscopically, is a procedure to treat and prevent further complications such as cholangitis, pancreatitis, and obstructive jaundice due to common bile duct stones. Choledocholithiasis may be a silent disease that, over time, generates further troubles [1]. Up to 20% of patients with cholelithiasis were classified as choledocholithiasis [2].

Following the CBDE, biliary drainage with T-tube or biliary stent is inserted within the common bile duct to prevent biliary leakage, stenosis, and other choledocholithiasis [3, 4]. This T-tube insertion was commonly performed before the endoscopic retrograde cholangiopancreatography (ERCP) era, the less invasive and renowned procedure in which safer internal biliary drainage could be achieved [5, 6]. Conversely, ERCP was not frequently utilized in developing countries due to the inequality of healthcare services.

The pioneer of T-tube drainage following choledochotomy was Deaver in 1904 [7]. Up to this day, T-tube has four types of material: polyvinyl chloride (PVC), red rubber, silicone, and latex. The latex T-tube is the most commonly used due to its superiority in complications, inflammatory reaction, and low bile sedimentation within the lumen. Generally, T-tube may be removed within 7 to 10 days if all clinical assessments improve [8]. It was a standard technique up to two decades ago due to its high association with numerous complications such as leakage, biliary peritonitis, T-tube dislodgment, vascular injury, and fistula. This procedure is a complex technique and requires an experienced operator to minimize the complications. Further damages, such as biliary leakage and biliary peritonitis, might occur when the T-tube is taken [9–11].

A biliary stent is a cylindrical device (plastic or metal) frequently used to establish the patency of the bile duct. Plastic and self-expanding metal stents (SEMs) showed a similar success rate in relieving bile duct stones and bile leakage [12]. Additionally, biliary stent removal can be removed safely endoscopically, which minimizes additional complications [13]. The previous report demonstrated the biliary complications from T-tube do not differ in open and laparoscopic procedures (15.5% vs 13.8%, respectively), and those complications were more related to the placement and removal of the T-tube drainage itself [14, 15]. Due to these controversies among surgeons, a shift towards biliary stenting occurred, demonstrating better outcomes and lower complication rates [16–18]. Previous meta-analysis conducted by Yin *et al.* 2013 showed primary duct closure (PDC) alone was superior to combination with T-tube insertion [19]. Moreover, Jiang *et al.* 2019 also stated that PDC alone or PDC combined with internal drainage (stenting) was superior to PDC with T-tube insertion [20].

Based on our awareness, we did not find any meta-analysis that purposefully compared T-tube complications and biliary stent drainage after CBDE (open or laparoscopically). In developed countries, T-tube insertion and open CBDE were left behind a decade ago. However, many developing countries still undergo those procedures due to the limitations of healthcare facilities. Consequently, we aimed to analyze and compare both procedures' complications, efficacy, efficiency, and feasibility.

## Materials & methods

### Eligibility criteria

The inclusion criteria are framed with the PICOS formula. (1) Population: patients with common bile duct (CBD) exploration (either open or laparoscopic), (2) Intervention: patients who were receiving biliary stenting, (3) Comparison: patients who were receiving T-tube drainage, (4) Primary outcomes: biliary leakage, wound infection, pancreatitis, overall postoperative

complications; secondary outcomes: operation time, blood loss volume, length of hospital stay, readmission rate, and reoperation rate, and lastly, (5) Study: randomized or non-randomized controlled trials, cohort, case-control, cross-sectional, and case-series.

We excluded any study with a population under 18 years old, pregnant women, and no control group. Non-English and Non-original studies, including review articles, editorials, letters to the editor, correspondences, and conference abstracts were also exclusions. This study concurs with the preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines and is registered in the international prospective register of systematic reviews (PROSPERO) CRD42022360170.

### Information sources

We conducted literature searches in four databases (EuroPMC, PubMed, Scopus, and Clinical-Trials.gov) until June 2022 with English-language restrictions.

### Search strategy

The following keywords of this study are “(biliary OR bile duct OR common bile duct OR CBD) AND (exploration OR surgery OR open-choledochotomy OR laparoscopic OR decompression) AND (stent OR stenting) AND (T-tube)”.

### Selection process

Initial screening of titles and abstracts with Covidence software (<https://covidence.org>) by two researchers (RR, RVH) and any discrepancies were solved with the third researcher (FH) discussion and consultation.

### Data collection process

Full-text screening and data extractions (author details, patient’s characteristics, study design, outcomes measured) were performed by RR and RVH independently. Furthermore, FH merged and validated the two extracted data. Any incongruity was resolved through discussion with the third researcher.

### Data items

The outcomes were operation time (from the first incision until wound closure), blood loss volume (total blood loss during operation), length of hospital stay (as defined by each study), biliary leakage (definition of leakage by each study), wound infection (definition of infection by each study), pancreatitis (as defined by each study), postoperative complications (determined by each study), readmission rate (hospitalized due to specific reason after operation), and reoperation rate (second operation due to particular reason after the procedure).

### Study risk of bias assessment

We used the Newcastle-Ottawa scale (NOS) to evaluate the quality of case-control and cohort studies. The assessment reviews each study’s comparability, selection, and outcome [21]. Meanwhile, a modified Jadad scale is a procedure to assess the quality of included clinical trial studies. It evaluates the four domains: the randomization process, allocation concealment, blindness, and withdrawals and drop-outs of the reported results [22]. Two researchers (RR, RVH) evaluated those risk of bias assessments, and a third researcher (FH) interceded with any divergence.

## Effect measures

The effect measures used in synthesizing results depend on the variable outcomes. We used a risk ratio (RR) with a 95% confidence interval (CI) for the dichotomous variable and standardized mean difference (SMD) and its standard deviations (SD) for the continuous variable. The summarized synthesis is presented in the forest plot.

## Synthesis methods

We utilized Review Manager 5.4 software for conducting meta-analysis. A Mantel-Haenszel formula with random-effect models was applied to calculate the RR and 95% CI. This formula allows to calculate an unconfounded, overall effect estimate of a given exposure for a specific outcome by combining the RR [23]. Meanwhile, an Inverse-Variance formula with random-effect models was used to assess the SMD and SD. This method involves a weighted average of the effect estimates from the separate studies, and the weight in each study is the inverse of the variance of the effect estimate [24]. We synthesized the heterogeneity using  $I^2$  statistics. A 50% heterogeneity is considered substantial heterogeneity, which indicates half of the variability among effect sizes is due to significant heterogeneity among studies. Moreover, we applied Wan X *et al.* [25] formula to convert means and standard deviations (SD) for meta-analysis data synthesis.

## Publication bias

We employed funnel plots analysis to visualize the qualitative risk of publication bias.

## Results

### Study selection

Our initial databases search revealed 838 titles and abstracts. After removing the duplicates, screening titles, and abstracts, 63 publications were eligible for further full-text assessment. The final 16 full-text publications were qualified for qualitative and quantitative synthesis [26–41]. However, we excluded 47 publications for several reasons, including 27 articles that did not include biliary stenting, 12 articles that did not have any comparison or control group, five articles that were not primary research, and three articles that were not English papers. The summarized PRISMA flow chart of study selection is presented in Fig 1.

### Study characteristics


The final 16 full-text publications included 1,080 patients (534 biliary stents and 546 T-tube), published between 2004–2022. The study design were five prospective RCT studies, eight retrospective cohort studies, and three prospective cohort studies. Most of the included study participants received 7 or 10-Fr biliary stents. The detailed study characteristics are summarized in Table 1.

### Risk of bias in studies

The results of NOS indicated all the included studies had good quality (Table 2). The modified Jadad scale revealed one RCT was graded as a high-quality study; on the other hand, four RCTs were graded as moderate-quality studies (Table 3).

### Biliary stent vs. T-tube

**Bile leakage.** The pooled analysis from 12 studies ( $n = 811$ ) demonstrated no differences between biliary stenting and T-tube drainage in bile leakage incidence [RR 0.67 (95% CI 0.36–1.27),  $p = 0.22$ ,  $I^2 = 0\%$ , random-effect modeling] (Fig 2A).



**Fig 1. The PRISMA flowchart of the studies selection process.**

<https://doi.org/10.1371/journal.pone.0280755.g001>

**Wound infection.** In the pooled analysis of six studies ( $n = 269$ ), biliary stenting and T-tube drainage after CBD exploration did not demonstrate any differences in wound infection outcome [RR 0.37 (95% CI 0.12–1.13),  $p = 0.08$ ,  $I^2 = 12\%$ , random-effect modeling] (Fig 2B).

**Pancreatitis.** In the pooled analysis of eight studies ( $n = 363$ ), the pancreatitis incidence after CBD exploration did not differ between biliary stenting and T-tube drainage [RR 0.55 (95% CI 0.13–2.28),  $p = 0.41$ ,  $I^2 = 41\%$ , random-effect modeling] (Fig 2C).

**Overall postoperative complications.** In the pooled analysis of 16 studies ( $n = 1,080$ ), the overall postoperative complications were lowered in the biliary stenting group than in the T-tube group after CBD exploration [RR 0.43 (95% CI 0.23–0.80),  $p = 0.007$ ,  $I^2 = 63\%$ , random-effect modeling] (Fig 2D).

Table 1. Data characteristics of the included studies.

First author, year	Sample size		Design	Age (years)		Gender (male/female)		Stent used	Outcomes*
	Stent	T-tube		Stent	T-tube	Stent	T-tube		
Abdelkader AM et al. [26] 2018	25	25	Prospective cohort	41.2 ± 8.1	40.6 ± 7.5	11/14	10/15	8 or 10 Fr Nelaton tube	1,2,3,4,7
Dietrich A et al. [27] 2014	13	35	Retrospective cohort	60.2 ± 12.5	64.1 ± 15	9/5	18/17	7-Fr plastic biliary stent (Cotton-Leung)	1,3,6,7,9
Gyalpo T et al. [28] 2018	20	20	Prospective RCT	51.1 ± 11.5	50.6 ± 12	4/16	3/17	7-Fr 10cm straight flap biliary stent (Indovasive, India)	1,3,4,5,7,8,9
Hassan AM et al. [29] 2017	15	9	Retrospective cohort	49.5 ± 6.6	49.7 ± 6	15/6	5/4	7-Fr plastic stent	1,3,6,7,9
Isla AM et al. [30] 2004	21	32	Retrospective cohort	54 ± 14	65.1 ± 17.7	5/16	9/23	10-Fr biliary stent (Olympus)	1,3,4,7
Kwon SU et al. [31] 2011	23	29	Retrospective cohort	73 ± 9.1	69.7 ± 10.5	5/18	12/17	Not mentioned in details	1,3,4,7
Xiao L et al. [32] 2018	120	116	Retrospective cohort	53.8 ± 12.8	51 ± 12.5	52/68	54/62	7 to 10-Fr biliary stent (ENDO-FLEX GmbH Company)	1,3,4,7,8
Lyon M et al. [33] 2015	82	34	Prospective cohort	43.8 ± 10	45.9 ± 12.5	21/61	6/28	7- Fr straight or duodenal curve biliary Diagmed™ stent	3,7,9
Mangla V et al. [34] 2012	31	29	Prospective RCT	46.8 ± 14.8	47.1 ± 12.3	9/22	5/24	7 or 10-Fr, 10-cm straight flap biliary stent (Devon, Agra, India)	1,2,3,4,5,7
Abd El Wahab A et al. [35] 2022	10	10	Prospective cohort	48.7 ± 4.9	53.1 ± 6.9	N/A	N/A	Not described in detail	1,2,3,4,5,6,7
Martinez-Baena D et al. [36] 2013	28	47	Retrospective cohort	61.9 ± 15.2	61.1 ± 16.9	9/19	24/23	Biliary stenting (Flexima®, Boston Scientific Corporation)	3,4,6,7,8,9
Mir IS et al. [37] 2016	11	20	Retrospective cohort	N/A	N/A	N/A	N/A	7 or 10-Fr biliary stent	1,3,4,5,6,7
Parra-Membrives P et al. [38] 2017	58	52	Retrospective cohort	63.2 ± 15.7	60.7 ± 15.2	18/40	25/27	6 to 8.5-Fr plastic biliary stent	3,4,7,8,9
Perez G et al. [39] 2005	37	44	Prospective RCT	56.9 ± 18.7	55 ± 16.3	34/10	28/9	5 or 6-Fr biliary stent (Zimmon, Wilson-Cook Medical Winston-Salem, NC, USA)	3,5,6,7,8,9
Redwan AA et al. [40] 2006	18	19	Prospective RCT	46 ± 12	46 ± 12	N/A	N/A	8.5-Fr biliary stent	1,3,4,5,6,7
Xu Y et al. [41] 2013	22	25	Prospective RCT	61.3 ± 21.8	62.1 ± 15.5	9/13	14/11	8.5 to 10-Fr biliary stent (Leadgem Medical Co., serial number FY0825)	1,2,3,4,6,7,9

\*Outcomes: 1 = operation time; 2 = blood loss volume; 3 = length of hospital stay; 4 = bile leakage; 5 = wound infection; 6 = pancreatitis; 7 = post-operative complications; 8 = readmission rate; 9 = reoperations rate

<https://doi.org/10.1371/journal.pone.0280755.t001>

**Operation time.** In the pooled analysis from 12 studies (n = 698), biliary stenting as the drainage method after CBD exploration demonstrated a shorter operative time than T-tube [SMD -1.02 (95% CI -1.53, -0.52),  $p < 0.0001$ ,  $I^2 = 88\%$ , random-effect modeling] (Fig 2E).

**Blood loss volume.** The pooled analysis from four studies (n = 177) demonstrated both biliary stenting and T-tube drainage after CBD exploration did not differ in terms of blood loss volume [SMD -0.45 (95% CI -1.11, 0.21),  $p = 0.18$ ,  $I^2 = 77\%$ , random-effect modeling] (Fig 3A).

**Length of hospital stay.** In the pooled analysis from 16 studies (n = 1,080), the length of hospital stay was shorter in the biliary stenting method than T-tube after CBD exploration [SMD -1.96 (95% CI -2.63, -1.29),  $p < 0.00001$ ,  $I^2 = 95\%$ , random-effect modeling] (Fig 3B).

**Readmission rate.** In the pooled analysis of four studies (n = 306), the biliary stenting group exhibited a lower hospital readmission rate compared with the T-tube group [RR 0.39 (95% CI 0.15–0.97),  $p = 0.04$ ,  $I^2 = 0\%$ , random-effect modeling] (Fig 3C).

Table 2. Newcastle-Ottawa quality assessment of observational studies.

First author, year	Study design	Selection <sup>a</sup>	Comparability <sup>b</sup>	Outcome <sup>c</sup>	Total score	Result
Abdelkader AM et al. [26] 2018	Cohort	***	**	**	7	Good
Dietrich A et al. [27] 2014	Cohort	***	**	***	8	Good
Hassan AM et al. [29] 2017	Cohort	***	**	**	7	Good
Isla AM et al. [30] 2004	Cohort	***	**	***	8	Good
Kwon SU et al. [31] 2011	Cohort	***	**	**	7	Good
Xiao L et al. [32] 2018	Cohort	***	**	***	8	Good
Lyon M et al. [33] 2015	Cohort	***	**	**	7	Good
Abd El Wahab A et al. [35] 2022	Cohort	***	**	**	7	Good
Martinez-Baena D et al. [36] 2013	Cohort	***	**	**	7	Good
Mir IS et al. [37] 2016	Cohort	***	**	**	7	Good
Parra-Membrives P et al. [38] 2017	Cohort	***	**	***	8	Good

<sup>a</sup>(1) representativeness of the exposed cohort; (2) selection of the non-exposed cohort; (3) ascertainment of exposure; (4) demonstration that outcome of interest was not present at the start of the study

<sup>b</sup>(1) comparability of cohorts based on design or analysis (maximum two stars)

<sup>c</sup>(1) assessment of outcome; (2) was follow-up long enough for outcomes to occur; (3) adequacy of follow-up of cohorts

<https://doi.org/10.1371/journal.pone.0280755.t002>

**Reoperation rate.** In the pooled analysis of nine studies ( $n = 677$ ), the reoperation rate between the biliary stenting and T-tube groups revealed no differences significantly [RR 0.59 (95% CI 0.25–1.42),  $p = 0.24$ ,  $I^2 = 0\%$ , random-effect modeling] (Fig 3D).

## Publication bias

We assessed the risk of bias using Funnel plot analysis for each synthesis. Our analysis illustrated a relatively symmetrical inverted plot for all syntheses. Thus, our Funnel plot analysis indicates no publication bias in this study (Fig 4A–4I).

## Discussion

Based on the included studies, T-tube insertion as biliary drainage after CBDE has been unpopular in the modern surgical era. Ten out of 16 studies were conducted before 2017. This trend is shifting towards the biliary stenting era in developed countries. A similar meta-analysis had been conducted by Yin *et al.* and Jiang *et al.*; nonetheless, our study did not discern the operative procedure (open or laparoscopic) because we aimed to solely analyze the safety and feasibility between T-tube and biliary stent insertion after CBDE [19, 20].

Table 3. Quality appraisal of studies included in the meta-analysis using the Jadad scale assessment.

Study	Random allocation	Concealment schemes	Blinding	Withdrawals and Drop-out	Total score	Interpretation
Gyalpo T et al. [28] 2018	2	0	1	1	4	Moderate quality
Mangla V et al. [34] 2012	2	2	2	1	7	High quality
Perez G et al. [39] 2005	1	1	1	0	3	Moderate quality
Redwan AA et al. [40] 2006	1	1	1	1	4	Moderate quality
Xu Y et al. [41] 2013	1	1	1	0	3	Moderate quality

Interpretation:






More than 4 points : High quality

3–4 points : Moderate quality

Less than 3 points : Low quality

<https://doi.org/10.1371/journal.pone.0280755.t003>







**Fig 2.** Forest plots of (A) bile leakage, (B) wound infection, (C) pancreatitis, (D) overall postoperative complications, and (E) operation time.

<https://doi.org/10.1371/journal.pone.0280755.g002>

The previous meta-analysis stated that T-tube insertion yielded higher biliary complications [19, 20]. On the other hand, our analytical synthesis demonstrated a similar outcome in terms of biliary leakage ( $n = 881$ ) and pancreatitis ( $n = 363$ ) with [RR 0.67 (95% CI 0.36–1.27),  $p = 0.22$ ,  $I^2 = 0\%$ , random-effect modeling] and [RR 0.55 (95% CI 0.13–2.28),  $p = 0.41$ ,  $I^2 = 41\%$ , random-effect modeling], respectively. These contradictory findings could be the limitation of the previous meta-analysis, which underwent the procedure laparoscopically. Visual limitation during a laparoscopic procedure when the CBD is fragile, edematous, and prone to bleeding might escalate higher complication rate [42]. Therefore, biliary complications will increase when the T-tube is inserted with the laparoscopic technique in the previous meta-analysis.

Another concern from the patient's point of view is having a 'tube' on their tummy skin for weeks or months. Hygiene, patient activities, and external exposures might raise the possibility of wound infection [43]. Based on our latest analysis, T-tube drainage did not increase the wound infection incidence compared to biliary stenting [RR 0.37 (95% CI 0.12–1.13),  $p = 0.08$ ,  $I^2 = 12\%$ , random-effect modeling]. Somehow, if we examine the overall postoperative complications (consisting of bile leakage, pancreatitis, wound infection, stricture, residual tone, bleeding, and bile duct injury), biliary stenting exhibited lower incidences [RR 0.43 (95% CI 0.23–0.80),  $p = 0.007$ ,  $I^2 = 63\%$ , random-effect modeling] compared to T-tube group. This analysis ensured that the complexity of T-tube insertion affected postoperative complications.


Regarding efficiency, our pooled analysis from 12 studies ( $n = 698$ ) showed operation time in biliary stenting as the drainage method after CBDE was shorter than the T-tube group. This finding is concurrent with the previous statement about the difficulty of T-tube insertion, and surgeons are prone to prefer biliary stenting as biliary drainage due to a shorter learning curve [44]. Shorter operation time and fewer complications mean a safer and preferable option for patients and operators.

Furthermore, we analyzed a blood loss volume that the previous meta-analysis had not conducted. This outcome is essential for patient safety and further complications. The pooled analysis from four studies ( $n = 177$ ) exhibited no difference in blood loss volume between both groups [SMD -0.45 (95% CI -1.11, 0.21),  $p = 0.18$ ,  $I^2 = 77\%$ , random-effect modeling].

Length of hospital stays and readmission rates are also crucial regarding potential nosocomial infections in the hospital [45, 46]. Our statistical analysis of hospital stay in 16 studies ( $n = 1,080$ ) showed a shorter duration of hospital stay in the biliary stent group [SMD -1.96 (95% CI -2.63, -1.29),  $p < 0.00001$ ,  $I^2 = 95\%$ , random-effect modeling] and the readmission rate analysis from four studies ( $n = 306$ ) was also significantly lower in the biliary stent group compared to the T-tube group [RR 0.39 (95% CI 0.15–0.97),  $p = 0.04$ ,  $I^2 = 0\%$ , random-effect modeling]. Our findings were equivalent to prior meta-analysis [19, 20].


Lastly, the reoperation rate did not significantly differ between both groups [RR 0.59 (95% CI 0.25–1.42),  $p = 0.24$ ,  $I^2 = 0\%$ , random-effect modeling]. Even though the overall postoperative complications were higher in the T-tube group, the reoperation rate did not differ statistically. This finding suggested that the postoperative complications could be managed with non-operative approaches.

We realize in today's era, a less invasive approach has been a favorable technique, and T-tube should be dropped off. We would like to emphasize to the physicians that biliary stent should be a preferred method due to its superiority in several aspects (shorter operation time,



**Fig 3.** Forest plots of (A) blood loss volume, (B) length of hospital stay, (C) readmission rate, and (D) reoperation rate.

<https://doi.org/10.1371/journal.pone.0280755.g003>



**Fig 4.** Funnel plots of (A) bile leakage, (B) wound infection, (C) pancreatitis, (D) overall postoperative complications, (E) operation time, (F) blood loss volume, (G) length of hospital stay, (H) readmission rate, and (I) reoperation rate.

<https://doi.org/10.1371/journal.pone.0280755.g004>

shorter hospital stay, and lower overall complications rate). Despite the advancement of the biliary stent in the last decade, these two candidates still can be compared. Our included studies in the last five years assessed both procedures, and T-tube remains in the modern era that gradually fades over time.

There are some considerations regarding our meta-analysis limitations. First, only five out of 16 studies are categorized as RCTs. Second, ten studies were conducted before 2017. More updated and high-quality RCTs are needed to improve further meta-analysis. Third, high heterogeneity among the included studies might affect the overall outcomes. The contributions of high heterogeneity derive from data characteristics, the definition of each outcome, and outcome parameters. Lastly, English-only studies were also limiting our meta-analysis to extend further included studies.

## Conclusions

Stenting as biliary drainage after CBDE was superior to T-tube insertion. A shorter operation time and hospital stay in biliary drainage resulted in a lower overall postoperative complication rate. Other influences, including the complexity and shorter learning curve, might also affect the superiority of biliary stenting. We recommend stenting over T-tube insertion as biliary drainage; even though, T-tube cannot be left out entirely in developing countries with limited healthcare facilities.

## Supporting information

**S1 Checklist. PRISMA 2020 checklist.**  
(PDF)

## Author Contributions

**Conceptualization:** Reno Rudiman, Ricarhdo Valentino Hanafi.

**Data curation:** Reno Rudiman, Ricarhdo Valentino Hanafi.

**Formal analysis:** Reno Rudiman, Ricarhdo Valentino Hanafi.

**Investigation:** Reno Rudiman, Ricarhdo Valentino Hanafi, Freda Halim.

**Methodology:** Reno Rudiman, Ricarhdo Valentino Hanafi, Freda Halim.

**Project administration:** Reno Rudiman.

**Resources:** Reno Rudiman.

**Software:** Reno Rudiman.

**Supervision:** Almawijaya.

**Validation:** Reno Rudiman, Ricarhdo Valentino Hanafi, Freda Halim.

**Visualization:** Reno Rudiman, Ricarhdo Valentino Hanafi, Freda Halim.

**Writing – original draft:** Reno Rudiman, Ricarhdo Valentino Hanafi, Almawijaya, Freda Halim.

**Writing – review & editing:** Reno Rudiman, Ricarhdo Valentino Hanafi, Almawijaya, Freda Halim.

## References

1. Gurusamy KS, Koti R, Davidson BR. T-tube drainage versus primary closure after laparoscopic common bile duct exploration. *Cochrane Database of Systematic Reviews* 2013; 6:CD005641. <https://doi.org/10.1002/14651858.CD005641.pub3> PMID: 23794201
2. Molvar C, Glaenger B. Choledocholithiasis: Evaluation, Treatment, and Outcomes. *Semin Intervent Radiol* 2016; 33:268–76. <https://doi.org/10.1055/s-0036-1592329> PMID: 27904245
3. Ahmed I, Pradhan C, Beckingham IJ, Brooks AJ, Rowlands BJ, Lobo DN. Is a T-tube Necessary after Common Bile Duct Exploration? *World J Surg* 2008; 32:1485–8. <https://doi.org/10.1007/s00268-008-9475-2> PMID: 18270647
4. Lee JS, Yoon YC. Laparoscopic common bile duct exploration using V-Loc suture with insertion of endobiliary stent. *Surg Endosc* 2016; 30:2530–4. <https://doi.org/10.1007/s00464-015-4518-x> PMID: 26310532
5. Baiu I, Hawn MT. Choledocholithiasis. *JAMA* 2018; 320:1506. <https://doi.org/10.1001/jama.2018.11812> PMID: 30304429
6. Hakuta R, Kawahata S, Kogure H, Nakai Y, Saito K, Saito T, et al. Endoscopic papillary large balloon dilation and endoscopic papillary balloon dilation both without sphincterotomy for removal of large bile duct stones: A propensity-matched analysis. *Digestive Endoscopy* 2019; 31:59–68. <https://doi.org/10.1111/den.13220> PMID: 29943385
7. Deaver J. The surgery of jaundice. *Ann Surg* 1925; 81:287–98. <https://doi.org/10.1097/00000658-192501010-00024> PMID: 17865173
8. Daldoul S, Moussi A, Zaouche A. T-tube drainage of the common bile duct choleperitoneum: Etiology and management. *J Visc Surg* 2012; 149:e172–8. <https://doi.org/10.1016/j.jvisc.2012.03.008> PMID: 22537812
9. Quaresima S, Balla A, Guerrieri M, Campagnacci R, Lezoche E, Paganini AM. A 23 year experience with laparoscopic common bile duct exploration. *HPB* 2017; 19:29–35. <https://doi.org/10.1016/j.hpb.2016.10.011> PMID: 27890483
10. Chen CC, Wu SD, Tian Y, Zeng XT, Siwo EA, Xian GZ. The Fading Role of T-Tube in Laparoscopic Choledochotomy: Primary Choledochorrhaphy and Over Pigtail J and Endonasobiliary Drainage Tubes. *Journal of Laparoendoscopic & Advanced Surgical Techniques* 2010; 20:807–11. <https://doi.org/10.1089/lap.2010.0075> PMID: 21029026
11. Gharaibeh KI, Heiss HA. Biliary leakage following T-tube removal. *Int Surg* 2000; 85:57–63. PMID: 10817434
12. Pfau PR, Pleskow DK, Banerjee S, Barth BA, Bhat YM, Desilets DJ, et al. Pancreatic and biliary stents. *Gastrointest Endosc* 2013; 77:319–27. <https://doi.org/10.1016/j.gie.2012.09.026> PMID: 23410693
13. Buxbaum JL, Abbas Fehmi SM, Sultan S, Fishman DS, Qumseyra BJ, Cortessis VK, et al. ASGE guideline on the role of endoscopy in the evaluation and management of choledocholithiasis. *Gastrointest Endosc* 2019; 89:1075–1105.e15. <https://doi.org/10.1016/j.gie.2018.10.001> PMID: 30979521
14. Garteiz Martínez D, Sánchez AW, López Acosta ME. Laparoscopic T-tube choledochotomy for biliary lithiasis. *JSLs* 2008; 12:326–31. PMID: 18765064
15. Wills VL, Gibson K, Karihaloo C, Jorgensen JO. Complications of biliary T-tubes after choledochotomy. *ANZ J Surg* 2002; 72:177–80. <https://doi.org/10.1046/j.1445-2197.2002.02308.x> PMID: 12071447
16. Zhu H, Wu L, Yuan R, Wang Y, Liao W, Lei J, et al. Learning curve for performing choledochotomy bile duct exploration with primary closure after laparoscopic cholecystectomy. *Surg Endosc* 2018; 32:4263–70. <https://doi.org/10.1007/s00464-018-6175-3> PMID: 29602995
17. Lou J, Zhao H, Chen W, Wang J. T tube sinus tract duodenal fistula: a rare complication of postoperative choledochoscopy for treating retained intrahepatic stones. *Surg Endosc* 2021; 35:5567–72. <https://doi.org/10.1007/s00464-020-08057-7> PMID: 33030589
18. Zhu QD, Tao CL, Zhou MT, Yu ZP, Shi HQ, Zhang QY. Primary closure versus T-tube drainage after common bile duct exploration for choledocholithiasis. *Langenbecks Arch Surg* 2011; 396:53–62. <https://doi.org/10.1007/s00423-010-0660-z> PMID: 20582601
19. Yin Z, Xu K, Sun J, Zhang J, Xiao Z, Wang J, et al. Is the End of the T-Tube Drainage Era in Laparoscopic Choledochotomy for Common Bile Duct Stones Is Coming? A Systematic Review and Meta-Analysis. *Ann Surg* 2013; 257:54–66. <https://doi.org/10.1097/SLA.0b013e318268314b> PMID: 23059495
20. Jiang C, Zhao X, Cheng S. T-Tube Use After Laparoscopic Common Bile Duct Exploration. *JSLs* 2019; 23:e2018.00077.
21. Margulis A v, Pladevall M, Riera-Guardia N, Varas-Lorenzo C, Hazell L, Berkman N, et al. Quality assessment of observational studies in a drug-safety systematic review, comparison of two tools: the

- Newcastle-Ottawa Scale and the RTI item bank. *Clin Epidemiol* 2014; 6:359–68. <https://doi.org/10.2147/CLEP.S66677> PMID: 25336990
22. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJM, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: Is blinding necessary? *Control Clin Trials* 1996; 17:1–12. [https://doi.org/10.1016/0197-2456\(95\)00134-4](https://doi.org/10.1016/0197-2456(95)00134-4) PMID: 8721797
  23. Tripepi G, Jager KJ, Dekker FW, Zoccali C. Stratification for Confounding—Part 1: The Mantel-Haenszel Formula. *Nephron Clin Pract* 2010; 116:c317–21. <https://doi.org/10.1159/000319590> PMID: 20664286
  24. Borenstein M, Hedges L v., Higgins JPT, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 2010; 1:97–111. <https://doi.org/10.1002/jrsm.12> PMID: 26061376
  25. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; 14:135. <https://doi.org/10.1186/1471-2288-14-135> PMID: 25524443
  26. Abdelkader AM, Zidan AM, Younis MT. Temporary CBD Stenting with a Nelaton Tube Is a More Practical and Safer Option Than T-Tube Drainage after Conventional CBD Exploration for Choledocholithiasis. *HPB Surgery* 2018; 2018:1–5. <https://doi.org/10.1155/2018/8035164> PMID: 30302070
  27. Dietrich A, Alvarez F, Resio N, Mazza O, de Santibañes E, Pekolj J, et al. Laparoscopic management of common bile duct stones: Transpapillary stenting or external biliary drainage? *Journal of the Society of Laparoendoscopic Surgeons* 2015; 18:1–5.
  28. Gyalpo T, Akram M, Dolma K, Azad TP. A Comparative Study of T-tube versus Ante grade Stenting of CBD after Open CBD Exploration for Choledocholithiasis. *Journal of Medical Science And clinical Research* 2018; 6:436–45.
  29. Hassan A, Mohammed A, Shawky MEG. Common bile duct exploration with transpapillary stenting versus T-tube drainage for management of irretrievable common bile duct stones. *Al-Azhar Assiut Medical Journal* 2017; 15:117–21.
  30. Isla AM, Griniatsos J, Karvounis E, Arbuckle JD. Advantages of laparoscopic stented choledochorrrhaphy over T-tube placement. *British Journal of Surgery* 2004; 91:862–6. <https://doi.org/10.1002/bjs.4571> PMID: 15227692
  31. Kwon SU, Choi IS, Moon JI, Ra YM, Lee SE, Choi WJ, et al. Comparison of Bile Drainage Methods after Laparoscopic CBD Exploration. *Korean J Hepatobiliary Pancreat Surg* 2011; 15:117–22. <https://doi.org/10.14701/kjhbps.2011.15.2.117> PMID: 26421026
  32. Xiao LK, Xiang JF, Wu K, Fu X, Zheng MY, Song XX, et al. The reasonable drainage option after laparoscopic common bile duct exploration for the treatment of choledocholithiasis. *Clin Res Hepatol Gastroenterol* 2018; 42:564–9. <https://doi.org/10.1016/j.clinre.2018.07.005> PMID: 30145281
  33. Lyon M, Menon S, Jain A, Kumar H. Use of biliary stent in laparoscopic common bile duct exploration. *Surg Endosc* 2015; 29:1094–8. <https://doi.org/10.1007/s00464-014-3797-y> PMID: 25249145
  34. Mangla V, Chander J, Vindal A, Lal P, Ramteke VK. A Randomized Trial Comparing the Use of Endobiliary Stent and T-Tube for Biliary Decompression After Laparoscopic Common Bile Duct Exploration. *Surg Laparosc Endosc Percutan Tech* 2012; 22:345–8. <https://doi.org/10.1097/SLE.0b013e31825b297d> PMID: 22874685
  35. Abd El Wahab A, Tawila M, Khalil O. Comparative study between primary common bile duct repair with internal stent insertion versus t-tube drainage after common bile duct exploration. *Al-Azhar International Medical Journal* 2022; 3:107–11.
  36. Martínez-Baena D, Parra-Membrives P, Díaz-Gómez D, Lorente-Herce JM. Laparoscopic common bile duct exploration and antegrade biliary stenting: Leaving behind the Kehr tube. *Revista Española de Enfermedades Digestivas* 2013; 105:125–30. <https://doi.org/10.4321/s1130-01082013000300002> PMID: 23735018
  37. Mir I, Hajini M, Rashid T, Sheikh V, Bhat S. Laparoscopic management of common bile duct stones: Stent Vs T-tube drainage. *Archives of International Surgery* 2016; 6:90–5.
  38. Parra-Membrives P, Martínez-Baena D, Lorente-Herce J, Jiménez-Riera G. Comparative Study of Three Bile Duct Closure Methods Following Laparoscopic Common Bile Duct Exploration for Choledocholithiasis. *Journal of Laparoendoscopic and Advanced Surgical Techniques* 2018; 28:145–51. <https://doi.org/10.1089/lap.2017.0433> PMID: 28976804
  39. Pérez G, Escalona A, Jarufe N, Ibáñez L, Viviani P, García C, et al. Prospective randomized study of T-tube versus biliary stent for common bile duct decompression after open choledocotomy. *World J Surg* 2005; 29:869–72. <https://doi.org/10.1007/s00268-005-7698-z> PMID: 15951939
  40. Redwan A, Omar M. Common bile duct clearance of stones by open surgery, laparoscopic surgery, and endoscopic approaches (comparative study). *The Egyptian Journal of Surgery* 2017; 36:76.



41. Xu Y, Dong C, Ma K, Long F, Jiang K, Shao P, et al. Spontaneously removed biliary stent drainage versus T-tube drainage after laparoscopic common bile duct exploration. *Medicine (United States)* 2016; 95. <https://doi.org/10.1097/MD.00000000000005011> PMID: 27684867
42. Mooney MJ, Deyo GA, O'Reilly MJ. T-tube placement during laparoscopic cholecystectomy. *Surg Endosc* 1992; 6:32–32. <https://doi.org/10.1007/BF00591184> PMID: 1344576
43. Asaduzzaman M, Mia MGM, Abdullah S, Islam MS, Hanif MA, Karim SR. Comparative Study between Primary Closure and T-tube Drainage after Open Choledochotomy. *Med Today* 2017; 29:15–8.
44. Kitagawa K, Mitoro A, Minami R, Nagamatsu S, Ozutsumi T, Fujinaga Y, et al. Efficacy of a dedicated plastic stent in endoscopic ultrasound-guided hepaticogastrostomy during the learning curve: cumulative multi-center experience. *Scand J Gastroenterol* 2022;1–8. <https://doi.org/10.1080/00365521.2022.2118557> PMID: 36073949
45. Jia H, Li L, Li W, Hou T, Ma H, Yang Y, et al. Impact of Healthcare-Associated Infections on Length of Stay: A Study in 68 Hospitals in China. *Biomed Res Int* 2019; 2019:1–7. <https://doi.org/10.1155/2019/2590563> PMID: 31119159
46. Stewart S, Robertson C, Pan J, Kennedy S, Haahr L, Manoukian S, et al. Impact of healthcare-associated infection on length of stay. *Journal of Hospital Infection* 2021; 114:23–31.