

A Sequence of Cauchy Sequences Conjectured to Converge to the Zeros of the Hardy Z Function

Stephen Crowley <stephencrowley214@gmail.com >

November 22, 2017

Abstract

A sequence of Cauchy sequences which conjecturally converge to the zeros of the Hardy Z function is constructed.

1 Preliminary Outline

1.1 Definitions

Let $\zeta(t)$ be the Riemann zeta function

$$\begin{aligned}\zeta(t) &= \sum_{n=1}^{\infty} n^{-s} & \forall \operatorname{Re}(s) > 1 \\ &= (1 - 2^{1-s}) \sum_{n=1}^{\infty} n^{-s} (-1)^{n-1} & \forall \operatorname{Re}(s) > 0\end{aligned}\tag{1}$$

and $\vartheta(t)$ be Riemann-Siegel vartheta function

$$\vartheta(t) = -\frac{i}{2} \left(\ln \Gamma \left(\frac{1}{4} + \frac{it}{2} \right) - \ln \Gamma \left(\frac{1}{4} - \frac{it}{2} \right) \right) - \frac{\ln(\pi)t}{2}\tag{2}$$

so that the Hardy Z function[2] can be defined by

$$Z(t) = e^{i\vartheta(t)} \zeta \left(\frac{1}{2} + it \right)\tag{3}$$

which is real-valued when t is real and satisfies the identity

$$\zeta(t) = e^{-i\vartheta(\frac{i}{2}-it)} Z \left(\frac{i}{2} - it \right)\tag{4}$$

where $\ln \Gamma(z)$ is the principal branch of the logarithm of the Γ function defined by

$$\ln \Gamma(z) = \ln(\Gamma(z)) = (z-1)! = \prod_{k=1}^{z-1} k \forall z \in \mathbb{R} > 0\tag{5}$$

which is analytically continued from the positive real axis when $z \in \mathbb{C}$ is complex. Each of the points $z \in \mathbb{Z} = \{0, -1, -2, \dots\}$ is a singularity and a branch point so that the union of the branch cuts is the negative real axis. On the branch cuts, the values of $\ln \Gamma(z)$ are determined by continuity from above. Let $S(t)$ denote the normalized argument of $\zeta(t)$ on the critical line

$$\begin{aligned}S(t) &= \pi^{-1} \arg \left(\zeta \left(\frac{1}{2} + it \right) \right) \\ &= -\frac{i}{2\pi} \left(\ln \zeta \left(\frac{1}{2} + it \right) - \ln \zeta \left(\frac{1}{2} - it \right) \right) \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0} \operatorname{Im} \left(\ln \zeta \left(\frac{1}{2} + it + \varepsilon \right) \right)\end{aligned}\tag{6}$$

1.2 Transcendental Equations Satisfied By The Nontrivial Riemann Zeros

Definition 1 The *critical line* is the line in the complex plane defined by $\text{Re}(t) = \frac{1}{2}$.

Definition 2 The *critical strip* is the strip in the complex plane defined by $0 < \text{Re}(t) < 1$.

Definition 3 The *exact equation* for the n -th zero of the Hardy Z function y_n is given by [1, Equation 20]

$$\vartheta(y_n) + S(y_n) = \left(n - \frac{3}{2}\right) \pi \quad (7)$$

where y_n enumerates the zeros of Z on the real line and the imaginary parts of the zeros of ζ on the critical line

$$Z(y_n) = 0 \text{ and } \zeta\left(\frac{1}{2} + iy_n\right) \forall n \in \mathbb{Z}^+ \quad (8)$$

where \mathbb{Z}^+ denotes the positive integers. [1, Equation 14]

By replacing the $\ln \Gamma$ function in (2) with Stirling's asymptotic expansion as in [1, Equation 13] we get

$$\tilde{\vartheta}(t) = \frac{t}{2} \ln\left(\frac{t}{2\pi e}\right) - \frac{\pi}{8} + O(t^{-1}) \quad (9)$$

and substitute $\vartheta(t)$ with $\tilde{\vartheta}(t)$ in Equation 7 which leads to

Definition 4 The *asymptotic equation* for the n -th zero of the Hardy Z function

$$\frac{t_n}{2\pi} \ln\left(\frac{t_n}{2\pi t}\right) + S(t_n) = n - \frac{11}{8} \quad (10)$$

[1, Equation 20]

Remark 1 The fact that the exact and asymptotic equations have two solutions when $n = 1$ can be understood by noting that Equations (7) and (10) are derived from the equation

$$n = \tilde{\vartheta}(t) + \frac{\pi}{8} - \frac{5}{8} + S(t) \quad (11)$$

which has a minimum in the interval $(-2, -1)$ and thus $n \geq -1$ so that, in order to follow the convention that the zeros are enumerated by the positive integers, the substitution $n \rightarrow n - 2$ is made in Equation (11) so that

$$n - 2 = \tilde{\vartheta}(t) + \frac{\pi}{8} - \frac{5}{8} + S(t) \quad (12)$$

[1, Equation 12]




Figure 1: The functions $\vartheta(y_n) + S(y_n) - (n - \frac{3}{2})\pi$ and $\tilde{\vartheta}(y_n) + S(y_n) - (n - \frac{3}{2})\pi$ for $n = 1, 2$ with the zeros at y_1 and y_2 marked with vertical lines.

Theorem 1 *If the limit*

$$\lim_{\delta \rightarrow 0^+} \arg \left(\zeta \left(\frac{1}{2} + \delta + it \right) \right) \quad (13)$$

exists and is well-defined $\forall t$ then the left-hand side of Equation (10) is well-defined $\forall t$, and due to monotonicity, there must be a unique solution for every $n \in \mathbb{Z}^+$. [1, II.A]

Corollary 1 *The number of solutions of Equation (10) over the interval $[0, t]$ is given by*

$$N_0(t) = \frac{t}{2\pi} \ln \left(\frac{t}{2\pi e} \right) + \frac{7}{8} + S(t) + O(t^{-1}) \quad (14)$$

which counts the number of zeros **on the critical line**.

Conjecture 1 *(The Riemann hypothesis) All solutions t of the equation*

$$\zeta(t) = 0 \quad (15)$$

besides the trivial solutions $t = -2n$ with $n \in \mathbb{Z}^+$ have real-part $\frac{1}{2}$, that is, $\text{Re}(t) = \frac{1}{2}$ when $\zeta(t) = 0$ and $t \neq -2n$.

Definition 5 *The Riemann-von-Mangoldt formula makes use of Cauchy's argument principle to count the number of zeros **inside the critical strip** $0 < \text{Im}(\rho_n) < t$ where $\zeta(\sigma + i\rho_n)$ with $0 < \sigma < 1$*

$$N(t) = \frac{t}{2\pi} \ln \left(\frac{t}{2\pi e} \right) + \frac{7}{8} + S(t) + O(t^{-1}) \quad (16)$$

and this definition does not depend on the Riemann hypothesis(Conjecture 1). This equation has exactly the same form as the asymptotic Equation 10. [1, Equation 15]

Lemma 1 *If the exact Equation (7) has a unique solution for each $n \in \mathbb{Z}^+$ then Conjecture 1, the Riemann hypothesis, follows.*

Proof If the exact equation has a unique solution for each n , then the zeros obtained from its solutions on the **critical line** can be counted since they are enumerated by the integer n , leading to the counting function $N_0(t)$ in Equation (14). The number of solutions obtained on the **critical line** would saturate counting function of the number of solutions on the **critical strip** so that $N(t) = N_0(t)$ and thus all of the non-trivial zeros of ζ would be enumerated in this manner. If there are zeros off of the critical line, or zeros with multiplicity $m \geq 2$, then the exact Equation (7) would fail to capture all the zeros on the critical strip which would mean $N_0(t) < N(t)$. [1, IX] \square

Corollary 2 *The Riemann hypothesis(RH) is not necessarily false if the exact Equation (7) does not have a unique solution for every n , since the solutions could still be on the critical line but not necessarily simple, that is, a root on the critical line could have multiplicity $m \geq 2$ and the RH would still be true.*

Corollary 3 *The Riemann hypothesis is true and all of the zeros on the critical line are simple if the exact Equation (7) has a unique solution for each $n \in \mathbb{Z}^+$. [1, IX]*

1.3 Iterated Function Systems

Definition 6 *A fixed-point α of a function $f(x)$ is a value α such that*

$$f(\alpha) = \alpha \quad (17)$$

[7, 3.]

Definition 7 *The multiplier $\lambda_f(\alpha)$ of a fixed point α of a map $f(x)$ is equal to the derivative $\dot{f}(\alpha)$ of the map evaluated at the point α which is the first term in the Taylor expansion at that point*

$$\lambda_f(\alpha) = \dot{f}(\alpha) \quad (18)$$

If $|\lambda_f(\alpha)| < 1$ then α is said to be an attractive fixed-point of $f(x)$. If $|\lambda_f(\alpha)| = 1$ then α is an indifferent fixed-point of $f(t)$ also known as a neutral fixed-point, and if $|\lambda_f(\alpha)| > 1$ then α is a repelling fixed-point of $f(t)$. When $|\lambda_f(\alpha)| = 0$ the fixed-point α is said to be a superattractive fixed-point of $f(t)$ [7, 3.]

Definition 8 *Let*

$$Y_{n,m}(t) = \begin{cases} t & m = 0 \\ t + h_{n,m} \cos(\pi n) \tanh\left(\frac{Z(Y_{n,m-1}(t))}{|\Omega(t)| \prod_{k=1}^{n-1} \tanh(Y_{n,m-1}(t) - y_k)}\right) & m \geq 1 \end{cases}$$

denote the m -th iterate of the n -th iteration function corresponding to the n -th zero of the Hardy Z function where

$$\Omega(t) = \begin{cases} 1 & t = e \\ e^{\frac{3}{4} \sqrt{\frac{\log(t)}{\log(\log(t))}}} & t \neq e \end{cases} \quad (19)$$

is a lower bound for the running maximum of $|Z(s)|$

$$\max_{0 \leq s \leq t} |Z(s)| > \Omega(t) \forall t \geq 45.590 \dots \quad (20)$$

ensuring that

$$\frac{|Z(t)|}{\Omega(t)} > 0 \forall t \geq 45.590 \dots \quad (21)$$

which normalizes the range of $Z(t)$ which is known to grow in both maximum and average value as $t \rightarrow \infty$ and $h_{n,m}$ is factor which influences the rate of convergence

$$h_{n,m} = \begin{cases} 1 & m \leq 2 \\ \frac{h_{n,m-1}}{2} & \begin{aligned} & \text{sign}(\Delta Y_{n,m-2}(t)) = \text{sign}(\Delta Y_{n,m-1}(t)) \\ & \text{sign}(\Delta Y_{n,m-2}(t)) \neq \text{sign}(\Delta Y_{n,m-1}(t)) \end{aligned} \end{cases} \quad (22)$$

where

$$\Delta Y_{n,m}(t) = Y_{n,m}(t) - Y_{n,m-1}(t) \quad (23)$$

is the 1-st difference of the m -th iterate for the n -th zero. [6, Theorem 3.2.3]

Lemma 2 The roots of $Z(t)$ are fixed-points of $Y_{n,m}(t) \forall n, m \in \mathbb{Z}^+$.

Proof If $Z(t) = 0$ then $\tanh\left(\frac{Z(t)}{|\Omega(t)| \prod_{k=1}^{n-1} \tanh(t-y_k)}\right) = \tanh\left(\frac{0}{|\Omega(t)| \prod_{k=1}^{n-1} \tanh(t-y_k)}\right) = \tanh(0) = 0$ so that $Y_n(t) = t + \cos(\pi n)0 = t + 0 = t$ when $Z(t) = 0$. \square

Theorem 2 $Y_{n,m}(t)$ has indifferent fixed-points at each point y_k where $k = 1 \dots n-1$

Proof The product in the denominator $\prod_{k=1}^{n-1} \tanh(t-y_k) \rightarrow 0$ smoothly as t approaches any $y_k \in \bigcup_{k=1}^{n-1} y_k$ since $\tanh(0) = 0$ and \tanh is a smooth function. When any element of the product is zero the value of the product is zero regardless of the values of any other elements of the product. Since $\frac{1}{s} \rightarrow \infty$ as $s \rightarrow 0$ and $\tanh(|x|) \rightarrow 1$ as $|x| \rightarrow \infty$ we have $\tanh(\infty) = 1$ and $\tanh(-\infty) = -1$ so that $Y_n(t) = t + \cos(\pi n) \forall t \in \bigcup_{k=1}^{n-1} y_k$. Since $Y_n(t) = t \pm 1 \forall t \in \bigcup_{k=1}^{n-1} y_k$ when n is an integer, we see that $\frac{d}{dt} Y_n(t) = \frac{d}{dt} (t \pm 1) = 1$ so that the multiplier $\lambda_{Y_n(t)} = \left| \frac{d}{dt} Y_n(t) \right| = 1 \forall t \in \bigcup_{k=1}^{n-1} y_k$. \square

Theorem 3 $Y_{n,m}(t)$ has indifferent fixed points at each trivial zero $-\frac{i}{2}(-4n-1)$ where $Z\left(-\frac{i}{2}(-4n-1)\right) = 0 \forall n \in \mathbb{Z}^+$.

Proof Since $\frac{d}{dt}(f(t)+g(t)) = \dot{f}(t)g(t) + f(t)\dot{g}(t)$ and $Z(t) = e^{i\vartheta(t)} \zeta\left(\frac{1}{2} + it\right)$ it suffices to show that $\lim_{t \rightarrow -\frac{i}{2}(-4n-1)} \left| \frac{d}{dt} e^{i\vartheta(t)} \right| = \infty$. Since $\frac{d}{dt} e^{i\vartheta(t)} = i \left(\frac{i\Psi\left(\frac{1}{4} - \frac{it}{2}\right)}{2} - \frac{i\Psi\left(\frac{1}{4} + \frac{it}{2}\right)}{2} - \frac{\ln(\pi)}{2} \right) e^{i\vartheta(t)}$ we only have to check that $\lim_{t \rightarrow -\frac{i}{2}(-4n-1)} \left| \Psi\left(\frac{1}{4} + \frac{it}{2}\right) \right| = \lim_{t \rightarrow -\frac{i}{2}(-4n-1)} \left| \Psi\left(\frac{1}{4} - \frac{it}{2}\right) \right| = \infty$ which is true since $\Psi(t)$ has poles at $t = 1 - n \forall n \in \mathbb{Z}^+$ where $\Psi(t) = \frac{\Gamma'(t)}{\Gamma(t)}$ and $\Gamma(t)$ has poles at $t = 1 - n \forall n \in \mathbb{Z}^+$. Since $|\tanh(\infty)| = 1$ the multiplier is equal to 1 at each $-\frac{i}{2}(-4n-1)$. \square

Proposition 1 When n is an odd number, $Y_n(t)$ has attractive fixed-points at the odd-numbered roots $y_{2k-1} \forall 2k-1 \geq n$ and repulsive fixed-points at the even-numbered roots $y_{2k} \forall 2k \geq n$.

Proposition 2 When n is an even number, $Y_n(t)$ has attractive fixed-points at the even-numbered roots $y_{2k} \forall 2k \geq n$ and repulsive fixed-points at the odd-numbered roots $y_{2k-1} \forall 2k-1 \geq n$.




Figure 2: The functions which are subtracted or added to t to get $Y_1(t), Y_2(t), Y_3(t), Y_4(t)$. When n is odd $\cos(\pi n) = -1$ so that the value is subtracted from t , when n is even $\cos(2\pi) = 1$ so it is added. It is plain to see that the curves $\tanh\left(\frac{Z(t)}{\Omega(t) \prod_{k=1}^{n-1} \tanh(t-y_k)}\right)$ do not cross the zero axis for any $t < y_n$

Remark 2 The function $h_{n,m}$ is defined to be 1 when $1 \leq m \leq 2$. If $\text{sign}(\Delta Y_{n,m-1}(t)) \neq \text{sign}(\Delta Y_{n,m}(t))$ then $h_{n,m+1} = \frac{h_{n,m}}{2}$ so that the convergence rate is halved when the sign of the difference between successive iterates changes, indicating that it jumped across the root. This prevents the sequence generated by the iteration from getting stuck in an artificial 2-cycle and jumping back and forth across the root with equal magnitude indefinitely when implementing this method with finite-precision arithmetic on a digital computer. Without this successive relaxation, the iterates still converge in theory however the number of iterations required could be several million or higher, while still having the difficulty of possibly getting stuck in a 2-cycle in computer implementations.

Theorem 4 The Lipschitz constant M of the map $Y_{n,m}(t) < 1 \forall t > e$ therefore $Y_{n,m}(t)$ is a contraction mapping

$$|Y_{n,m}(t) - Y_{n,m}(s)| \leq M|t - s| \quad (24)$$

Proof The Lipschitz constant of a continuous differentiable function $f(x)$ is equal to the maximum absolute value of its derivative

$$M = \sup_x \left| \frac{d}{dx} f(x) \right| \quad (25)$$




Figure 3: Multiplier of the maps $Y_1(t)$ and $Y_3(t)$

The derivative of $t - \tanh(t)$ is $\tanh(t)^2$. Since the maximum absolute value of $\tanh(t)$ is 1 then the maximum value of its square is also 1. Since $\Omega(t) > 1$ and $h_{n,m} \leq 1$ the derivative $\frac{d}{dt}Y_{n,m}(t)$ can never have an absolute value ≥ 1 since that would require $\left| \tanh\left(\frac{Z(t)}{|\Omega(t)| \prod_{k=1}^{n-1} \tanh(t-y_k)}\right) \right| = 1$ which is only possible if $Z(t) = \pm\infty$ which is only the case when $t = \pm\frac{i}{2}$ which corresponds to the pole at $\zeta(1)$. Since $Z(t) \in \mathbb{R}$ when $t \in \mathbb{R}$ it can never be the case that $Z(t) = \infty$ so that $\left| \frac{d}{dt}Y_{n,m}(t) \right| \neq 1 \forall t \in \mathbb{R}$ and the Lipschitz constant M is strictly less than 1. \square

Proposition 3 *The limit*

$$y_n = \lim_{m \rightarrow \infty} Y_{n,m}(s_n) \quad (26)$$

where

$$s_n = \begin{cases} 14 & n = 1 \\ 21 & n = 2 \\ \frac{y_{n-1} + y_{n-2}}{2} & n \geq 3 \end{cases} \quad (27)$$

exists and is equal to the n -th zero of the Hardy Z function for all integer $n \in \mathbb{Z}^+$. That is, $Y_{n,m}(z_n)$ forms a Cauchy sequence, due to the contraction mapping property proved in Theorem 4 whose elements are indexed by m converging to the n -th root y_n where the n -th starting point is defined to be half-way between the $(n-2)$ -th and the $(n-1)$ -th root y_n when $n > 2$ and equal to a point close to the first known zero at 14.134... when $n = 1$ and a point close to the 2nd zero at 21.022... when $n = 2$

Remark 3 The mid-way point between the nearest neighbors to the left of y_n is used as the starting point for the iteration since any point less than y_n and greater than e is within the immediate basin of attraction of y_n . The precise location of any roots y_p where $p < n$ cannot be used as a starting point since the map $Y_{p,m}(t)$ is a non-expansive mapping with Lipschitz constant precisely equal to 1 when $t \in \bigcup_{k=1}^{p-1} y_k$ so that the hyperbolic tangent has an argument of infinity resulting in a value of 1. Trajectories are neither attracted or repelled to any point $\bigcup_{k=1}^{n-1} y_k$ under the action of the map $Y_{n,m}(t)$ however, trajectories started precisely on any point $t \in \bigcup_{k=1}^{n-1} y_k$ will never attain a value other than t since any y_k is a fixed-point of $Y_{n,m}(t)$.

Note 1 The truth of Proposition 3 has been verified computationally up to $n = 800,000$ with a computer program which implements the methods described here using the arbitrary precision complex ball arithmetic library arblib[3] and compares the results against the tables published by Andrew Odlyzko[5].

Theorem 5 *The Cauchy sequence $\lim_{m \rightarrow \infty} Y_{n,m}(s_n)$ will never converge to any y_k where $k < n$.*

Proof All y_k are indifferent fixed-points of $Y_{n,m}(t)$ and the trajectories generated by $Y_{n,m}(s_n)$ are never started from a point y_k since $s_n \notin \bigcup_{k=1}^{n-1} y_k$ and the only way $Y_{n,m}(t)$ would "converge" to an indifferent fixed-point is if it was started precisely on one, and s_n is by definition equal to the mid-point between successive y_n . \square

Theorem 6 *The Cauchy sequence $Y_{n,m}(s_n)$ will never converge to any $y_{n+2k-1} \forall k \in \mathbb{Z}^+$ if Proposition 1 is true.*

Proof If Proposition 1 is true then y_{n+2k-1} are repelling fixed-points for $Y_{n,m}(t)$. \square

Note 2 If Propositions 1 and 2 are true then $Y_{n,m}(s_n)$ will never converge to y_q with q odd and n even nor to y_r with r even and n odd. It suffices to prove that $Y_{n,m}(s_n) < y_{n+1} \forall n, m \in \mathbb{Z}^+$ which would mean that $Y_{n,m}(s_n)$ can never jump across the repelling fixed-point at y_{n+1} to land on any of the attractive fixed-points in $\bigcup_{k=1}^{\infty} y_{n+2k}$

Lemma 3 *Let*

$$Y_{n,m}^+(t) = \begin{cases} t & m = 0 \\ t + h_{n,m} \cos(\pi n) \tanh\left(\frac{Z(Y_{n,m-1}^+(t))}{|\Omega(t)| \prod_{k=1}^{n-1} \tanh(Y_{n,m-1}^+(t) - y_k)}\right) & m \geq 1 \end{cases} \quad (28)$$

$$Y_{n,m}^-(t) = \begin{cases} t & m = 0 \\ t - h_{n,m} \cos(\pi n) \tanh\left(\frac{Z(Y_{n,m-1}^-(t))}{|\Omega(t)| \prod_{k=1}^{n-1} \tanh(Y_{n,m-1}^-(t) - y_k)}\right) & m \geq 1 \end{cases} \quad (29)$$

and

$$z_n = \min\left(\lim_{m \rightarrow \infty} Y_{n,m}^+(t), \lim_{m \rightarrow \infty} Y_{n,m}^-(t)\right) \quad (30)$$

which must exist because there is known to be an infinity of zeros on the critical line.

Proof The only way z_n would not exist is if all the roots y_k were indifferent fixed-points $\forall k > n$ but that is impossible since there are no indifferent fixed-points of $Y_{n,m}(t)$ because for a fixed-point y_k to be indifferent would require $\tanh\left(\frac{Z(Y_{n,m-1}(t))}{|\Omega(t)|}\right) = 1$ which is only possible if $|Z(Y_{n,m-1}(t))| = \infty$ for some $m \in \mathbb{Z}^+$ and the Z function only takes on the value ∞ when $t = -\frac{i}{2}$ which corresponds to the pole at $\zeta(1)$ since $\frac{1}{2} + i\left(-\frac{i}{2}\right) = 1$. \square

Definition 9 *The multiplicity $m_f(t)$ of a root α is a root $f(\alpha) = 0$ such that its Taylor expansion about the point α has the form*

$$f(t) = c(t - \alpha)^{m_f(t)} + (\text{higher order terms} \dots) \quad (31)$$

where $c \neq 0$ and $m \geq 1$. The multiplicity of a root t is related to the multiplier $\lambda_f(t)$ through the formula

$$m_f(t) = \frac{1}{1 - \lambda_{N_f}(t)} \quad (32)$$

where

$$\lambda_{N_f}(t) = \frac{f(t)\ddot{f}(t)}{\dot{f}(t)} \quad (33)$$

is the first derivative of the Newton map of $f(t)$

$$N_f(t) = t - \frac{f(t)}{\dot{f}(t)} \quad (34)$$

Lemma 4 (Milnor's Lemma) *Every **simple root** of $f(t)$ is a super-attractive fixed-point of $N_f(t)$ since a superattractive fixed-point is one such that its multiplier $\lambda_f(t) = 0$ so that its multiplicity is*

$$m_f(t) = \frac{1}{1 - \lambda_{N_f}(t)} = \frac{1}{1 - 0} = \frac{1}{1} = 1 \quad (35)$$

See [4, p.52]

Proof Let α be a root $Z(\alpha) = 0$ then the multiplier of its Newton map is $\lambda_{N_Z}(\alpha) = \lambda_f(\alpha) = \frac{Z(\alpha)\ddot{Z}(\alpha)}{\dot{Z}(\alpha)} = 0$ since $Z(\alpha) = 0$ the entire expression $\frac{Z(\alpha)\ddot{Z}(\alpha)}{\dot{Z}(\alpha)}$ is equal to 0 since due to the ordering of operations the value of $\dot{Z}(t)$ or $\ddot{Z}(t)$ is never required to be known in order to know the value of $\lambda_{N_Z}(t)$ when $Z(t) = 0$. If any term in the product is 0 then the entire product takes the value 0. The multiplicity is related to the multiplier by $m_Z(t) = \frac{1}{1 - \lambda_{N_Z}(t)} = 1$ and therefore simple. Since $m_Z(t) = \frac{1}{1 - \lambda_{N_Z}(t)} \forall \lambda_{N_Z}(t) \neq 1$ then it is known that $\lambda_{N_Z}(t) = 0$ when $Z(t) = 0$ therefore the point α is a superattractive fixed-point corresponding to a simple zero at α . Since we now know that $m_Z = (\alpha)$ and therefore the zero at $Z(\alpha) = 0$ is simple, we therefore know that the denominator $\dot{Z}(t)$ of the multiplier $\lambda_{N_Z}(t)$ cannot vanish so that $\dot{Z}(\alpha) \neq 0$ since that would imply that α is not a simple root, which would be a contradiction to the already established fact that $m_Z(\alpha) = 1$ when $Z(\alpha) = 0$. \square

Conjecture 2 *Let*

$$c_n(\varepsilon) = \frac{Z(\max_{t \in [0, y_n]} \{Y_{n+1,1}(t) \geq t\} + \varepsilon) - Z(\min_{t \in [y_n, \infty]} \{Y_{n+1,1}(t) \leq t\} - \varepsilon)}{2\varepsilon + \max_{t \in [0, y_n]} \{Y_{n+1,1}(t) \geq t\} - \min_{t \in [y_n, \infty]} \{Y_{n+1,1}(t) \leq t\}} \quad (36)$$

denote the Lipschitz constant in Formula 38 then it is always possible to choose a small enough positive ε such that $0 < c_n(\varepsilon) < 1$.

2 Appendix

Lemma 5 *The Banach Fixed-Point Theorem*

If $f(x)$ is a continuous function defined on $[a, b]$ and

$$f(x) \in [a, b] \forall x \in [a, b] \quad (37)$$

and there exists some constant $0 < c < 1$ such that

$$\frac{|f(x) - f(y)|}{x - y} \leq c \quad (38)$$

then $f(x)$ has a unique fixed-point $x \in [a, b]$ and the sequence $f(x_0), f(f(x_0)), f(f(f(x_0))), \dots$ converges to the unique fixed-point of $f(x)$ in the interval $[a, b]$.

References

- [1] Guilherme França and André LeClair. Transcendental equations satisfied by the individual zeros of riemann zeta, dirichlet and modular l-functions. *Communications in Number Theory and Physics*, 2015.
- [2] A. Ivić. *The Theory of Hardy's Z-Function*. Cambridge Tracts in Mathematics. Cambridge University Press, 2013.
- [3] F. Johansson. Arb: a C library for ball arithmetic. *ACM Communications in Computer Algebra*, 47(4):166–169, 2013.
- [4] John Willard Milnor. *Dynamics in one complex variable*, volume 160. Springer, 2006.
- [5] Andrew Odlyzko. Tables of zeros of the riemann zeta function. http://www.dtc.umn.edu/~odlyZko/zeta_tables/.
- [6] K Ramachandra. *Lectures on the Mean-value and Omega-theorems for the Riemann Zeta-function*. Lectures on mathematics and physics. Tata Institute of Fundamental Research, 1995.
- [7] Hans Rådström. On the iteration of analytic functions. *Mathematica Scandinavica*, pages 85–92, 1953.