Leak-free data preprocessing for short-horizon PM2.5 forecasting on the Beijing PM2.5 dataset

Vetriselvan R, Hrishi Dongre, Pampana Sree Adithya, Ishan Goyal, Vedant Madne
Team T024, Section A - AML/DL Project 19: Air quality forecasting via gradient boosting and LSTM models

Abstract—We describe the preprocessing pipeline that turns the UCI Beijing PM2.5 hourly series into model-ready data for gradient boosting (Phase 1) and recurrent models (Phase 2). An audit of the raw file shows a complete hourly grid with missing values only in the target, a right-skewed target, and three weather columns stored as running totals rather than instantaneous values. The pipeline decodes these totals, keeps the target label untouched while building a causal input copy with a missingness flag, defines expanding-window chronological folds with a 24-hour embargo, and fits all scaling statistics on training dates only. Every guarantee is checked by automated tests.

I. Introduction

The project forecasts PM2.5 concentration a few hours ahead from past pollution and weather, and compares an XGBoost baseline with an LSTM or temporal CNN under rolling chronological backtesting. Forecast comparisons are only meaningful if no information from the evaluation period reaches training, including through filling, scaling or the split itself [4], [5]. This report covers the data side: what the raw data contains, which problems it has, and how each step avoids look-ahead leakage.

II. Dataset

The dataset [1] holds hourly PM2.5 measured at the US Embassy in Beijing together with meteorology from Beijing Capital International Airport, from 2010-01-01 00:00 to 2014-12-31 23:00 (43 824 rows). It was introduced in [2]. Table I lists the variables after renaming.

TABLE I
Variables after renaming

Name	Raw	Meaning (unit)
pm25	pm2.5	PM2.5 concentration ($\mu g\,m^{-3}$), target
dewp	DEWP	dew point (°C)
temp	TEMP	temperature (°C)
pres	PRES	pressure (hPa)
cbwd	cbwd	combined wind direction: NE, NW, SE, cv
ws_cum	Iws	cumulated wind speed (m/s)
snow_cum	Is	cumulated hours of snow
rain_cum	Ir	cumulated hours of rain

III. Data audit

- Time grid. Timestamps form a complete, unique hourly grid, so no resampling is needed. The time zone is not stated in the source.




Fig. 1. Missing pm25 hours per month. Gaps concentrate in 2010-2012.

- Missing values. Only pm25 is missing: 2067 hours (4.72%) in 214 gaps. Of these, 112 gaps last one hour, 143 last at most three hours, and 20 exceed 24 hours. The longest gap is 155 hours from 2010-09-21 05:00. Missingness falls from 669, 728 and 489 hours in 2010-2012 to 82 and 99 in 2013-2014 (Fig. 1).
- Skew. pm25 has median 72, 99th percentile 420 and maximum 994, with skewness 1.80; log1p reduces it to −0.30 (Fig. 3).
- Extremes. The two highest values, 994 at 2012-01-23 01:00 and 980 at 2010-02-14 01:00, fall in the first hours of the Lunar New Year; the next highest is 886 on 2013-01-12. Neighbouring hours rise and fall with them, so we treat the extremes as real episodes and keep them.
- Cumulative columns. Iws increases in every hour that cbwd stays the same and restarts when cbwd changes. Is and Ir count consecutive snow and rain hours (98.4% and 97.1% of non-zero values equal the previous value plus one).

IV. Preprocessing pipeline

A. Schema and grid validation

The loader checks the exact column list, builds a timestamp index from year, month, day and hour, and stops with an error if the hourly grid has holes or duplicates or if any weather value is missing. Later steps rely on these conditions, so the pipeline fails rather than silently repairing them.




Fig. 2. First three days of 2010. Top: raw Iws, a running total that restarts at each direction change (vertical lines; long runs labelled). Bottom: hourly speed recovered with (1), in m/s.

B. Decoding cumulative weather variables

With d_t the wind direction and I_t the Iws value at hour t, the hourly wind speed is

$$ws_t = \begin{cases} I_t & d_t \neq d_{t-1} \\ I_t - I_{t-1} & d_t = d_{t-1} \end{cases} \tag{1}$$

The decoded series lies in $[0.45, 20.12]$ m/s with mean 2.94; calm-variable (cv) hours average 0.99 m/s, consistent with the label. Fig. 2 shows the running total and the recovered hourly speed. The running total is kept as ws_cum because it also encodes how long the wind has blown from one direction. Snow and rain become binary flags (snow, rain = 1 when the counter is positive) next to the original counters.

C. Wind direction encoding

cbwd has four levels and is one-hot encoded in a fixed order (wd_ne, wd_nw, wd_se, wd_cv), so column positions do not depend on which levels appear in a fold. Unknown levels raise an error.

D. Missing target values

The label pm25 is never filled; a missing label means that hour is excluded from training loss and from evaluation. A flag pm25_miss records the original missingness. A separate input column pm25_in carries the last observed value forward for at most $L = 3$ hours:

$$\tilde{y}_t = y_{t-k}, \quad k = \min\{j \geq 0 : y_{t-j} \text{ observed}\}, \quad k \leq L, \tag{2}$$

and stays missing otherwise. Linear interpolation was rejected because it fills a gap using the first value after it, which leaks future information into inputs near the forecast origin. Forward filling recovers 386 input hours; 1681 remain missing and are left for the model stage (XGBoost handles them natively; windows for the LSTM can be dropped or masked).




Fig. 3. Distribution of pm25 before and after log1p.




Fig. 4. Expanding-window folds. The 24-hour embargo before each evaluation year is too short to see at this scale.

E. Chronological backtest folds

Following rolling-origin evaluation [4], [3], fold k evaluates on calendar year Y_k and trains on all hours in $[t_0,\ s_k - G)$, where s_k is 1 January of Y_k and $G = 24$ hours is an embargo. With G at least the forecast horizon, no training target can fall inside the evaluation year. Whole years are used so that every fold sees a full seasonal cycle. 2014 is held out as the final test year (Fig. 4, Table II).

F. Leak-free scaling

For each fold a scaler is fitted on that fold's training rows only and saved to disk:

$$z = \frac{g(x) - \mu_{\text{train}}}{\max(\sigma_{\text{train}}, \epsilon)}, \tag{3}$$

where g is log1p for the skewed columns (pm25, pm25_in, ws, ws_cum, snow_cum, rain_cum, all with skewness above 1) and the identity otherwise. Tree models can ignore it; recurrent models use it, and predictions are mapped back to $\mu g\,m^{-3}$ with the inverse transform before computing MAE, RMSE and R^2.

V. Implementation and verification

The pipeline is a Python package (src/aqf/data) driven by one configuration file (configs/data.yaml) and run with python -m aqf.data.pipeline. It writes the cleaned hourly frame (data/interim/hourly.parquet), the fold boundaries (data/processed/folds.json), one scaler per fold (data/processed/scalers) and the audit numbers used

TABLE II
Fold sizes (hours)

Fold	Train until	Evaluate	Train	Labelled eval
cv1	2010-12-30 23:00	2011	8736	8032
cv2	2011-12-30 23:00	2012	17 496	8295
cv3	2012-12-30 23:00	2013	26 280	8678
test	2013-12-30 23:00	2014	35 040	8661

in this report (reports/preprocessing/qa.json). Nine tests check that: the grid is complete and hourly; only the target is missing; summing decoded speeds within each direction run rebuilds Iws exactly; the one-hot columns are exclusive; the label is unchanged and the flag matches it; filling at hour t is identical whether or not later hours exist (causality); folds never overlap and respect the embargo; scaler statistics do not change when evaluation rows are altered; and the scaler round-trips through disk and its inverse.

VI. Next steps

Feature engineering (lags, rolling means and calendar terms, built after sorting by time), the persistence and seasonal-naive baselines, and the XGBoost model build on these outputs. The audit suggests a 24-hour seasonal-naive baseline rather than a weekly one, since PM2.5 autocorrelation is 0.40 at lag 24 but 0.02 at lag 168. The forecast horizon and the AQI standard still have to be fixed and stated.

References

[1] S. Chen, "Beijing PM2.5," UCI Machine Learning Repository, 2015, doi: 10.24432/C5JS49.

[2] X. Liang, T. Zou, B. Guo, S. Li, H. Zhang, S. Zhang, H. Huang, and S. Chen, "Assessing Beijing's PM2.5 pollution: severity, weather impact, APEC and winter heating," Proc. Royal Society A, 2015.

[3] R. J. Hyndman and G. Athanasopoulos, Forecasting: Principles and Practice, 3rd ed. Melbourne, Australia: OTexts, 2021.

[4] L. J. Tashman, "Out-of-sample tests of forecasting accuracy: an analysis and review," Int. J. Forecasting, vol. 16, no. 4, pp. 437–450, 2000.

[5] C. Bergmeir and J. M. Benítez, "On the use of cross-validation for time series predictor evaluation," Information Sciences, vol. 191, pp. 192–213, 2012.