

Design of experiments: Lab 3

1.1. The patients are a blocking variable and the gas is the treatment. We only have $r = 1$, so we'll fit an additive model:

$$y_{ijk} = \mu + \beta_i + \tau_j + \varepsilon_{ijk}, \quad \varepsilon_{ijk} \stackrel{\text{ind}}{\sim} N(0, \sigma^2)$$

with sum-to-zero restrictions

$$\sum_{i=1}^7 \beta_i = \sum_{j=1}^4 \tau_j = 0.$$

Here's the ANOVA table:


```
library(tidyverse)
gasos = read.csv2("http://vicpena.github.io/doe/lab3/Gases.csv")
gasos$Gas = factor(gasos$Gas)
gasos$Sujeto = factor(gasos$Sujeto)
options(contrasts = c("contr.sum", "contr.poly"))
mod_add = aov(Valor ~ Sujeto+Gas, data = gasos)
summary(mod_add)
```

	##	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
Sujeto	##	6	1471772	245295	270.61	< 2e-16 ***						
Gas	##	3	44827	14942	16.48	2.11e-05 ***						
Residuals	##	18	16316	906								
---	##											
Signif. codes:	##	0	'***'	0.001	'**'	0.01	'*'	0.05	'.'	0.1	' '	1

Both the block and the treatment are significant.

Let's take a look at the residual plots:


```
par(mfrow = c(1,2))
plot(mod_add, 1:2)
```



The residual plots look fine.

Let's take a look at the effect plots

```
library(emmeans)
emmip(mod_add, Sujeto ~ Gas)
```



There are obvious differences between subjects. It seems that gas C might be significantly worse than the others (the response is distance walked in 12 minutes). We can compare the gases with TukeyHSD:

```
TukeyHSD(mod_add, which = "Gas")
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = Valor ~ Sujeto + Gas, data = gasos)
##
## $Gas
```

```
##           diff           lwr           upr           p adj
## B-A    45.00000    -0.4840367    90.48404    0.0530703
## C-A   -61.57143   -107.0554653   -16.08739    0.0061872
## D-A    24.57143   -20.9126082    70.05547    0.4429649
## C-B  -106.57143  -152.0554653   -61.08739    0.0000178
## D-B   -20.42857   -65.9126082    25.05547    0.5929753
## D-C    86.14286    40.6588204   131.62689    0.0002338
```


The p -values for the tests comparing gas C to the others are significant, confirming our initial intuition.

1.2. The rats are blocks and the zones are treatments. We only have one replicate ($r = 1$), so we fit an additive model with both variables. The p -values are all significant. The residuals look fine, with the exception of observation 11, which seems to be badly predicted by the model (large residual).

```
rates = read.csv2("http://vicpena.github.io/doe/lab3/Rates.csv")
rates$Sujeto = factor(rates$Sujeto)
rates$Zona = factor(rates$Zona)
mod_add = aov(Cobre ~ Sujeto+Zona, data = rates)
summary(mod_add)
```

```
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Sujeto      7  703.3   100.48    3.938 0.00678 **
## Zona       3  565.9   188.63    7.393 0.00146 **
## Residuals   21  535.8    25.51
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

par(mfrow = c(1, 2))
plot(mod_add, 1:2)
```



Since we're interested in comparing zones, let's run TukeyHSD:

```
TukeyHSD(mod_add, which = "Zona")

##      Tukey multiple comparisons of means
##      95% family-wise confidence level
##
## Fit: aov(formula = Cobre ~ Sujeto + Zona, data = rates)
```

```
##
## $Zona
##      diff      lwr      upr      p adj
## Z1-N 10.7125   3.672793 17.752207 0.0019145
## Z2-N  9.3000   2.260293 16.339707 0.0069970
## Z3-N  4.6750  -2.364707 11.714707 0.2786427
## Z2-Z1 -1.4125  -8.452207  5.627207 0.9429572
## Z3-Z1 -6.0375 -13.077207  1.002207 0.1098481
## Z3-Z2 -4.6250 -11.664707  2.414707 0.2872316
```

There are significant differences between Z1 and N and also between Z2 and N.

1.3. We'd put the loaves randomly to avoid systematic biases (for example, some parts of the oven might be hotter than other). The batches are a block effect and the recipes are the treatment. This is another complete block design with $r = 1$ – we'll fit an additive model. There are significant batch and recipe effects. The residuals look fine (not including them for concreteness).

```
pa = read.csv2("http://vicpena.github.io/doe/lab3/Pan.csv")
pa$Receta = factor(pa$Receta)
pa$Hornada = factor(pa$Hornada)
mod_add = aov(Densidad ~ Receta + Hornada, data = pa)
summary(mod_add)
```

```
##      Df Sum Sq Mean Sq F value Pr(>F)
## Receta      2 0.08657  0.04329   8.137 0.0118 *
## Hornada      4 0.09884  0.02471   4.645 0.0312 *
## Residuals    8 0.04256  0.00532
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Let's compare the recipes with TukeyHSD:

```
TukeyHSD(mod_add, which = "Receta")

##      Tukey multiple comparisons of means
##      95% family-wise confidence level
##
## Fit: aov(formula = Densidad ~ Receta + Hornada, data = pa)
##
## $Receta
##      diff      lwr      upr      p adj
## B-A -0.088 -0.2198146  0.04381462 0.1983163
## C-A -0.186 -0.3178146 -0.05418538 0.0093756
## C-B -0.098 -0.2298146  0.03381462 0.1460142
```

There are significant differences between recipes A and C.

1.4. Now we have $r = 2$, so we can fit a model with an interaction and see if we need it

```
options(contrasts = c("contr.sum", "contr.poly"))
aigua = read.csv2("http://vicpena.github.io/doe/lab3/Aigua.csv")
mod_inter = aov(Reduccio ~ Accio*Densitat, data = aigua)
summary(mod_inter)
```

```
##      Df Sum Sq Mean Sq F value    Pr(>F)
## Accio      2 182.36    91.18   32.242 1.49e-05 ***
## Densitat    3 260.32    86.77   30.685 6.55e-06 ***
## Accio:Densitat 6  30.39     5.07    1.791   0.184
## Residuals   12  33.94     2.83
```

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

It turns out that the interaction isn't significant, so we go ahead and fit an additive model:

```
mod_add = aov(Reducchio ~ Accio + Densitat, data = aigua)
summary(mod_add)
```

```
##              Df Sum Sq Mean Sq F value    Pr(>F)
## Accio         2  182.36    91.18    25.51 5.58e-06 ***
## Densitat      3  260.32    86.77    24.28 1.50e-06 ***
## Residuals    18   64.33     3.57
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Our final model is

$$y_{ijk} = \mu + \alpha_i + \tau_j + \varepsilon_{ijk}, \quad \varepsilon_{ijk} \stackrel{\text{iid}}{\sim} N(0, \sigma^2),$$

where τ_i represents the effect of the “action” and τ_j the population density. The model has sum-to-zero restrictions on the effects, as usual. The hypothesis tests are


$$H_{0,\alpha} : \alpha_i = 0 \text{ for all } i, \quad H_{1,\alpha} : \alpha_i \neq 0 \text{ for at least one } i,$$

and

$$H_{0,\tau} : \tau_j = 0 \text{ for all } j, \quad H_{1,\tau} : \tau_j \neq 0 \text{ for at least one } j.$$


Here are the residual plots, which look fine:

```
par(mfrow = c(1, 2))
plot(mod_add, which = c(1,2))
```



And here's the effects plot:

```
emmip(mod_add, Accio ~ Densitat)
```



It seems that action S (a subsidy for changing old equipment) works best. This can be confirmed with TukeyHSD. If we want to find intervals for the actions, we can use `emmeans`:

```
emmeans(mod_add, ~ Accio)
```

```
## Accio emmean SE df lower.CL upper.CL
## C 7.46 0.668 18 6.06 8.87
## P 8.47 0.668 18 7.07 9.88
## S 13.75 0.668 18 12.35 15.15
##
## Results are averaged over the levels of: Densitat
## Confidence level used: 0.95
```

The 95% confidence interval for S goes from 12.35 to 15.15.

Finally, we can find the estimated mean for action S and density A with `emmeans`, which is 19.15:

```
emmeans(mod_add, ~ Accio + Densitat)
```

```
## Accio Densitat emmean SE df lower.CL upper.CL
## C A 12.87 0.945 18 10.88 14.85
## P A 13.88 0.945 18 11.89 15.87
## S A 19.15 0.945 18 17.17 21.14
## C B 7.28 0.945 18 5.30 9.27
## P B 8.30 0.945 18 6.31 10.28
## S B 13.57 0.945 18 11.58 15.56
## C C 4.35 0.945 18 2.36 6.34
## P C 5.36 0.945 18 3.38 7.35
## S C 10.64 0.945 18 8.65 12.62
## C D 5.35 0.945 18 3.36 7.34
## P D 6.36 0.945 18 4.38 8.35
## S D 11.64 0.945 18 9.65 13.62
##
## Confidence level used: 0.95
```