



Projected Entangled Pair States (PEPS)

Victor Vanthilt, Niccolò Laurora, Michael Shatnev
Ghent University, Università degli Studi di Milano, Kharkiv Institute of Physics and Technology


I. DEFINITION

Projected Entangled Pair States (PEPS) are a class of tensor network states that efficiently parametrise quantum states with finite entanglement. They are a generalization of Matrix Product States (MPS).

- Given a physical system composed of local sites with site-Hilbertspaces $\mathcal{H}_i \equiv \mathbb{C}^{d_i}$ situated on a graph with edges $E = \{e_{i,j}\}$ and vertices $V = \{v_i\}$.
- For each vertex v_i , and for each edge $e_{i,j}$ connected to v_i (and v_j), associate an ancilla degree of freedom $a_{i,j} \in \mathcal{H}_{i,j} \equiv \mathbb{C}^{d_{i,j}}$.
- Maximally entangle ancillae $a_{i,j}$ and $a_{j,i}$ associated with each edge $e_{i,j}$:

$$|\Phi_{i,j}\rangle = \sum_{l=1}^{d_{i,j}} |l\rangle \otimes |l\rangle \quad (1)$$

- Apply a linear map $\mathcal{P}_i : \bigotimes_{j_i} \mathcal{H}_{a_{i,j}} \rightarrow \mathcal{H}_i$ to the ancillae of site i , to obtain the projected entangled pair states (PEPS) $|\Psi\rangle = \bigotimes_{e \in E} \mathcal{P}_e |\Phi_e\rangle$.
- The final PEPS is a tensor network state that has the same connectivity as the original graph and that lives in the total Hilbert space $\mathcal{H} = \bigotimes_i \mathcal{H}_i$.



[Cirac et. al. 2021]

II. ENTANGLEMENT AREA LAW

Entanglement Area Law

The entanglement entropy of a region \mathcal{A} of quantum state with finite (local) entanglement scales as $\partial\mathcal{A}$, the boundary of \mathcal{A} .

$$S_{\mathcal{A}} \sim \partial\mathcal{A} \quad (2)$$

This is in contrast with the volume law most states follow.

Given the Schmidt decomposition of a state $|\Psi\rangle$ across a bipartition of the system into the “In” system \mathcal{A} and the “Out” system \mathcal{B} (where $\sum_i \lambda_i^2 = 1$):

$$|\Psi\rangle = \sum_i \lambda_i |I_i\rangle \otimes |O_i\rangle \quad (3)$$


The entanglement entropy of the region \mathcal{A} is given by:

$$S_{\mathcal{A}} = - \sum_i \lambda_i^2 \log(\lambda_i^2) \quad (4)$$

The maximal value this can take is when all the $\lambda_i = \frac{1}{N_S}$, with N_S the Schmidt rank. For finitely entangled systems, $S_{\mathcal{A}}$ is thus bounded by the finite Schmidt rank N_S .

$$S_{\mathcal{A}} \leq \log(N_S) \quad (5)$$

It turns out that PEPS satisfy an **area law** for their entanglement entropy **by construction**. This property makes PEPS an efficient representation of quantum states with finite (local) entanglement.



The entanglement entropy of the shaded area \mathcal{A} is the sum of the entanglement entropy across the cut virtual bonds, which by construction all have a finite bond dimension (and Schmidt rank) χ .

The entanglement entropy of a single cut is bounded by $\log(\chi)$. The total entanglement entropy of the region \mathcal{A} is thus proportional to the number of cut virtual bonds times $\log(\chi)$.

$$S_{\mathcal{A}} \sim \log(\chi) \cdot \partial\mathcal{A} \quad (6)$$

III. PARENT HAMILTONIANS

Parent Hamiltonians for PEPS

Every injective PEPS is the unique ground state of a *local, frustration free* parent Hamiltonian.

Consider the same shaded region \mathcal{A} as before. The support space $\mathcal{S}_{\mathcal{A}}$ of the reduced density matrix $\rho_{\mathcal{A}}$ grows as $\partial\mathcal{A}$ while the Hilbert space $\mathcal{H}_{\mathcal{A}}$ grows as \mathcal{A} .

Therefore we can always find a sufficiently large region such that $\rho_{\mathcal{A}}$ has a non-trivial kernel $\ker(\rho_{\mathcal{A}}) = \mathcal{H}_{\mathcal{A}} \setminus \mathcal{S}_{\mathcal{A}}$.

It is thus the groundstate of every Hamiltonian h with:

- $h \geq 0$
- $\ker(h) = \mathcal{S}_{\mathcal{A}}$

since by construction $\text{tr}(\rho_{\mathcal{A}} h) = 0$.

The Hamiltonian can be extended to the entire lattice by assigning a region \mathcal{R}_v to every vertex v of the lattice and defining

$$H = \sum_v h_v \otimes \mathbb{1} \quad (7)$$

The Hamiltonian is *local* since it is the sum of local terms h_v acting on the region \mathcal{R}_v , and it is *frustration free* since every term h_v is minimized by the PEPS.

[Perez-Garcia et. al. 2007]

IV. COMPUTATION OF LOCAL EXPECTATION VALUES

COMPUTATION OF EXPECTATION VALUES?????????

V. COMPUTATIONAL COMPLEXITY OF PEPS CONTRACTIONS

How Complex are these peps really? It can't be that NP-Hard.