


What is the rate of the reaction?

May 30, 2017 by Veerendra — Leave a Comment

What is the rate of the reaction?


- **Rate of reaction** measures the speed at which the reactants are converted to the products in a chemical reaction.



- For a reaction that occurs **rapidly**, the rate of reaction is **high**. Conversely, for a reaction that occurs **slowly**, the rate of reaction is low.
- The **time taken** for a **fast** reaction is **short**, whereas the time taken for a slow reaction is **long**.
- Hence, the **rate** of a particular reaction is **inversely proportional** to the **time taken** for the reaction.

Rate of reaction $\propto \frac{1}{\text{time taken}}$

- Different chemical reactions occur at different rates. Some examples are illustrated in Table.



Kindergarten

4th Grade

8th Grade

1st Grade

5th Grade

X² Algebra 1 & 2

2nd Grade

6th Grade

Geometry

3rd Grade

7th Grade

Precalculus

Type of reaction	Fast reaction	Slow reaction
Reaction involving liberation of a gas	Bubbles of carbon dioxide gas liberate rapidly when sodium carbonate powder reacts with dilute hydrochloric acid. $\text{Na}_2\text{CO}_3(\text{s}) + 2\text{HCl}(\text{aq}) \rightarrow 2\text{NaCl}(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$	In photosynthesis, carbon dioxide reacts with water very slowly in the presence of sunlight and chlorophyll to produce glucose and oxygen gas. $6\text{CO}_2(\text{g}) + 6\text{H}_2\text{O}(\text{l}) \rightarrow \text{C}_6\text{H}_{12}\text{O}_6(\text{s}) + 6\text{O}_2(\text{g})$
Precipitation reaction	When silver nitrate solution is added to sodium chloride solution, a white precipitate of silver chloride is formed immediately. $\text{AgNO}_3(\text{aq}) + \text{NaCl}(\text{aq}) \rightarrow \text{AgCl}(\text{s}) + \text{NaNO}_3(\text{aq})$	When dilute hydrochloric acid is added to sodium thiosulphate solution, a yellow precipitate of sulphur appears only after a few seconds. $\text{Na}_2\text{S}_2\text{O}_3(\text{aq}) + 2\text{HCl}(\text{aq}) \rightarrow 2\text{NaCl}(\text{aq}) + \text{S}(\text{s}) + \text{SO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
Heating a metal in air	When a small piece of potassium is heated in air, it burns rapidly to form a white solid of potassium oxide. $4\text{K}(\text{s}) + \text{O}_2(\text{g}) \rightarrow 2\text{K}_2\text{O}(\text{s})$	When a small piece of copper is heated in air, it reacts slowly with oxygen in the air to form a black solid of copper(II) oxide. $2\text{Cu}(\text{s}) + \text{O}_2(\text{g}) \rightarrow 2\text{CuO}(\text{s})$

People also ask

SEE ALSO

Online IT Courses

2020 Accredited Online Colleges

Accredited Schools Online

Online Graduate Degrees

Download Free PDF Books

10 Best Masters Degrees

Affordable Online Universities

Free Math Lessons

- How do you calculate the reaction rate?
- What factors affect the rate of a reaction?
- How does the surface area affect the rate of reaction?
- Explain the effect of concentration on the rate of reaction?

CBSE CLASS 10
SAMPLE PAPERS BUNDLE



BIG SALE

40 Min Revision Book worth ₹95 for FREE

₹ 614 399*

* Includes All Subjects

DOWNLOAD NOW

RS Aggarwal Solutions

RS Aggarwal Solutions Class 10

RS Aggarwal Solutions Class 9

RS Aggarwal Solutions Class 8

RS Aggarwal Solutions Class 7

RS Aggarwal Solutions Class 6

ICSE Solutions

Selina ICSE Solutions

Concise Mathematics Class 10 ICSE Solutions

Concise Physics Class 10 ICSE Solutions

Concise Chemistry Class 10 ICSE Solutions

Concise Biology Class 10 ICSE Solutions

Concise Mathematics Class 9 ICSE Solutions

Concise Physics Class 9 ICSE Solutions

Concise Chemistry Class 9 ICSE Solutions

Concise Biology Class 9 ICSE Solutions

ML Aggarwal Solutions

ML Aggarwal Class 10 Solutions

ML Aggarwal Class 9 Solutions

ML Aggarwal Class 8 Solutions

ML Aggarwal Class 7 Solutions

ML Aggarwal Class 6 Solutions

HSSLive Plus One

HSSLive Plus Two

Kerala SSLC



RECENT POSTS

Kurukshetra University Distance Education Admissions 2020 Dates, Course & Fees

Resignation Letter | How To Write a Letter Resignation To Company, Format & Samples

Osmania University Distance Education Admission 2020 Courses & Fee Details

What does the rate of reaction measure?

Observable changes for measuring the rate of reaction:

1. When a reaction occurs, **two obvious changes** that occur are:

- the **quantity of a reactant decreases** with **time**
- the **quantity of a product increases** with **time**

2. The **quantity** of a reactant/product can be the

- number of moles** of a substance
- mass** of a solid
- volume** of a gas
- concentration** of a solution

3. If the **changes** of any of these quantities are **visible** and **measurable** during a reaction, then it can be used to **measure** the **rate** of that reaction.

4. Suitable measurable visible changes in a chemical reaction are:

- volume** of a gas liberated
- formation** of a precipitate
- changes in the **mass** during a reaction
- colour** changes
- changes in the **electrical conductivity** of the solution
- temperature** changes
- pressure** changes
- changes in **concentration** of the solution of a reactant
- pH** changes


5. **One** of these measurable visible changes can be selected as a **suitable quantity** to determine the rate of a particular reaction.

The **changes** in this **selected quantity** can be measured by carrying out **an experiment** and the results are then **analysed** to determine the rate of that reaction.

6. **Definition: Rate of reaction** is defined as the **change** in a **selected quantity** during a reaction **per unit time** whereby the selected quantity can be any of the measurable visible changes in the reaction.

7. **Two examples to illustrate the meaning of rate of reaction.**

Reaction between magnesium and dilute sulphuric acid



- In the reaction between dilute sulphuric acid and a magnesium ribbon, the following **two changes** are observed:
 - The **mass of magnesium** (the reactant) **decreases** with **time**.
 - The **volume of hydrogen gas** (the product) **increases** with **time**.
- Hence, the **rate of reaction** between dilute sulphuric acid and magnesium can be **determined** by measuring the **change in the mass** of magnesium or the **volume** of hydrogen gas per **unit time**.
Quantitatively,

NCVT MIS ITI Result Portal


NCVT ITI MIS Admit Card 2020 | Download NCVT ITI MIS Hall Ticket from Here

Madras University Distance Education Admission 2020 Dates, Courses & Fees Details

Anna University Distance Education Admission Application Form 2020 Eligibility, Courses, Dates

CG ITI Admission 2020 | Eligibility, Dates, Application Form, Selection Process

ITI Admission Process 2020 | State Wise ITI Admissions, Application Form, Eligibility



- When excess aqueous ethanedioic acid is added to acidified potassium manganate(VII) solution, the **purple** colour of the solution **slowly decolourises** at room temperature.
- By **measuring the time taken** for the purple colour to decolourise, the rate of reaction can be determined.
- Rate of reaction is **inversely proportional** to the time taken for the purple colour to decolourise.

- Rate of reaction is reflected by the value of 1/time taken. The **larger** the value time taken of **1/time taken**, the **higher** the **rate** of reaction.
- Table shows the units for the rate of reaction measured in different ways.

Change in a selected quantity per unit time	Units for the rate of reaction
Change in mass per unit time	g s^{-1} or g min^{-1}
Change in volume of a gas liberated per unit time	$\text{cm}^3 \text{ s}^{-1}$ or $\text{cm}^3 \text{ min}^{-1}$
Change in concentration of a reactant per unit time	$\text{mol dm}^{-3} \text{ s}^{-1}$ or $\text{mol dm}^{-3} \text{ min}^{-1}$
Change in number of moles of a reactant per unit time	mol s^{-1} or mol min^{-1}

Rate of Reaction Experiment

Aim: To compare the rates of a few reactions.

Materials: Marble chips, 2 mol dm⁻³ hydrochloric acid, 0.1 mol dm⁻³ sodium thiosulphate solution, 1 mol dm⁻³ lead(II) nitrate solution.

Apparatus: 50 cm³ beakers, test tubes.

Procedure:

- 5 cm³ of 2 mol dm⁻³ hydrochloric acid is poured into each of the three test tubes on a rack.
- The test tubes are labelled I to III respectively.
- One piece of marble chip is added into test tube I.
- About 2 cm³ of 0.1 mol dm⁻³ sodium thiosulphate solution is poured into test tube II and the mixture is shaken well.
- About 2 cm³ of 1 mol dm⁻³ lead(II) nitrate solution is poured into test tube III and the mixture is shaken well.
- The changes are observed carefully. The rates of reactions in the three test tubes are compared.

Observations:

Test tube	Reactants	Observation
I	Marble chip and hydrochloric acid	Bubbles of a colourless gas are liberated rapidly, that is, effervescence occurs rapidly.
II	Sodium thiosulphate solution and hydrochloric acid	A yellow precipitate appears only after about 12 seconds.
III	Lead(II) nitrate solution and	A white precipitate is formed immediately.

1. The reaction between lead(II) nitrate solution and hydrochloric acid is very fast.
2. The reaction between the marble chip and hydrochloric acid is moderately fast.

Excellent Support G

Ad

Cheapest SSL Certifi

Web Security Solutions I

Truehost Cloud

Shop Now

3. The reaction between sodium thiosulphate solution and hydrochloric acid is slow.

Discussion:

1. The chemical equation for the reaction in

2. The rate of reaction in ascending order is: rate in test tube II < rate in test tube I < rate in test tube III
3. The observable change that can be used to compare the rate of reaction in
- (a) test tube I is the time taken for the effervescence to stop completely.
- (b) test tube II is the time taken for the appearance of a yellow precipitate.
- (c) test tube III is the time taken for the appearance of a white precipitate.

Conclusion:

1. The rate of reaction between sodium thiosulphate solution and hydrochloric acid is the lowest.
2. The rate of reaction between the marble chip and hydrochloric acid is moderately high.
3. The rate of reaction between lead(II) nitrate solution and hydrochloric acid is the highest.

Filed Under: **Chemistry**

Tagged With: **compare the rates of a few reactions, rate of reaction, rate of reaction calculation, rate of reaction concentration, rate of reaction examples, rate of reaction experiment, rate of reaction factors, rate of reaction temperature, rate of reaction units, Units for the rate of reaction, What does the rate of reaction measure?, What is the rate of the reaction?**

Leave a Reply

Your email address will not be published. Required fields are marked *

Comment

Name *

Email *

Website

POST COMMENT

RS Aggarwal Solutions

RS Aggarwal Solutions Class 10

RS Aggarwal Solutions Class 9

RS Aggarwal Solutions Class 8

RS Aggarwal Solutions Class 7

RS Aggarwal Solutions Class 6

ICSE Solutions

Selina ICSE Solutions

ML Aggarwal Solutions

HSSLive Plus One

HSSLive Plus Two

Kerala SSLC

Disclaimer

Privacy Policy