

Lecture 20: Numerical Analysis (UMA011)

Dr. Meenu Rani

School of Mathematics
TIET, Patiala
Punjab-India

System of linear equations:

Example:

Suppose that

$$2x_1 + x_2 + 3x_3 = 1$$

$$4x_1 + 6x_2 + 8x_3 = 5$$

$$6x_1 + \alpha x_2 + 10x_3 = 5,$$

with $|\alpha| < 10$. For which of the following values of α will there be no row interchange required when solving this system using scaled partial pivoting?

a.) $\alpha = 6,$

b.) $\alpha = 9,$

c.) $\alpha = -3.$

System of linear equations

Solution:

$$[A:b] = \begin{matrix} E_1 \\ E_2 \\ E_3 \end{matrix} \left[\begin{array}{ccc|c} 2 & 1 & 3 & 1 \\ 4 & 6 & 8 & 5 \\ 6 & \alpha & 10 & 5 \end{array} \right]$$

To find the scaled factors

$$s_1 = \max \{ |2|, |1|, |3| \} = 3$$

$$s_2 = 8$$

$$s_3 = \max \{ |6|, |\alpha|, |10| \} = 10$$

$$\max \left\{ \frac{|a_{11}|}{s_1}, \frac{|a_{21}|}{s_2}, \frac{|a_{31}|}{s_3} \right\} = \max \left\{ \frac{2}{3}, \frac{4}{8}, \frac{6}{10} \right\} = \frac{2}{3} = \frac{|a_{11}|}{s_1}$$

$$E_2 \rightarrow E_2 - 2E_1, \quad E_3 \rightarrow E_3 - 3E_1$$

$$[A:b] = \begin{matrix} E_1 \\ E_2 \\ E_3 \end{matrix} \left[\begin{array}{ccc|c} 2 & 1 & 3 & 1 \\ 0 & 4 & 2 & 3 \\ 0 & \alpha-3 & 1 & 2 \end{array} \right]$$

$$\begin{aligned} \max \left\{ \frac{|a_{22}|}{s_2}, \frac{|a_{32}|}{s_3} \right\} &= \max \left\{ \frac{4}{8}, \frac{|\alpha-3|}{10} \right\} \\ &= \max \left\{ \frac{1}{2}, \frac{|\alpha-3|}{10} \right\} \end{aligned}$$

If there is no row interchange, then

$$\frac{|\alpha-3|}{10} < \frac{1}{2}$$

$$|\alpha-3| < 5$$

$$-5 < \alpha-3 < 5$$

$$-5 < \alpha - 3 < 5$$

$$-2 < \alpha < 8.$$

a) when $\alpha = 6$, then no row interchange required.

b) when $\alpha = 9$, then row interchange required.

c) when $\alpha = -3$, then row interchange required.

System of linear equations

LU Factorization:

Procedure

1st step.

Reduce A into U

$$E_3 \rightarrow E_3 - \left(\frac{a_{32}}{a_{22}} \right) E_2$$

$$E_n \rightarrow E_n - \left(\frac{a_{n2}}{a_{22}} \right) E_2$$

$$AX = b$$

$$X = ?$$

$$A = \begin{matrix} E_1 \\ E_2 \\ \vdots \\ E_n \end{matrix} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ - & - & - & - \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

If $a_{11} \neq 0$

$$E_2 \rightarrow E_2 - \left(\frac{a_{21}}{a_{11}} \right) E_1$$

$$E_3 \rightarrow E_3 - \frac{a_{31}}{a_{11}} E_1$$

$$E_n \rightarrow E_n - \frac{a_{n1}}{a_{11}} E_1$$

$$E_j \rightarrow E_j - \frac{a_{j1}}{a_{11}} E_1$$

$$= L * U$$

Lower
triangular
matrix

upper
triangular
matrix

$$\begin{bmatrix} \text{Lower triangular matrix} \end{bmatrix} * \begin{bmatrix} \text{Upper triangular matrix} \end{bmatrix}$$

L U

$$E_j \rightarrow E_j - \left(\frac{a_{ji}}{a_{ii}} \right) E_i, \quad i=1, 2, \dots, n-1$$

$$j=i+1, i+2, \dots, n.$$

$$A \sim \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \vdots & 0 & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix} = U$$

2nd step. construct L

$$L = \begin{bmatrix} l_{11} & 0 & 0 & \cdots & 0 \\ l_{21} & l_{22} & \cdots & \cdots & 0 \\ l_{31} & l_{32} & l_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ l_{n1} & \cdots & \cdots & \cdots & l_{nn} \end{bmatrix}$$

$$L = \begin{bmatrix} 1 & 0 & - & 0 \\ l_{21} & 1 & 0 & - & 0 \\ l_{31} & l_{32} & 1 & & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ l_{n1} & l_{n2} & - & - & - & 1 \end{bmatrix}$$

$$l_{21} = \frac{a_{21}}{a_{11}}$$

$$l_{32} = \frac{a_{32}}{a_{22}}$$

Hy get all the entries
of L

$$l_{31} = \frac{a_{31}}{a_{11}}$$

$$l_{n2} = \frac{a_{n2}}{a_{22}}$$

In general, $l_{ji} = \frac{a_{ji}}{a_{ii}}$

$$l_{n1} = \frac{a_{n1}}{a_{11}}$$

How can we get the entries of L

$$L * U = \begin{bmatrix} l_{11} & 0 & \dots & 0 \\ l_{21} & l_{22} & & 0 \\ l_{31} & l_{32} & l_{33} & 0 \\ \vdots & \vdots & \vdots & \vdots \\ l_{n1} & \dots & \dots & l_{nn} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & & a_{2n} \\ & & \dots & \\ 0 & & & a_{nn} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \hline \hline a_{n1} & \dots & \dots & a_{nn} \end{bmatrix} = A$$

$$l_{11} a_{11} = a_{11}$$

$$l_{11} = 1$$

$$l_{22} = 1$$

$$l_{21} a_{11} = a_{21}$$

$$l_{21} = \frac{a_{21}}{a_{11}}$$

$$l_{31} = \frac{a_{31}}{a_{11}}$$

$$l_{32} = \frac{a_{32}}{a_{22}}$$

- - -

- - -

3rd step

$$A = \check{L} \check{U}$$

$$AX = b$$

$$\Rightarrow (\check{L}\check{U})X = b$$

$$\check{L}(\check{U}X) = b$$

$$\text{Put } \underset{n \times n}{\check{U}} \underset{n \times 1}{X} = \underset{n \times 1}{Y} \text{ (say)}$$

$$\check{L}Y = b$$

$$\begin{bmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \hline l_{m1} & l_{m2} & \dots & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

where

$$Y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}_{n \times 1}$$

by using forward sub.

$$1 * y_1 = b_1 \checkmark$$

$$l_{21} * y_1 + y_2 = b_2$$

$$\underline{\hspace{1cm}} \\ y_n = \dots$$

$$(AB)C \\ = A(BC)$$

4th step

$$UX = y$$

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$$

Use backward sub.

System of linear equations

Example:

Determine the LU factorization for matrix A in the linear system

$$Ax = b, \text{ where } A = \begin{bmatrix} 1 & 1 & 0 & 3 \\ 2 & 1 & -1 & 1 \\ 3 & -1 & -1 & 2 \\ -1 & 2 & 3 & -1 \end{bmatrix}, B = \begin{bmatrix} 1 \\ 1 \\ -3 \\ 4 \end{bmatrix}. \text{ Then}$$

use the factorization to solve the system.

System of linear equations

Solution:

$$A = \begin{matrix} E_1 \\ E_2 \\ E_3 \\ E_4 \end{matrix} \begin{bmatrix} 1 & 1 & 0 & 3 \\ 2 & 1 & -1 & 1 \\ 3 & -1 & -1 & 2 \\ -1 & 2 & 3 & -1 \end{bmatrix}$$

$$E_2 \rightarrow E_2 - \textcircled{2}E_1, \quad E_3 \rightarrow E_3 - \textcircled{3}E_1, \quad E_4 \rightarrow E_4 + E_1$$

$$A \sim \begin{matrix} E_1 \\ E_2 \\ E_3 \\ E_4 \end{matrix} \begin{bmatrix} 1 & 1 & 0 & 3 \\ 0 & -1 & -1 & -5 \\ 0 & -4 & -1 & -7 \\ 0 & \textcircled{3} & 3 & 2 \end{bmatrix}$$

$$E_3 \rightarrow E_3 - \textcircled{4}E_2, \quad E_4 \rightarrow E_4 + \textcircled{3}E_2$$

$$A \sim \begin{bmatrix} 1 & 1 & 0 & 3 \\ 0 & -1 & -1 & -5 \\ 0 & 0 & 3 & 13 \\ 0 & 0 & 0 & -13 \end{bmatrix} = U$$

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 3 & 4 & 1 & 0 \\ -1 & -3 & 0 & 1 \end{bmatrix}$$

$$AX = b$$

$$(LU)X = b$$

$$L(UX) = b$$

$$\text{Let } UX = Y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix} \text{ (say)}$$

$$\Rightarrow LY = b$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 3 & 4 & 1 & 0 \\ -1 & -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ -3 \\ 4 \end{bmatrix}$$

$$\begin{aligned} y_1 &= 1 \\ 2y_1 + y_2 &= 1 \\ y_2 &= -1 \\ 3y_1 + 4y_2 + y_3 &= -3 \end{aligned}$$

$$y_3 = -2$$

$$-y_1 - 3y_2 + y_4 = 4$$

$$-1 + 3 + y_4 = 4$$

$$y_4 = 2$$

$$\Rightarrow Y = \begin{bmatrix} 1 \\ -1 \\ -2 \\ 2 \end{bmatrix}$$

Take

$$UX = Y \Rightarrow$$

$$\begin{bmatrix} 1 & 1 & 0 & 3 \\ 0 & -1 & -1 & -5 \\ 0 & 0 & 3 & 13 \\ 0 & 0 & 0 & -13 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -2 \\ 2 \end{bmatrix}$$

Use backward sub.

$$-13x_4 = 2$$

$$x_4 = -2/13,$$

from 3rd eqⁿ.

$$3x_3 + 13x_4 = -2$$

$$3x_3 + 13 \times \left(-\frac{2}{13}\right) = -2$$

$$\Rightarrow x_3 = 0$$

from 2nd eqⁿ.

$$-x_2 - x_3 - 5x_4 = -1$$

$$-x_2 - 0 - 5 \times \frac{-2}{13} = -1$$

$$-x_2 = -1 + \frac{10}{13} = -\frac{3}{13}$$

$$\Rightarrow x_2 = \frac{3}{13},$$

from 1st eqⁿ

$$x_1 + x_2 + 3x_4 = 1$$

$$x_1 + \frac{3}{13} + 3 \times \frac{-2}{13} = 1$$

$$x_1 = 1 + \frac{3}{13} = \frac{16}{13}$$

$$X = \begin{bmatrix} 16/13 \\ 3/13 \\ 0 \\ -2/13 \end{bmatrix} \sim \begin{bmatrix} 1.2308 \\ 0.2308 \\ 0 \\ -0.1538 \end{bmatrix}$$

System of linear equations:

Exercise:

- 1 Modify the LU Factorization Algorithm so that it can be used to solve a linear system, and then solve the following linear system:

$$2x_1 - x_2 + x_3 = -1$$

$$3x_1 + 3x_2 + 9x_3 = 0$$

$$3x_1 + 3x_2 + 5x_3 = 4.$$

- 2 Given the linear system

$$x_1 - x_2 + \alpha x_3 = -2$$

$$-x_1 + 2x_2 - \alpha x_3 = 3$$

$$\alpha x_1 + x_2 + x_3 = 2.$$

System of linear equations:

Exercise (continued):

- a** Find value(s) of α for which the system has no solution.
- b** Find value(s) of α for which the system has infinite number of solutions.
- c** Assuming a unique solution exists for a given α , find the solution.