

Risk Behavior Activity – Answer Key

A BIGGER RISK	A RISK	A SMALLER RISK	NO RISK
Sharing needles to inject drugs because it provides direct access of blood from the shared needle into the other person's body.	Drinking alcohol or using drugs at a party that can reduce a person's inhibitions and make them more likely to engage in risky sexual behavior.	Having vaginal or anal sex with a condom since a latex condom does not allow HIV to pass through, if used correctly and every time you have sex.	Abstaining from all types of sex and drugs since there would be no risk behavior or contact with a fluid that can transmit HIV.
Having vaginal or anal sex without a condom because these two behaviors are most risky due to the presence of semen, vaginal fluids and possibly blood and the delicate mucus membrane inside the vagina and anus that can tear easily.	Having oral sex without a condom or dental dam allows the semen or vaginal fluids of the partner into the mouth, which can have tears in the mucus membrane lining the mouth, of the partner who is giving oral sex.	Having sex with only the same partner and both people have been tested since the fewer sexual partners the less chance there is to be exposed to someone who is infected with HIV and both partners have been tested.	Being with someone who is crying, coughing or sneezing since none of those behaviors of fluids can transmit HIV.
Breastfeeding from a mother with HIV since breast milk can transmit HIV.	Dating someone who is a lot older since often they have had more sexual partners and have a position of power over the young partner who may not feel comfortable making the older partner use a condom or protection.	Kissing with an open mouth since small cuts inside the mouth might allow blood to transfer from one person to the other.	Donating blood since sterile equipment is used every time.
	Cutting with a shared knife or razor because blood is fluid with highest concentration of HIV and can infect another.		Kissing with a closed mouth since saliva alone can not transmit HIV.

Risk Behavior Activity – Answer Key

A BIGGER RISK	A RISK	A SMALLER RISK	NO RISK
	Sharing a toothbrush since sometimes there can be very tiny amounts of blood that can be transmitted into the mouth of another person.		Getting a mosquito bite since insects can not transmit HIV.
	Piercing or tattooing with a needle someone else already used because of the blood that can be present on the used needle.		Touching doorknobs, toilet seats or dishes since HIV can not live on surfaces like these.
			Going to school with a person who has HIV since there is no risk behavior or fluid involved.
			Shaking hands with someone with HIV since there is no risk behavior or fluid involved.
			Going swimming with someone who has HIV since there is no risk behavior or fluid involved.