

Estimating accuracy of hypotheses

Relevant Readings: 5.1, 5.2 in Mitchell

CS495 - Machine Learning, Fall 2009

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?
 - ▶ The difference of one data point may not be significant

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?
 - ▶ The difference of one data point may not be significant
 - ▶ Maybe this happened by “bad luck” and B is actually better in general

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?
 - ▶ The difference of one data point may not be significant
 - ▶ Maybe this happened by “bad luck” and B is actually better in general
- ▶ What if the testing set had 100,000 instances?

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?
 - ▶ The difference of one data point may not be significant
 - ▶ Maybe this happened by “bad luck” and B is actually better in general
- ▶ What if the testing set had 100,000 instances?
 - ▶ Then we have high confidence that A is more accurate than B for the measured task

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?
 - ▶ The difference of one data point may not be significant
 - ▶ Maybe this happened by “bad luck” and B is actually better in general
- ▶ What if the testing set had 100,000 instances?
 - ▶ Then we have high confidence that A is more accurate than B for the measured task
- ▶ There is an issue of *variance in the estimate*. How do we quantify all of this?

How much can we trust the measured accuracy?

- ▶ Algorithm A an accuracy of 100% on the testing set
- ▶ Algorithm B an accuracy of 90% on the testing set
- ▶ Question: Which algorithm is better?
- ▶ Answer: We can't really say without more information.
- ▶ What if the testing set had only 10 instances?
 - ▶ The difference of one data point may not be significant
 - ▶ Maybe this happened by “bad luck” and B is actually better in general
- ▶ What if the testing set had 100,000 instances?
 - ▶ Then we have high confidence that A is more accurate than B for the measured task
- ▶ There is an issue of *variance in the estimate*. How do we quantify all of this?

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit
- ▶ Suppose sample S is drawn from X according to \mathcal{D}

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit
- ▶ Suppose sample S is drawn from X according to \mathcal{D}
- ▶ The error rate that some hypothesis makes on S is called the *sample error*

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit
- ▶ Suppose sample S is drawn from X according to \mathcal{D}
- ▶ The error rate that some hypothesis makes on S is called the *sample error*
 - ▶ Easy to measure; just use the testing set

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit
- ▶ Suppose sample S is drawn from X according to \mathcal{D}
- ▶ The error rate that some hypothesis makes on S is called the *sample error*
 - ▶ Easy to measure; just use the testing set
- ▶ The error rate that some hypothesis makes on X (under distribution \mathcal{D}) is called the *true error*

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit
- ▶ Suppose sample S is drawn from X according to \mathcal{D}
- ▶ The error rate that some hypothesis makes on S is called the *sample error*
 - ▶ Easy to measure; just use the testing set
- ▶ The error rate that some hypothesis makes on X (under distribution \mathcal{D}) is called the *true error*
 - ▶ Harder to measure, but more important to know

The skiing example [Mitchell, 5.2]

- ▶ The instance space X is the set of all people
 - ▶ This includes some attributes (age, time since last visit, etc.)
- ▶ There is a probability distribution \mathcal{D} (unknown to us) that gives the probability that person $x \in X$ is the next one to walk in the ski shop
- ▶ Target concept $f : X \rightarrow \{0, 1\}$ is whether or not each person plans to buy skis when they visit
- ▶ Suppose sample S is drawn from X according to \mathcal{D}
- ▶ The error rate that some hypothesis makes on S is called the *sample error*
 - ▶ Easy to measure; just use the testing set
- ▶ The error rate that some hypothesis makes on X (under distribution \mathcal{D}) is called the *true error*
 - ▶ Harder to measure, but more important to know

Estimating true error (confidence intervals)

- ▶ The sample error provides an estimate of true error, but how good is the estimate?

Estimating true error (confidence intervals)

- ▶ The sample error provides an estimate of true error, but how good is the estimate?
- ▶ Under some reasonable assumptions (see [Mitchell, 5.2.2]), we can say with 99% probability that:

- ▶ $e - 2.58\sqrt{e(1-e)/n} \leq E \leq e + 2.58\sqrt{e(1-e)/n}$
- ▶ where e is the sample error,
- ▶ E is the true error, and
- ▶ $n \geq 30$ is the number of samples

Estimating true error (confidence intervals)

- ▶ The sample error provides an estimate of true error, but how good is the estimate?
- ▶ Under some reasonable assumptions (see [Mitchell, 5.2.2]), we can say with 99% probability that:
 - ▶ $e - 2.58\sqrt{e(1-e)/n} \leq E \leq e + 2.58\sqrt{e(1-e)/n}$
 - ▶ where e is the sample error,
 - ▶ E is the true error, and
 - ▶ $n \geq 30$ is the number of samples
- ▶ The 2.58 constant can be adjusted according to the certainty we require:

Estimating true error (confidence intervals)

- ▶ The sample error provides an estimate of true error, but how good is the estimate?
- ▶ Under some reasonable assumptions (see [Mitchell, 5.2.2]), we can say with 99% probability that:
 - ▶ $e - 2.58\sqrt{e(1-e)/n} \leq E \leq e + 2.58\sqrt{e(1-e)/n}$
 - ▶ where e is the sample error,
 - ▶ E is the true error, and
 - ▶ $n \geq 30$ is the number of samples
- ▶ The 2.58 constant can be adjusted according to the certainty we require:
 - ▶ For 99% certainty use 2.58
 - ▶ For 98% certainty use 2.33
 - ▶ For 95% certainty use 1.96
 - ▶ For 90% certainty use 1.64
 - ▶ For 80% certainty use 1.28
 - ▶ For 68% certainty use 1.00
 - ▶ For 50% certainty use 0.67

Estimating true error (confidence intervals)

- ▶ The sample error provides an estimate of true error, but how good is the estimate?
- ▶ Under some reasonable assumptions (see [Mitchell, 5.2.2]), we can say with 99% probability that:
 - ▶ $e - 2.58\sqrt{e(1-e)/n} \leq E \leq e + 2.58\sqrt{e(1-e)/n}$
 - ▶ where e is the sample error,
 - ▶ E is the true error, and
 - ▶ $n \geq 30$ is the number of samples
- ▶ The 2.58 constant can be adjusted according to the certainty we require:
 - ▶ For 99% certainty use 2.58
 - ▶ For 98% certainty use 2.33
 - ▶ For 95% certainty use 1.96
 - ▶ For 90% certainty use 1.64
 - ▶ For 80% certainty use 1.28
 - ▶ For 68% certainty use 1.00
 - ▶ For 50% certainty use 0.67