

Random variables, probability distributions,
expected value, variance, binomial distribution,
Gaussian distribution (normal distribution)

Relevant Readings: Table 5.2, Section 5.3 up through 5.3.4 in
Mitchell

CS495 - Machine Learning, Fall 2009

Prob and stats

- ▶ Probability, statistics, and sampling theory aren't only just useful in ML. These are generally useful in computer science (such as in the study of randomized algorithms)

Prob and stats

- ▶ Probability, statistics, and sampling theory aren't only just useful in ML. These are generally useful in computer science (such as in the study of randomized algorithms)

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die
- ▶ A random variable’s behavior is governed its *probability distribution*

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die
- ▶ A random variable’s behavior is governed its *probability distribution*
 - ▶ The probability distribution is a specification for the likelihood of the random variable taking on different values

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die
- ▶ A random variable’s behavior is governed its *probability distribution*
 - ▶ The probability distribution is a specification for the likelihood of the random variable taking on different values
 - ▶ Example: For rolling a die, the probability distribution is *uniform*, meaning that each possibility (1, 2, 3, 4, 5, 6) has equal probability of coming up ($1/6$).

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die
- ▶ A random variable’s behavior is governed its *probability distribution*
 - ▶ The probability distribution is a specification for the likelihood of the random variable taking on different values
 - ▶ Example: For rolling a die, the probability distribution is *uniform*, meaning that each possibility (1, 2, 3, 4, 5, 6) has equal probability of coming up ($1/6$).
 - ▶ Random variables can be discrete or continuous

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die
- ▶ A random variable’s behavior is governed its *probability distribution*
 - ▶ The probability distribution is a specification for the likelihood of the random variable taking on different values
 - ▶ Example: For rolling a die, the probability distribution is *uniform*, meaning that each possibility (1, 2, 3, 4, 5, 6) has equal probability of coming up (1/6).
 - ▶ Random variables can be discrete or continuous
 - ▶ The probability that random variable X takes on the value x is denoted $\Pr(X = x)$

Some basics

- ▶ A *random variable* is a numerical outcome of some random “experiment”
 - ▶ It can be thought of as an unknown value that changes every time it is observed
 - ▶ Similar to `Math.random()`
 - ▶ Example: The result of rolling a die
- ▶ A random variable’s behavior is governed its *probability distribution*
 - ▶ The probability distribution is a specification for the likelihood of the random variable taking on different values
 - ▶ Example: For rolling a die, the probability distribution is *uniform*, meaning that each possibility (1, 2, 3, 4, 5, 6) has equal probability of coming up (1/6).
 - ▶ Random variables can be discrete or continuous
 - ▶ The probability that random variable X takes on the value x is denoted $\Pr(X = x)$

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$
 - ▶ So then for continuous random variables, we work in terms of ranges of values, rather than the probability of equaling some value exactly

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$
 - ▶ So then for continuous random variables, we work in terms of ranges of values, rather than the probability of equaling some value exactly
 - ▶ Think of it this way: what's the probability of throwing a dart and hitting *exactly* the right spot on the wall?

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$
 - ▶ So then for continuous random variables, we work in terms of ranges of values, rather than the probability of equaling some value exactly
 - ▶ Think of it this way: what's the probability of throwing a dart and hitting *exactly* the right spot on the wall?
 - ▶ Zero.

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$
 - ▶ So then for continuous random variables, we work in terms of ranges of values, rather than the probability of equaling some value exactly
 - ▶ Think of it this way: what's the probability of throwing a dart and hitting *exactly* the right spot on the wall?
 - ▶ Zero.
 - ▶ But after it's thrown, it actually hit *some* exact spot.

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$
 - ▶ So then for continuous random variables, we work in terms of ranges of values, rather than the probability of equaling some value exactly
 - ▶ Think of it this way: what's the probability of throwing a dart and hitting *exactly* the right spot on the wall?
 - ▶ Zero.
 - ▶ But after it's thrown, it actually hit *some* exact spot.
 - ▶ Therefore zero-probability doesn't necessarily mean "impossible"

Continuous probability distributions

- ▶ Consider a uniform continuous random variable X over the range $[0,2]$
 - ▶ What is $\Pr(X = 1/2)$?
 - ▶ Zero.
 - ▶ What is $\Pr(X = 1/4)$?
 - ▶ Zero.
 - ▶ What is $\Pr(1/4 \leq X \leq 1/2)$?
 - ▶ $1/8$
 - ▶ So then for continuous random variables, we work in terms of ranges of values, rather than the probability of equaling some value exactly
 - ▶ Think of it this way: what's the probability of throwing a dart and hitting *exactly* the right spot on the wall?
 - ▶ Zero.
 - ▶ But after it's thrown, it actually hit *some* exact spot.
 - ▶ Therefore zero-probability doesn't necessarily mean "impossible"

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$
- ▶ The *expected value* (aka. *mean*) of a random variable Y is $E[Y] = \sum_i y_i \cdot \Pr(Y = y_i)$, where the sum is over all possible outcomes

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$
- ▶ The *expected value* (aka. *mean*) of a random variable Y is $E[Y] = \sum_i y_i \cdot \Pr(Y = y_i)$, where the sum is over all possible outcomes
 - ▶ Just a weighted average (the continuous case would be an integral).

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$
- ▶ The *expected value* (aka. *mean*) of a random variable Y is $E[Y] = \sum_i y_i \cdot \Pr(Y = y_i)$, where the sum is over all possible outcomes
 - ▶ Just a weighted average (the continuous case would be an integral).
- ▶ The *variance* of a random variable Y is $\text{Var}(Y) = E[(Y - E[Y])^2]$

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$
- ▶ The *expected value* (aka. *mean*) of a random variable Y is $E[Y] = \sum_i y_i \cdot \Pr(Y = y_i)$, where the sum is over all possible outcomes
 - ▶ Just a weighted average (the continuous case would be an integral).
- ▶ The *variance* of a random variable Y is $\text{Var}(Y) = E[(Y - E[Y])^2]$
 - ▶ This is a measure of how much Y deviates from the mean

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$
- ▶ The *expected value* (aka. *mean*) of a random variable Y is $E[Y] = \sum_i y_i \cdot \Pr(Y = y_i)$, where the sum is over all possible outcomes
 - ▶ Just a weighted average (the continuous case would be an integral).
- ▶ The *variance* of a random variable Y is $Var(Y) = E[(Y - E[Y])^2]$
 - ▶ This is a measure of how much Y deviates from the mean
- ▶ The *standard deviation* of a random variable Y is simply $\sqrt{Var(Y)}$

Some basic notions

- ▶ Summation notation: $\sum_{i=0}^n a_i$ means $a_0 + a_1 + \cdots + a_n$
- ▶ The *expected value* (aka. *mean*) of a random variable Y is $E[Y] = \sum_i y_i \cdot \Pr(Y = y_i)$, where the sum is over all possible outcomes
 - ▶ Just a weighted average (the continuous case would be an integral).
- ▶ The *variance* of a random variable Y is $Var(Y) = E[(Y - E[Y])^2]$
 - ▶ This is a measure of how much Y deviates from the mean
- ▶ The *standard deviation* of a random variable Y is simply $\sqrt{Var(Y)}$

Discrete uniform distribution

- ▶ Suppose X is a discrete random variable with uniform probability distribution over the set $\{1, 2, 3, \dots, m\}$

Discrete uniform distribution

- ▶ Suppose X is a discrete random variable with uniform probability distribution over the set $\{1, 2, 3, \dots, m\}$
- ▶ What is $E[X]$?

Discrete uniform distribution

- ▶ Suppose X is a discrete random variable with uniform probability distribution over the set $\{1, 2, 3, \dots, m\}$
- ▶ What is $E[X]$?
 - ▶ $(m + 1)/2$

Discrete uniform distribution

- ▶ Suppose X is a discrete random variable with uniform probability distribution over the set $\{1, 2, 3, \dots, m\}$
- ▶ What is $E[X]$?
 - ▶ $(m + 1)/2$
- ▶ What is $Var[X]$?

Discrete uniform distribution

- ▶ Suppose X is a discrete random variable with uniform probability distribution over the set $\{1, 2, 3, \dots, m\}$
- ▶ What is $E[X]$?
 - ▶ $(m + 1)/2$
- ▶ What is $Var[X]$?
 - ▶ $(m^2 - 1)/12$

Discrete uniform distribution

- ▶ Suppose X is a discrete random variable with uniform probability distribution over the set $\{1, 2, 3, \dots, m\}$
- ▶ What is $E[X]$?
 - ▶ $(m + 1)/2$
- ▶ What is $Var[X]$?
 - ▶ $(m^2 - 1)/12$

Continuous uniform distribution

- ▶ Suppose X is a continuous random variable with uniform probability distribution over the interval $[a, b]$

Continuous uniform distribution

- ▶ Suppose X is a continuous random variable with uniform probability distribution over the interval $[a, b]$
- ▶ What is $E[X]$?

Continuous uniform distribution

- ▶ Suppose X is a continuous random variable with uniform probability distribution over the interval $[a, b]$
- ▶ What is $E[X]$?
 - ▶ $(a + b)/2$

Continuous uniform distribution

- ▶ Suppose X is a continuous random variable with uniform probability distribution over the interval $[a, b]$
- ▶ What is $E[X]$?
 - ▶ $(a + b)/2$
- ▶ What is $Var[X]$?

Continuous uniform distribution

- ▶ Suppose X is a continuous random variable with uniform probability distribution over the interval $[a, b]$
- ▶ What is $E[X]$?
 - ▶ $(a + b)/2$
- ▶ What is $Var[X]$?
 - ▶ $(b - a)^2/12$

Continuous uniform distribution

- ▶ Suppose X is a continuous random variable with uniform probability distribution over the interval $[a, b]$
- ▶ What is $E[X]$?
 - ▶ $(a + b)/2$
- ▶ What is $Var[X]$?
 - ▶ $(b - a)^2/12$

Binomial distribution

- Imagine an unfair coin n times that has probability p of landing on heads

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete
- ▶ Suppose X is a random variable with binomial probability distribution

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete
- ▶ Suppose X is a random variable with binomial probability distribution
- ▶ What is $E[X]$?

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete
- ▶ Suppose X is a random variable with binomial probability distribution
- ▶ What is $E[X]$?
 - ▶ np

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete
- ▶ Suppose X is a random variable with binomial probability distribution
- ▶ What is $E[X]$?
 - ▶ np
- ▶ What is $Var[X]$?

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete
- ▶ Suppose X is a random variable with binomial probability distribution
- ▶ What is $E[X]$?
 - ▶ np
- ▶ What is $Var[X]$?
 - ▶ $np(1 - p)$

Binomial distribution

- ▶ Imagine an unfair coin n times that has probability p of landing on heads
- ▶ A binomial distribution is: “how many times did we get heads?”
- ▶ Is this a discrete or continuous distribution?
 - ▶ Discrete
- ▶ Suppose X is a random variable with binomial probability distribution
- ▶ What is $E[X]$?
 - ▶ np
- ▶ What is $Var[X]$?
 - ▶ $np(1 - p)$

Gaussian (normal) distribution

- ▶ If you add lots of random variables together, the sum is in itself a random variable whose probability distribution tends to look like a Gaussian.

Gaussian (normal) distribution

- ▶ If you add lots of random variables together, the sum is in itself a random variable whose probability distribution tends to look like a Gaussian.
- ▶ Thus, Gaussian distributions show up a lot in nature

Gaussian (normal) distribution

- ▶ If you add lots of random variables together, the sum is in itself a random variable whose probability distribution tends to look like a Gaussian.
- ▶ Thus, Gaussian distributions show up a lot in nature
- ▶ It's a “bell shaped curve”

Gaussian (normal) distribution

- ▶ If you add lots of random variables together, the sum is in itself a random variable whose probability distribution tends to look like a Gaussian.
- ▶ Thus, Gaussian distributions show up a lot in nature
- ▶ It's a “bell shaped curve”
- ▶ The probability distribution for a Gaussian with mean μ and variance σ^2 is defined as:

Gaussian (normal) distribution

- ▶ If you add lots of random variables together, the sum is in itself a random variable whose probability distribution tends to look like a Gaussian.
- ▶ Thus, Gaussian distributions show up a lot in nature
- ▶ It's a “bell shaped curve”
- ▶ The probability distribution for a Gaussian with mean μ and variance σ^2 is defined as:
 - ▶ $e^{-(x-\mu)^2/(2\sigma^2)} / (\sigma\sqrt{2\pi})$

Gaussian (normal) distribution

- ▶ If you add lots of random variables together, the sum is in itself a random variable whose probability distribution tends to look like a Gaussian.
- ▶ Thus, Gaussian distributions show up a lot in nature
- ▶ It's a “bell shaped curve”
- ▶ The probability distribution for a Gaussian with mean μ and variance σ^2 is defined as:
 - ▶ $e^{-(x-\mu)^2/(2\sigma^2)} / (\sigma\sqrt{2\pi})$

Application

- ▶ Suppose we are programming Naive Bayes and want to handle a continuous variable

Application

- ▶ Suppose we are programming Naive Bayes and want to handle a continuous variable
- ▶ One way we could do this: measure the mean and variance of the relevant examples, model them with a probability distribution (such as a Gaussian), and use that distribution to help determine whether “yes” or “no” is more likely for the given instance

Application

- ▶ Suppose we are programming Naive Bayes and want to handle a continuous variable
- ▶ One way we could do this: measure the mean and variance of the relevant examples, model them with a probability distribution (such as a Gaussian), and use that distribution to help determine whether “yes” or “no” is more likely for the given instance