

# My paper on the cosmic microwave background and formation of structures in our Universe

V. A. Vikenes<sup>1</sup>

Institute of Theoretical Astrophysics, University of Oslo, 0315 Oslo, Norway  
e-mail: v.a.vikenes@astro.uio.no

May 25, 2023

## ABSTRACT

The code for this project can be found on my GitHub repository: <https://github.com/Vikenes/AST5220/>

**Key words.** cosmic microwave background – large-scale structure of Universe

## 1. Introduction

Write an introduction here. Give context to the paper. Citations to relevant papers. You only need to do this in the end for the last milestone.

## 2. Milestone I

In this section we will examine the evolution of the Universe's uniform background. Our primary objective is to develop methods for computing the Hubble parameter and related time- and distance measures. These methods provide a first step towards further investigations and modelling of the early Universe. To compute the background cosmology, we will solve ordinary differential equations (ODEs) numerically, using cosmological parameters obtained from the Planck Collaboration (Planck Collaboration et al. 2020). The parameters we will use are listed in Eq. (A.1) in Appendix A. One crucial aspect in the process is validating our model. We will therefore develop some simple methods for comparing our result. This will mainly involve considering simplified cases where analytical solutions can be obtained.

Our primary focus in this section concerns methods where the cosmological parameters are given from the start. Another interesting aspect is to use data to constrain cosmological parameters. To do this, we will use data from supernova observations (Betoule et al. 2014), containing luminosity distance associated with different values of redshift. By employing the numerical methods we develop initially, we will try to estimate optimal values of three cosmological parameters, by implement a simple Markov chain Monte Carlo (MCMC) algorithm. The parameters we will be sampling are  $h$ ,  $\Omega_{m0}$  and  $\Omega_{\Lambda0}$ . From these results, we will investigate confidence regions of  $\Omega_{m0}$  and  $\Omega_{\Lambda0}$ , and try to estimate a probability distribution function (PDF) for the Hubble parameter.

## 2.1. Theory

### 2.1.1. Density parameters and Hubble factor

The Friedmann equation can be written in terms of density parameters,  $\Omega_i \equiv \rho_i/\rho_c$ , where  $\rho_c \equiv 3H^2/8\pi G$  is the critical density. The density of a given species,  $i$ , evolves as (Dodelson 2020, Eq. (2.61))

$$\rho_i(t) \propto a(t)^{-3(1+w_i)}, \quad (1)$$

where we have assumed that the equation of state (EoS) parameter,  $w_i \equiv P_i/\rho_i$  (Dodelson 2020, Eq. 2.60), is constant.  $P_i$  denotes the pressure of the species. We will limit ourselves to consider three types of species in this report: matter, radiation and dark energy. We will only consider baryons and cold dark matter (CDM) for the matter component, which we express as  $\Omega_{m0} = \Omega_{b0} + \Omega_{CDM0}$ . The subscript 0 is used to refer to today's value. The radiation component we consider is  $\Omega_{r0} = \Omega_{\gamma0} + \Omega_{\nu0}$ , corresponding to photons and neutrinos, respectively. For the dark energy, we only have the cosmological constant  $\Omega_{\Lambda0}$ .

Matter, radiation and dark energy have densities evolving according to Eq. (1) with  $w_i = 0, 1/3$  and  $-1$ , respectively. Neutrinos having  $w = 1/3$  only holds since we will assume that the neutrinos are massless. Curvature can be described by  $w_i = -1/3$ , with  $\Omega_{k0} \equiv -kc^2/H_0^2$ . The parameter  $k$  represents the curvature of the Universe, where  $k = 0$  corresponds to a flat Universe. With these parameters, the Friedmann Equation can be written as (Dodelson 2020, Eq. (3.14))

$$H = H_0 \sqrt{\Omega_{m0}a^{-3} + \Omega_{r0}a^{-4} + \Omega_{k0}a^{-2} + \Omega_{\Lambda0}}, \quad (2)$$

where  $H \equiv \dot{a}/a$  is the Hubble parameter, with the dot denoting a derivative with respect to cosmic time,  $t$ . For the radiation,  $\Omega_{\gamma0}$  and  $\Omega_{\nu0}$  follow from the temperature of the CMB today,  $T_{CMB0}$ , and the effective number of massless neutrinos,  $\mathcal{N}_{\text{eff}}$ . They are given by

$$\Omega_{\gamma0} = 2 \cdot \frac{\pi^2}{30} \frac{(k_b T_{CMB0})^4}{\hbar^3 c^5} \cdot \frac{8\pi G}{3H_0^2}, \quad (3)$$

$$\Omega_{\nu0} = \mathcal{N}_{\text{eff}} \cdot \frac{7}{8} \cdot \left(\frac{4}{11}\right)^{4/3} \Omega_{\gamma0}. \quad (4)$$

The value of  $\Omega_{\Lambda 0}$  is fixed by the requirement that  $H(a = 1) = H_0$ , yielding

$$\Omega_{\Lambda 0} = 1 - (\Omega_{m0} + \Omega_{r0} + \Omega_{k0}). \quad (5)$$

We also introduce the scaled Hubble factor,  $\mathcal{H} \equiv aH$ . Rather than working with the scale factor,  $a(t)$ , we will mainly be working with the logarithm of the scale factor

$$x \equiv \ln a, \quad ' \equiv \frac{d}{dx}. \quad (6)$$

The resulting expression for  $\mathcal{H}(x)$  is thus

$$\mathcal{H}(x) = H_0 \sqrt{\Omega_{m0}e^{-x} + \Omega_{r0}e^{-2x} + \Omega_{k0} + \Omega_{\Lambda 0}e^{2x}}. \quad (7)$$

This form of the Hubble factor is the one we will focus on for the majority of this report. In terms of  $\mathcal{H}(x)$ , the value of the density parameters can be obtained at any given  $x$ , with

$$\Omega_k(x) = \frac{\Omega_{k0}}{\mathcal{H}(x)^2/H_0^2}, \quad (8)$$

$$\Omega_m(x) = \frac{\Omega_{m0}}{e^x \mathcal{H}(x)^2/H_0^2}, \quad (9)$$

$$\Omega_r(x) = \frac{\Omega_{r0}}{e^{2x} \mathcal{H}(x)^2/H_0^2}, \quad (10)$$

$$\Omega_{\Lambda}(x) = \frac{\Omega_{\Lambda 0}}{e^{-2x} \mathcal{H}(x)^2/H_0^2}, \quad (11)$$

From these expressions, we can identify the epochs during which the Universe was dominated by an equal amount of matter and radiation, and by an equal amount of matter and dark energy. These epochs are defined by the time when  $\Omega_m = \Omega_r$  and  $\Omega_m = \Omega_{\Lambda}$ , respectively, and are a valuable asset towards understanding the physics governing the evolution of the Universe. Another time of interest is the onset of acceleration, defined as the time when  $\ddot{a} = 0$ . In terms of  $\mathcal{H}$  and  $x$ , this corresponds to

$$\ddot{a} = \frac{dx}{dt} \frac{da}{dx} = \frac{d \ln a}{dt} \frac{d\mathcal{H}(x)}{dx} = e^{-x} \mathcal{H}(x) \frac{d\mathcal{H}(x)}{dx}. \quad (12)$$

In Sect. 2.1.4 we will derive an expression for  $\mathcal{H}'(x)$ .

### 2.1.2. Conformal time

We now want to relate the Hubble factor to some time variables. The main one we will consider is the conformal time,  $\eta$ . It is a measure of the distance light has been able to travel since  $t = 0$ , where  $t$  is the cosmic time. Using its definition in terms of  $t$  (Dodelson 2020, Eq. (2.90)), we can express it in terms of  $x$  as

$$\eta = \int_0^t \frac{c dt'}{a(t')} = \int_{-\infty}^{x'} \frac{c dx'}{\mathcal{H}(x')}. \quad (13)$$

This leads us to the following differential equation that we will solve numerically

$$\frac{d\eta}{dx} = \frac{c}{\mathcal{H}(x)}. \quad (14)$$

The initial condition we have is  $\eta(-\infty) = 0$ . Noting from Eq. (7) that  $\mathcal{H}(x) \rightarrow H_0 \sqrt{\Omega_{r0}e^{-2x}}$  as  $x \rightarrow -\infty$ , we get an analytical approximation for the initial condition of  $\eta$  at early times

$$\eta(x_{\text{start}}) \approx \int_{-\infty}^{x_{\text{start}}} \frac{c dx'}{H_0 \sqrt{\Omega_{r0}}} e^{x'} = \frac{c}{\mathcal{H}(x_{\text{start}})}. \quad (15)$$

Note that  $\eta(x)\mathcal{H}(x)/c \rightarrow 1$  at low  $x$ , which provides a natural way of validating our implementation.

For the cosmic time,  $t$ , starting from  $H = \dot{a}/a$  and applying the chain rule yields the desired differential equation for  $t(x)$ , which we will solve numerically,

$$\frac{dt}{dx} = \frac{1}{H(x)}. \quad (16)$$

To get an initial condition for  $t$ , we consider the radiation dominating era, with the following integral expression

$$t(x) = \int_{-\infty}^x \frac{dx'}{H(x')}. \quad (17)$$

Comparing with Eq. (15), we see that the two integrands only differ by a factor  $e^x$ . The initial condition for  $t$  is therefore easily seen to be

$$t(x_{\text{start}}) = \frac{1}{2H(x_{\text{start}})}. \quad (18)$$

### 2.1.3. Distance measures

The supernova data we will study has distances measured in terms of luminosity distance,  $d_L$ . Expressing it in terms of the angular distance,  $d_A = ar$ , it becomes

$$d_L(a) = \frac{d_A}{a^2} = \frac{r}{a} \implies d_L(x) = e^{-x}r. \quad (19)$$

Here,  $r$  represents the radial coordinate of the emitted photon. To get an expression for  $r$ , we consider a photon's line-element in spherical coordinates,

$$ds^2 = -c^2 dt^2 + a^2 \left( \frac{dr^2}{1 - kr^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right). \quad (20)$$

For photons travelling radially towards us, we have  $d\theta = d\phi = 0$ . Since  $ds^2 = 0$  for photons, integrating the line-element of a photon emitted at,  $(t, r)$ , reaching an observer at  $(t_0, 0)$ , yields

$$\int_0^r \frac{dr'}{\sqrt{1 - kr'^2}} = \int_t^{t_0} \frac{c dt}{a}. \quad (21)$$

The RHS of Eq. (21) is known as the co-moving distance,  $\chi$ , which in terms of conformal time is given as

$$\chi = \int_t^{t_0} \frac{c dt}{a} = \int_x^0 \frac{c dx'}{\mathcal{H}(x')} = \eta(0) - \eta(x). \quad (22)$$

Solving Eq. (21) with respect to  $r$ , we get

$$r = \begin{cases} \chi \cdot \frac{\sin(\sqrt{|\Omega_{k0}|}H_0\chi/c)}{(\sqrt{|\Omega_{k0}|}H_0\chi/c)}, & \Omega_{k0} < 0, \\ \chi, & \Omega_{k0} = 0, \\ \chi \cdot \frac{\sinh(\sqrt{|\Omega_{k0}|}H_0\chi/c)}{(\sqrt{|\Omega_{k0}|}H_0\chi/c)}, & \Omega_{k0} > 0. \end{cases} \quad (23)$$

Eq. (19) can now be used to compute  $d_L$ , and the expression to use depends on the curvature.

#### 2.1.4. Analytical solutions

In Sect. 2.1.2 we discussed how  $\eta(x)$  can be used to test our implementation in the radiation dominating era. To test our solutions in other regimes, we will need the first and second derivative of  $\mathcal{H}(x)$ . To simplify the resulting expressions, we define the function,  $g(x)$ , as the derivative of the term inside the square root in Eq. (7), namely

$$g(x) \equiv -\Omega_{m0}e^{-x} - 2\Omega_{r0}e^{-2x} + 2\Omega_{\Lambda0}e^{2x}. \quad (24)$$

The first two derivatives of  $\mathcal{H}(x)$  are easily seen to be

$$\frac{d\mathcal{H}(x)}{dx} = \frac{H_0^2}{2\mathcal{H}(x)}g(x), \quad (25)$$

$$\frac{d^2\mathcal{H}(x)}{dx^2} = \frac{H_0^2}{2\mathcal{H}(x)} \left[ g'(x) - \frac{1}{2} \left( \frac{H_0 g(x)}{\mathcal{H}(x)} \right)^2 \right]. \quad (26)$$

Now we will consider the situation where the Universe is dominated by a single fluid with a constant EoS parameter,  $w$ . In that case we have  $H(t)^2 \propto \rho_i(t)^2$  (Dodelson 2020, Eq. (3.13)). Using Eq. (1), the Hubble parameter expressed in terms of  $w_i$  becomes

$$H(t)^2 \propto a^{-3(1+w)} \implies \mathcal{H}(x) = c_1 e^{-\frac{3}{2}(1+w)x}, \quad (27)$$

where  $c_1$  is some constant. The reason for doing this, is that both  $c_1$  and the exponential factor drops out when we consider  $\mathcal{H}'(x)/\mathcal{H}(x)$  and  $\mathcal{H}''(x)/\mathcal{H}(x)$ . For different values of  $w_i$ ,  $\mathcal{H}'(x)/\mathcal{H}(x)$  becomes

$$\frac{1}{\mathcal{H}(x)} \frac{d\mathcal{H}}{dx} = -\frac{1+3w}{2} = \begin{cases} -1, & w = 1/3, \\ -1/2, & w = 0, \\ 1, & w = -1. \end{cases} \quad (28)$$

Similarly, the expression for  $\mathcal{H}''(x)/\mathcal{H}(x)$  becomes

$$\frac{1}{\mathcal{H}(x)^2} \frac{d^2\mathcal{H}}{dx^2} = \frac{(1+3w)^2}{2} = \begin{cases} 1, & w = 1/3 \\ 1/4, & w = 0 \\ 1, & w = -1 \end{cases} \quad (29)$$

Equations (28) and (29) offer a means to evaluate the accuracy of our numerical solution at different regimes. Each density parameter evolve differently with  $x$ , as seen from Eqs. (8)-(11). Certain ranges of  $x$ -values will therefore closely resemble a Universe that is dominated by a single fluid. By computing  $\mathcal{H}$ ,  $\mathcal{H}'$ , and  $\mathcal{H}''$ , we can examine whether these quantities exhibit the expected behaviour. This allows us to assess the validity of our model and ensure that it is consistent with the underlying physical principles.

#### 2.2. Implementation details

To solve the differential equations for  $\eta$  and  $t$  (Eq. (14) and (16)) we use the C++ library GSL Galassi (2009), and use their Runge-Kutta4 solver. From the solution we create a spline of the results for the given  $x$  domain we have considered.

We will consider three different ranges of  $x$ -values. For the initial testing, we will use  $x \in [\ln 10^{-10}, 5]$ . For fitting cosmological parameters to the supernova data, we will use  $x \in [\ln 10^{-2}, 0]$ . When we want to estimate important times during the cosmic evolution, we will consider  $x \in [-10, 1]$ , for increased resolution, as the result may vary by a noticeable amount between step sizes. In all cases, we use  $N_x = 10^5$  number of points.

The cosmological parameters we consider assume  $\Omega_{k0} = 0$ . In Sect. 2.2.1 we discuss how we will use supernova data to estimate a value for  $\Omega_{k0}$ . Curvature is therefore implemented in all the relevant methods, but we set  $\Omega_{k0} = 0$  when we're not dealing with supernova fitting.

##### 2.2.1. Supernova fitting and parameter sampling

The supernova data we will use contains  $N = 31$  data points of luminosity distance,  $d_L^{\text{obs}}(z_i)$ , with associated measurement errors,  $\sigma_i$ , at different redshifts,  $z_i \in [0.01, 1.30]$ . This corresponds to  $x \sim [-9.95 \cdot 10^{-3}, -0.833]$ . Using these measurements, we want to constrain the three-dimensional parameter space

$$C = \{\hat{h}, \hat{\Omega}_{m0}, \hat{\Omega}_{k0}\}, \quad (30)$$

where the hat is used to distinguish the estimated parameters from the fiducial ones. We use  $\Omega_{b0} = 0.05$  for this analysis, so  $\hat{\Omega}_{m0}$  enters via  $\Omega_{\text{CDM}0} = \hat{\Omega}_{m0} - \Omega_{b0}$ . Additionally, the neutrinos are not relevant at the small scale considered here, and we therefore set  $N_{\text{eff}} = 0$  for this analysis.

We will assume that the measurements at different redshifts are normal distributed and uncorrelated. The likelihood function is then given by  $L \propto e^{-\chi^2/2}$ , where

$$\chi^2(C) = \sum_{i=1}^N \frac{[d_L(z_i, C) - d_L^{\text{obs}}(z_i)]^2}{\sigma_i^2}, \quad (31)$$

is the function we want to minimize. To do this, we will sample parameter values randomly by a Markov chain Monte Carlo (MCMC) process. We also restrict the parameter space to sample, with the following limits:

$$\begin{aligned} 0.5 < \hat{h} < 1.5, \\ 0 < \hat{\Omega}_{m0} < 1, \\ -1 < \hat{\Omega}_{k0} < 1. \end{aligned} \quad (32)$$

To generate a new sample, we update each parameter by generating a random number  $P \sim \mathcal{N}(0, 1)$ , and multiplying it by a step size. We will use step sizes of  $\Delta \hat{h} = 0.007$ ,  $\Delta \hat{\Omega}_{m0} = 0.05$ ,  $\Delta \hat{\Omega}_{k0} = 0.05$ . To determine whether a new configuration should be included in the sample we use the Metropolis algorithm, where we always accept a state if it yields a lower value of  $\chi^2$  compared to the previous state that was accepted. If the new value of  $\chi^2$  is greater than the old one, we accept it if the ratio of the likelihood functions  $L(\chi_{\text{new}}^2)/L(\chi_{\text{old}}^2) > p$ , where  $p \sim \mathcal{U}(0, 1)$ . We continue drawing samples until we get a total of  $\hat{n} = 10^4$  samples. For the samples generated, we omit the first 1000 samples of the chain from our analysis.

With our generated samples, we can use the best fit,  $\chi_{\text{min}}^2$ , to find the  $1\sigma$  and  $2\sigma$  confidence regions. For the  $\chi^2$  distribution with 3 parameters, these regions are given by  $\chi^2 - \chi_{\text{min}}^2 < 3.53$  and  $\chi^2 - \chi_{\text{min}}^2 < 8.02$ , respectively. We will plot the  $1\sigma$  and  $2\sigma$  constraint in the  $(\Omega_{m0}, \Omega_{\Lambda0})$  plane. Since  $\Omega_{r0} < 10^{-4}$ ,  $\Omega_{\Lambda0}$  can be approximated well by  $\Omega_{\Lambda0} = 1 - \Omega_{m0}$ . After that we will plot the posterior probability distribution function (PDF) for  $H_0$ .

To compare our fit with the Planck data, we will plot  $d_L^{\text{obs}}(z_i)$  together with  $d_L^{\text{fit}}(z)$  and  $d_L^{\text{Planck}}(z)$ . We obtain the former by solving the background cosmology with  $h$ ,  $\Omega_{m0}$ ,  $\Omega_{k0}$  replaced by the configuration  $\hat{h}$ ,  $\hat{\Omega}_{m0}$ ,  $\hat{\Omega}_{k0}$  that yielded the lowest value of  $\chi^2$ .

### 2.3. Results

In this section we present the results from our numerical simulations. Most of our results concern the evolution of various parameters as a function of  $x$ . Whenever it's relevant for interpreting and understanding the plot, we mark the point where we have matter-radiation equality and matter-dark energy equality, corresponding to  $\Omega_m = \Omega_r$  and  $\Omega_m = \Omega_\Lambda$ , respectively. These points can be seen directly in Fig. 5.

#### 2.3.1. Analytical and numerical comparisons


The dimensionless quantity  $\eta\mathcal{H}/c$  is shown in Fig. 1. At the lowest values of  $x \lesssim -10$ , we see that  $\eta\mathcal{H}/c = 1$ , as expected. Slightly before matter-radiation equality takes place, we see a slight increase towards higher  $x$ . As we approach higher  $x$ ,  $\Omega_\Lambda$  starts dominating, and the solution eventually diverges, as expected.



**Fig. 1.** The dimensionless quantity  $\eta\mathcal{H}/c$  as a function of  $x$ . At low  $x$  it has a value of 1, as expected. The most significant changes occur near regions where there is a change in which density parameter is dominating.

In Fig. 2 we have plotted  $\mathcal{H}'/\mathcal{H}$  and  $\mathcal{H}''/\mathcal{H}$ , where we include the analytical approximation from Eq. (28) and Eq. (29), respectively. The different values of  $w$  are drawn over the whole range of  $x$  where their related density parameter is larger than the other two. This is done for visibility purposes, and we only expect approximations to be reasonable whenever a density parameter is close to 1.

Towards the smallest values of  $x$  we see that both quantities are well approximated by the analytical solutions for  $w = 1/3$ . As we reach  $x \gtrsim -12$ , matter becomes increasingly dominant, and the solution deviates from being purely dominated by a  $w = 1/3$  fluid. Towards the highest values of  $x$ , we see that both quantities reach a constant value of 1 for  $w = -1$ . At higher values of  $x$ , we know that both  $\Omega_m(x)$  and  $\Omega_r(x)$  should vanish, eventually, while  $\Omega_\Lambda \rightarrow 1$ . This behaviour is thus present in our implementation. When matter dominates, we see that both functions exhibit inferior agreement with the approximations compared to the other regimes. This can be understood from Fig. 5, where the vanishing contribution of  $\Omega_r$  occurs around the same time as  $\Omega_\Lambda$  starts contributing. The maximum value reached for this particular configuration is  $\Omega_m \approx 0.995$ . Nonetheless, significant deviations from the analytical solutions are not evident.




**Fig. 2.**  $\mathcal{H}'/\mathcal{H}$  and  $\mathcal{H}''/\mathcal{H}$  compared with analytical expressions in the case a single fluid with a given EoS parameter,  $w$ , shown by the dashed lines. The transition between the dashed lines are chosen as the corresponding epochs of equality.

#### 2.3.2. Hubble factor and time variables

Having checked that our implementation is physical, we now proceed by studying the evolution of the background, starting with a plot of the conformal Hubble factor,  $\mathcal{H}(x)$ , shown in Fig. 3.

The local minima taking place before matter-dark energy equality corresponds to the onset of acceleration, where  $\ddot{a} = 0$ . For  $x > 0$ ,  $\Omega_\Lambda$  will dominate the conformal Hubble factor, where we have  $\mathcal{H}(x) \propto e^x$ , as seen from Eq. (7).




**Fig. 3.** Evolution of the conformal Hubble factor. The minimum point marks the onset of acceleration, after which, dark energy dominates its evolution.

The evolution of  $\eta(x)$  and  $t(x)$  is shown in Fig. 4. At high values of  $x$ , the  $e^x$  dependence of  $\mathcal{H}(x)$  causes  $\eta(x)$  to grow as  $e^{-x}$  at late times, suppressing its growth. The cosmic time, on the other hand, does not have an exponential dependence in the integrand at high values of  $x$ . This yields the linear growth we see at late times. The different  $x$ -dependence of  $t$  and  $\eta$  results in the two quantities to be approximately equal at  $x \sim 2.75$ .



**Fig. 4.** The cosmic time,  $t$  and conformal time  $\eta/c$ . We emphasize that  $\eta \neq t$  at small  $x$ , but the difference is discernable on the scale considered.



**Fig. 5.** Evolution of the density parameters over time. The points where  $\Omega_m = \Omega_r$  marks the epoch of matter-radiation equality, and the intersection between  $\Omega_m$  and  $\Omega_\Lambda$  marks the epoch of matter-dark energy equality. The sum of the three quantities never exceed 1, as expected.


### 2.3.3. Supernova fitting

A plot showing the luminosity distance as a function of redshift is shown in Fig. 6, where we plot  $d_L(z)/z$  to better compare the simulations with the data. There is a noticeable discrepancy between the simulated luminosity distance and the data, with most redshifts causing the simulation to fall outside the uncertainties.


The  $1\sigma$  and  $2\sigma$  confidence regions in the  $\Omega_\Lambda - \Omega_m$  plane is shown in Fig. 7. In the figure we have also indicated the parameter configuration for a flat Universe. A majority of the configurations seem to favour a non-flat Universe, with  $\Omega_{k0} \approx 0.0674$ . This could be a result of local variations in the gravitational field at small scales.

The posterior PDF of  $H_0$  is shown in Fig. 8, where we have included the resulting Gaussian distribution from the mean and variance of the sampled  $H_0$  values. This shows further discrepancy from the value of  $H_0 = 67 \text{ km/s/Mpc}$ , given by Planck, while the mean value we obtain is  $\hat{H}_0 = 70.1 \text{ km/s/Mpc}$ , with a corresponding standard deviation of  $\hat{\sigma}_{H_0} = 0.64 \text{ km/s/Mpc}$ .

With the fitted parameters, we can solve the background cosmology once again, and compute the resulting luminosity dis-



**Fig. 6.** Supernova data compared with the predicted luminosity distance obtained from simulations using Planck cosmology. Note that we use a logarithmic scale for the y-axis. **To be removed**



**Fig. 7.** Confidence regions of  $\Omega_{m0}$  and  $\Omega_{\Lambda0}$  obtained from fitting the supernova data. The best fit parameter values are shown in the plot, as well as the configurations that would give a flat Universe.

tance. The result is shown in Fig. 9, where the previous result from Fig. 6 is included for comparison purposes. The simulation with the best fit parameters yields a luminosity distance function that is much more consistent with data. However, there is still some discrepancy at lower redshift, but the associated data points in this regime have large uncertainties.


There are three important periods we have discussed, the times of matter-radiation equality, acceleration onset and matter-dark energy equality. The times when these incidents occur are listed in table 2.3.3, in terms of  $x$ , redshift  $z$  and cosmic time  $t$ . We also include the age of the Universe today,  $t_0 \equiv t(x = 0)$ , and the conformal time today,  $\eta_0/c$ .

	Matter-radiation equality $\Omega_m = \Omega_r$	Acceleration onset $\ddot{a} = 0$	Matter-dark energy equality $\Omega_m = \Omega_\Lambda$
$x$	-8.132	-0.487	-0.256
$z$	3400.3	0.627	0.291
$t$ [Gyr]	51 028.9 yr	7.750	10.371

Age of Universe today:  $t_0 = 13.848$  Gyr

Conformal time today:  $\eta_0/c = 46.287$  Gyr

**Table 1.** Important times during the evolution of the Universe, expressed in terms of  $x$ , redshift and cosmic time. In the last two rows we also present today's time values



**Fig. 8.** Posterior PDF of  $H_0$ . The histogram shows the sampled values, while the blue curve is the corresponding normal distribution obtained from the mean and variance of the data.



**Fig. 9.** The luminosity distance function obtained from the best fit values of  $C$  (blue line), compared with the result from the Planck cosmology (dashed green line), which is also shown in Fig. 6.

### 3. Milestone II

Having successfully implemented the background cosmology, the next step in developing our model is to include interactions between particles. After the Big Bang, the early universe was highly ionized. Due to Thompson scattering, photons were strongly coupled to baryons. As the Universe expanded, and the temperature dropped, neutral atoms were able to form, and the photons were able to escape from the plasma. These are the CMB photons we observe today. The period where neutral atoms formed is called recombination, and will be the main topic of this section. Our goal is to compute the number density of free electrons in the Universe, and use this to estimate when recombination occurred. The evolution of free electrons will affect both structure formation and the resulting power spectrum of the CMB photons, as we will study later.

We will use both the Saha and Peebles equation to compute the electron number density, and use this to compute the optical depth. From the optical depth, we will compute the so-called *visibility function*. In this section we will adapt natural units, where we set  $c = \hbar = k_B = 1$ . Additionally, we will make the assumption that Hydrogen is the only element present in the universe. Hence, we use  $Y_p = 0$ , rather than the value from Planck, given in Eq. (A.1), as we neglect Helium and heavier elements.

#### 3.1. Theory

##### 3.1.1. Optical depth and visibility function

A source that emits light with an intensity  $I_0$  is attenuated by a factor  $e^{-\tau(x)}$  as it travels through a medium, where  $\tau$  is the optical depth of the medium. In the early universe we have  $\tau \gg 1$ , and the universe is said to be optically thick. As the universe expands and recombination takes place, the universe becomes optically thin,  $\tau \ll 1$ , and light is able to escape the primordial plasma and reach us. Considering Thompson scattering only, the optical depth is defined in terms of the scale factor,  $a$ , as (Callin 2006, Eq. (5))

$$\tau(\eta) = \int_{\eta}^{\eta_0} d\eta' n_e \sigma_T a, \quad (33)$$

where  $n_e$  is the number density of free electrons,

$$\sigma_T = \frac{8\pi\alpha^2}{3m_e^2} = 6.6524587158 \cdot 10^{-29} \text{ m}^2 \quad (34)$$

is the Thompson cross-section and  $a$  is the scale factor.  $\alpha$  is the fine-structure constant and  $m_e$  is the electron mass. Using Eq.

(14), we can rewrite Eq. (33) as a differential equation

$$\frac{d\tau}{dx} = -\frac{n_e \sigma_T}{H}, \quad (35)$$

which we can solve numerically once we know  $n_e$ . The initial condition is  $\tau(x=0) = 0$ , as the universe is transparent today.

From the optical depth, we obtain the so-called *visibility function* (Callin 2006, Eq. (8))

$$\tilde{g}(x) = -\tau' e^{-\tau}, \quad (36)$$

which is normalized as

$$\int_{-\infty}^0 dx \tilde{g}(x) = 1. \quad (37)$$

The normalization means that  $\tilde{g}(x)$  is a probability distribution, and we may interpret it as the probability of an observed CMB photon today having experienced its last scattering at a time  $x$ . The visibility function is sharply peaked around the time when the photons decouple from the baryons, and is often referred to as the surface of last scattering. We may therefore use the peak of  $\tilde{g}(x)$  to estimate the time when decoupling took place in the early universe.

### 3.1.2. Electron density

The final thing we need to compute  $\tau$  and  $\tilde{g}$  is the number density of free electrons. To compute the evolution of free electrons, we have to consider the Boltzmann equation (Dodelson 2020, Eq. (3.19))

$$\frac{df}{dt} = C[f], \quad (38)$$

where  $f$  is the distribution function and  $C[f]$  is a collision term.

We will not consider the full Boltzmann equation, but an approximated version relevant for the recombination. Following the notation and definitions from (Dodelson 2020, Eq. (4.5)-(4.9)), the equation we consider is given as

$$a^{-3} \frac{d(n_i a^3)}{dt} = n_i^{(0)} n_2^{(0)} \langle \sigma v \rangle \left\{ \frac{n_3 n_4}{n_3^{(0)} n_4^{(0)}} - \frac{n_1 n_2}{n_1^{(0)} n_2^{(0)}} \right\}, \quad (39)$$

where  $n_i$  denotes the number density of a particle species  $i$ , with  $n_i^{(0)}$  referring to its value in chemical equilibrium, and  $\langle \sigma v \rangle$  is the thermally averaged cross-section. The only reaction we will consider for recombination is

$$e^- + p^+ \leftrightarrow H + \gamma. \quad (40)$$

For the photons we will assume  $n_\gamma = n_\gamma^{(0)}$ . From now on we use  $e$  and  $p$  as subscripts to denote free electrons and protons, respectively, and a subscript  $H$  to denote neutral Hydrogen.

Instead of computing  $n_e$  directly, we will compute the fractional electron density

$$X_e \equiv \frac{n_e}{n_e + n_H} \approx \frac{n_e}{n_b}, \quad (41)$$

where  $n_b$  is the total baryon density of the universe. Since the universe must be electrically neutral, we have  $n_e = n_p$ , and assuming no heavier elements than Hydrogen gives  $n_b \approx n_p + n_H$ . Neglecting the small mass difference between the proton and the Hydrogen, the baryon density can be written as

$$n_b \approx \frac{\rho_b}{m_H} = \frac{\Omega_{b0} \rho_{c0}}{m_H a^3}, \quad (42)$$

where  $\rho_{c0} \equiv \frac{H_0^2}{8\pi G}$  is the critical density of the universe today and  $m_H$  is the Hydrogen mass.

Before recombination occurs, the interaction rates greatly exceed the expansion rate of the universe, and for Eq. (39) to be valid we must have that

$$\frac{n_e n_p}{n_H} = \frac{n_e^{(0)} n_p^{(0)}}{n_H^{(0)}}. \quad (43)$$

In terms of  $X_e$ , this reduces to the Saha equation

$$\frac{X_e^2}{1 - X_e} = \frac{1}{n_b} \left( \frac{m_e T_b}{2\pi} \right)^{3/2} e^{-\epsilon_0/T_b}, \quad (44)$$

where  $T_b$  is the temperature of the baryons, and  $\epsilon_0$  is the Hydrogen ionization energy. The time evolution of  $T_b$  is governed by a differential equation coupled to  $X_e$ . However, we will assume that it follows the photon temperature,  $T_\gamma$ , evolving as

$$T_b = T_\gamma = T_{\text{CMB}0} e^{-x}. \quad (45)$$

At later times, Eq. (43) is no longer a valid approximation, and we have to solve Eq. (39). Additionally, we have to take processes related to atomic physics into account, so the differential equation we will consider is the Peebles equation

$$\frac{dX_e}{dx} = \frac{C_r(T_b)}{H} [\beta(T_b)(1 - X_e) - n_H \alpha^{(2)}(T_b) X_e^2], \quad (46)$$

where  $C_r(T_b)$ ,  $\beta(T_b)$  and  $\alpha^{(2)}(T_b)$  are quantities related to interaction effects, and are given by

$$C_r(T_b) = \frac{\Lambda_{2s \rightarrow 1s} + \Lambda_\alpha}{\Lambda_{2s \rightarrow 1s} + \Lambda_\alpha + \beta^{(2)}(T_b)}, \quad (47a)$$

$$\Lambda_{2s \rightarrow 1s} = 8.227 \text{ s}^{-1}, \quad (47b)$$

$$\Lambda_\alpha = H \frac{(3\epsilon_0)^3}{(8\pi)^2 n_{1s}}, \quad (47c)$$

$$n_{1s} = (1 - X_e) n_H, \quad (47d)$$

$$\beta^{(2)}(T_b) = \beta(T_b) e^{3\epsilon_0/4T_b}, \quad (47e)$$

$$\beta(T_b) = \alpha^{(2)}(T_b) \left( \frac{m_e T_b}{2\pi} \right)^{3/2} e^{-\epsilon_0/T_b}, \quad (47f)$$

$$\alpha^{(2)} = \frac{64\pi}{\sqrt{27}\pi} \frac{\alpha^2}{m_e^2} \sqrt{\frac{\epsilon_0}{T_b}} \phi_2(T_b), \quad (47g)$$

$$\phi_2(T_b) = 0.448 \ln(\epsilon_0/T_b). \quad (47h)$$

The main reason behind these additional equations is that Hydrogen production is inefficient at  $T_b \approx \epsilon_0$ . Direct recombination to the Hydrogen ground state is likely to produce a photon with energy greater than  $\epsilon_0$ , which will ionize another nearby Hydrogen atom, resulting in no net Hydrogen production. Recombination is achieved when an electron and proton combine to an excited Hydrogen atom, followed by the atom's decay into the ground state, which is a slow process.

In Sect. 3.2.2 we discuss how combine the Saha and Peebles equation to compute  $X_e$ .

### 3.1.3. Sound Horizon at decoupling

Before recombination happens, baryons and photons are tightly coupled. For this reason, the baryons and photons behave as if



they were a single fluid, and the sound speed of this fluid is given by (Dodelson 2020, Eq. (9.21)) (where we use  $R \rightarrow 1/R$ )

$$c_s(x) = \frac{c}{\sqrt{3}} \sqrt{\frac{R(x)}{1+R(x)}}, \quad R(x) = \frac{4\Omega_\gamma(x)}{3\Omega_b(x)}. \quad (48)$$

The total co-moving distance a sound wave in this plasma could have travelled since the Big Bang is known as the sound-horizon, which is given as

$$s(x) = \int_{-\infty}^x \frac{dx' c_s}{\mathcal{H}}. \quad (49)$$

From this we can compute the sound horizon at decoupling,  $r_s = s(x_{\text{rec}})$ , which is an important quantity of the CMB. Thus, we have an additional ODE to solve,

$$\frac{ds(x)}{dx} = \frac{c_s}{\mathcal{H}}, \quad (50)$$

with  $s(x_{\text{ini}}) = \frac{c_s(x_{\text{ini}})}{\mathcal{H}(x_{\text{ini}})}$  as the initial condition, following the same reasoning as we did for  $\eta'(x)$  in Eq. (15).

### 3.2. Implementation details

To compute  $X_e$ , we use the Saha equation initially, as it is a good approximation at early times when  $X_e \approx 1$ . This is also the regime where the Peebles equation is unstable, and we therefore consider the Saha equation for  $X_e > X_e^{\text{tol}}$ . Once we reach  $X_e < X_e^{\text{tol}}$  we use the final value from the Saha equation as our initial condition to solve the Peebles equation. The Peebles equation is then used all the way to today, at  $x = 0$ . For this report, we choose  $X_e^{\text{tol}} = 0.99$ . For comparison with the Saha equation alone, we set  $X_e^{\text{tol}} = 10^{-6}$ . Once  $X_e$  is computed we get  $n_e(x)$  from Eq. (41).

#### 3.2.1. Solving the Saha Equation

Solving the Saha equation is done by solving a quadratic formula for  $X_e$ . At early times, however, the RHS of Eq. (44) will be enormous, and may cause numerical errors when solving the quadratic formula. To avoid this, we use the first order approximation  $\sqrt{1+x} \approx 1 + \frac{x}{2}$  for  $|x| \ll 1$  at early times. The Saha equation is thus implemented as

$$X_e = \begin{cases} 1, & y > 10^7, \\ \frac{y}{2} \left[ -1 + \sqrt{1 + 4/y} \right], & y \leq 10^7, \end{cases} \quad (51)$$

where  $y$  refers to the RHS of Eq. (44). Since  $X_e$  is strictly positive, we have omitted the negative solution. The exact value of  $10^7$  is chosen to ensure both  $X_e \not\approx 1$ , and  $X_e \not\ll X_e^{\text{tol}}$  when the quadratic formula is to be used.

#### 3.2.2. Solving the Peebles Equation

To solve Eq. (46) numerically, we follow the same procedure as we did for  $\eta(x)$ , but for the initial condition we use the final value of  $X_e$  that we obtained from the Saha equation.

At late times, when the baryon temperature gets low, the exponential term in Eq. (47e) for  $\beta^{(2)}(T_b)$  become sufficiently large to yield an overflow. However, this is also where  $\beta(T_b) \rightarrow 0$ , due to its exponential factor (Eq. (47f)). This exponential factor causes  $\beta^{(2)}(T_b) \rightarrow 0$  at late times. To avoid overflow, we implement the equation for  $\beta^{(2)}(T_b)$  as

$$\beta^{(2)}(T_b) = \begin{cases} 0, & \epsilon_0/T_b > 200, \\ \beta(T_b)e^{3\epsilon_0/4T_b}, & \epsilon_0/T_b \leq 200. \end{cases} \quad (52)$$

#### 3.2.3. Optical depth and visibility function

With  $n_e$  we can then solve Eq. (35) for  $\tau(x)$  with the aforementioned initial condition of  $\tau(x=0) = 0$ . We therefore integrate backwards, starting from  $x = 0$ . The visibility function is now easily obtained, as  $\tau'(x)$  is given analytically by Eq. (35). From this, we get immediately  $\tilde{g}(x)$ .

We also need the second derivative of  $\tau$ , as well as the first two derivatives of  $\tilde{g}$ . For  $\tau''(x)$ , we compute it from numerically differentiating the  $\tau'(x)$  data.<sup>1</sup> We use this to compute

$$\tilde{g}'(x) = \left( \tau'(x)^2 - \tau''(x) \right) e^{-\tau(x)}, \quad (53)$$

and obtain  $\tilde{g}''(x)$  by numerically differentiating  $\tilde{g}'(x)$ . This is done to avoid potential errors that may occur when numerically computing the second derivative of  $\tilde{g}(x)$ , if  $\tilde{g}(x)$  is somewhat ill-behaved at certain times.<sup>2</sup>

#### 3.2.4. Determining the time of recombination and decoupling

To estimate the times when recombination takes place, we define this as

$$X_e(x = x_{\text{recombination}}) = 0.1, \quad (54)$$

where the value  $X_e = 0.1$  is chosen arbitrarily.

For decoupling, we find  $x_{\text{decoupling}}$  as the point where  $\tilde{g}'(x_{\text{decoupling}}) = 0$ . However, we limit the time values to  $x \in [x_0 \pm 0.1]$ , where  $x_0$  is defined by  $\tau(x_0) = 1$ . This is done to avoid  $x$  values where  $\tilde{g}(x) = 0$ .

### 3.3. Results

#### 3.3.1. Electron fraction

The evolution of the electron fraction with time is shown in Fig. 10. Here, we have included the solution obtained through the Saha equation alone for comparison. In the figure we also show the time periods when recombination occurs according to the two solutions, i.e. when  $X_e = 0.1$ . The time of decoupling has been omitted, since it is very close to the recombination time, as shown in table 2.

We see that the Peebles equation predicts the production of neutral Hydrogen to take much longer time than the Saha equation. Recombination therefore occurs at a much lower temperature than the binding energy of Hydrogen, as expected. Towards  $x = 0$  the Peebles equation give  $X_e \ll 1$ , but we never reach  $X_e = 0$  entirely. From our solution we find that the current value is  $X_e(x=0) \approx 2.026 \cdot 10^{-4}$ . On the other hand, the Saha equation reaches  $X_e = 0$  before  $x = 0$  is reached, predicting now free electrons in the universe.

#### 3.3.2. Optical depth and visibility function

The optical depth, and its first two derivatives is shown in Fig. 11. We see that the three quantities rapidly drop by several orders of magnitude near  $x = -7$ , around the time of recombination. Looking at Fig. 12 we see that the rapidly varying optical depth results in a sharply peaked  $\tilde{g}$ .

<sup>1</sup> Couldn't we just compute it analytically? I get a tiny "bump" in  $\tau''$  near  $X_e^{\text{tol}}$ , and I don't know if this will cause problems later on, and therefore didn't want to spend too much time on it if wasn't needed.

<sup>2</sup> I don't know if this is useful, or helps, at all. I don't know if we care about the derivatives at  $x$ -values where this might be an issue.





**Fig. 10.** Free electron fraction computed from the Saha equation only (dashed green curve) and from the Peebles equation (solid blue curve), where the Saha equation was used at early times, until  $X_e < 0.99$ . The recombination times is shown for both solutions.



**Fig. 11.** The optical depth,  $\tau(x)$  and its two first derivatives with respect to  $x$ .

### 3.3.3. Times of recombination and decoupling

The period of recombination and decoupling, as computed from the Peebles equation, is given in table 2. We see that the photons and baryons decouple later than recombination occurs, as is expected. This does depend on how we define the two periods, and is thus not a very conclusive result on its own. Using a higher value of  $X_e$  for instance would yield a larger separation between the two events.

**Table 2.** Times when decoupling and recombination occurs, computed from the Peebles equation.

Peebles	$x$	$z$	$t$ [yr]	$r_s$ [Mpc]
Decoupling	-6.98534	1079.7	377 782	145.3
Recombination	-6.98549	1079.8	377 690	



**Fig. 12.** The visibility function,  $\tilde{g}(x)$  (solid curve), and its derivatives,  $\tilde{g}'(x)/10$  (dashed curve) and  $\tilde{g}''(x)/300$  (dotted curve). The derivatives have been scaled in order to view them all in the same plot.

**Table 3.** Times when decoupling and recombination occurs, computed from the Saha equation.

Saha	$x$	$z$	$t$ [yr]
Decoupling	-7.15485	1279.3	283 611
Recombination	-7.14035	1260.9	290 686

## 4. Milestone III

So far we have only considered the flat FLRW metric for a homogeneous Universe, and used this to study the recombination history of the Universe. In this section we will consider small perturbations to the background, and thereby introduce inhomogeneities. We will consider perturbations to the FLRW metric and the matter contents of the Universe, where we limit ourselves to photons, baryons and CDM. We will restrict ourselves to linear perturbations only, which allows us to derive good approximations for the Boltzmann and Einstein equations. From this we are able to study the evolution of structure in the early Universe, which we later will connect to statistical observables in the Universe. This means that we ignore contributions from neutrinos and polarization. This means that we set  $N_{\text{eff}} = 0$  when we compute relevant quantities from Sect. 2 and Sect. 3 that we will need in this section.

### 4.1. Theory

In this section we present the equations governing the evolution of the perturbed quantities. We will only present the relevant equations here, which are all obtained from Dodelson (2020) and Callin (2006), unless otherwise stated. We refer to Dodelson for a detailed derivation of the equations. For the notation, we adapt that of Callin. Throughout this section, we will consider perturbations in Fourier space.

#### 4.1.1. Perturbation quantities

For the perturbed metric we consider the Newtonian gauge, and write it as

$$g_{\mu\nu} = \begin{pmatrix} -(1 + 2\Psi) & 0 \\ 0 & a^2\delta_{ij}(1 + 2\Phi) \end{pmatrix}, \quad (55)$$

where we have introduced the scalar perturbations  $\Psi$  and  $\Phi$ , which are functions of both position and time. For  $\Psi = \Phi = 0$  we obtain the FLRW metric. Photon perturbations are defined in terms of the relative temperature variations,  $\Theta \equiv \delta T/T$ , via

$$T(\mathbf{k}, \mu, \eta) = T^{(0)}(\eta) [1 + \Theta(\mathbf{k}, \mu, \eta)], \quad (56)$$

where  $\mathbf{k}$  is the Fourier transformed variable corresponding to position  $\mathbf{x}$ , and  $\mu \equiv \frac{\mathbf{k} \cdot \mathbf{p}}{kp}$ . The temperature perturbations only depend on the photon momentum in terms of their directions, and we therefore expand  $\Theta$  in terms of multipoles as

$$\Theta_\ell = \frac{i^\ell}{2} \int_{-1}^1 \mathcal{P}_\ell(\mu) \Theta(\mu) d\mu \Leftrightarrow \Theta(\mu) = \sum_{\ell=0}^{\infty} \frac{2\ell+1}{i^\ell} \Theta_\ell \mathcal{P}_\ell(\mu), \quad (57)$$

where  $\mathcal{P}_\ell(\mu)$  are Legendre polynomials. The first three multipoles are the monopole,  $\Theta_0$ , the dipole,  $\Theta_1$  and the quadrupole,  $\Theta_2$ . The monopole represents the background temperature of the CMB and the dipole represents our relative velocity with respect to the CMB.

We denote density perturbations of CDM and baryons as  $\delta_{\text{CDM}}$  and  $\delta_b$ , and their velocity perturbations as  $v_{\text{CDM}}$  and  $v_b$ . From the perturbed Einstein equations one can show that  $\delta_\gamma = 4\Theta_0$  and  $v_\gamma = -3\Theta_1$  for the photons.

#### 4.1.2. Perturbation equations

The system of ODEs we will solve for the evolution of the perturbed quantities are given in (Callin 2006, Eq. (22)), where we neglect neutrinos and polarization and set  $\Theta_p = \mathcal{N} = 0$ .

$$\Theta'_0 = -\frac{k}{\mathcal{H}} \Theta_1 - \Phi', \quad (58a)$$

$$\Theta'_1 = \frac{k}{3\mathcal{H}} \Theta_0 - \frac{2k}{3\mathcal{H}} \Theta_2 + \frac{k}{3\mathcal{H}} \Psi + \tau' \left[ \Theta_1 + \frac{1}{3} v_b \right], \quad (58b)$$

$$\Theta'_\ell = \frac{\ell k}{(2\ell+1)\mathcal{H}} \Theta_{\ell-1} - \frac{(\ell+1)k}{(2\ell+1)\mathcal{H}} \Theta_{\ell+1} + \tau' \left[ \Theta_\ell - \frac{1}{10} \Pi \delta_{\ell,2} \right], \quad 2 \leq \ell < \ell_{\text{max}}, \quad (58c)$$

$$\Theta'_\ell = \frac{k}{\mathcal{H}} \Theta_{\ell-1} - c \frac{(\ell+1)}{\mathcal{H}\eta(x)} \Theta_\ell + \tau' \Theta_\ell, \quad \ell = \ell_{\text{max}}, \quad (58d)$$

$$\delta'_{\text{CDM}} = \frac{k}{\mathcal{H}} v_{\text{CDM}} - 3\Phi', \quad (58e)$$

$$v'_{\text{CDM}} = -v_{\text{CDM}} \frac{k}{\mathcal{H}} \Psi, \quad (58f)$$

$$\delta'_b = \frac{k}{\mathcal{H}} v_b - 3\Phi', \quad (58g)$$

$$v'_b = -v_b - \frac{k}{\mathcal{H}} \Psi + \tau' R(3\Theta_1 + v_b), \quad (58h)$$

$$\Phi' = -\Psi - \frac{k^2}{3\mathcal{H}^2} \Phi + \frac{H_0^2}{2\mathcal{H}^2} [\Omega_{\text{CDM}0} a^{-1} \delta_{\text{CDM}} + \Omega_{b0} a^{-1} \delta_b + 4\Omega_{\gamma0} a^{-2} \Theta_0], \quad (58i)$$

$$\Psi = -\Phi - \frac{12H_0^2}{k^2 a^2} \Omega_{\gamma0} \Theta_2. \quad (58j)$$

We have  $\Pi = \Theta_2$  due to  $\Theta_p = 0$ , and have written  $\Pi$  for clarity. The equation for  $\Psi$  is an algebraic equation, and its value is inserted in the other equations. Noting that  $\Phi'$  is the only derivative of a perturbed quantity found on the RHS of Eqs. (58), we solve for it first, and insert this value into the other equations.

In Sect. 5 we discuss the main topic of this report, which is to compute the power spectrum,  $C_\ell$ , which involves integrating over  $\Theta_\ell$ . Since we are interested in probing the power spectrum at scales where  $\ell > 1000$  (Mention how  $\ell$  relates to scale), we need a vast number of photon multipoles. From the ODE, we see that  $\Theta'_\ell$  is recursive, and computing  $\Theta_\ell$  directly from the ODEs will thus require us to solve thousands of ODEs. This is a highly tedious and computationally expensive process. However, as we will explain in Sect. 5.1.3, by exploiting the line-of-sight (LOS) integration technique Seljak & Zaldarriaga (1996), we can compute any order of  $\Theta_\ell$  for  $\ell > 2$  from  $\Theta_0$  and  $\Theta_2$ . Thus, we only need to use the ODEs to accurately compute  $\Theta_0$  and  $\Theta_2$ . For this, we need to truncate the series at some point. Truncating the series by setting  $\Theta_{\ell_{\text{max}}} = 0$  would lead to numerical errors propagating down to  $\Theta_2$ . Eq. (58d) comes from assuming  $\Theta_\ell(k, \eta) \sim j_\ell(k\eta)$  at high  $\ell$ , where  $j_\ell(k\eta)$  is the spherical Bessel function, and using the recurrence relation for  $j_\ell$  (see Callin 2006, Eq. (46)-(49)). By an appropriate choice of  $\ell_{\text{max}}$ , we greatly reduce the number of ODEs needed to accurately compute  $C_\ell$ .

At early times, photons and electrons are tightly coupled, and  $\tau$  is therefore very high. In this regime, Eq. (58) becomes numerically unstable, so we have to resort to approximations, which we discuss in the next section.

#### 4.1.3. Initial conditions

We will use adiabatic initial conditions, and follow the convention used by Winther (2023), with  $f_\nu = 0$  as we omit neutrinos<sup>3</sup>. The set of ODEs we solve are linear, so we can choose a particular normalization, which we adjust for when computing the Power spectrum later on. The initial conditions are

$$\Psi = -\frac{2}{3}, \quad (59a)$$

$$\Phi = -\Psi, \quad (59b)$$

$$\delta_{\text{CDM}} = \delta_b = -\frac{3}{2} \Psi, \quad (59c)$$

$$v_{\text{CDM}} = v_b = -\frac{k}{2\mathcal{H}} \Psi, \quad (59d)$$

$$\Theta_0 = -\frac{1}{2} \Psi, \quad (59e)$$

$$\Theta_1 = \frac{k}{6\mathcal{H}} \Psi. \quad (59f)$$

Initially,  $\tau$  is very large, so photons and baryons behave as a single fluid, and is referred to as the *Tight coupling regime*. This implies that temperature gradients are washed out, as the system is in thermodynamic equilibrium. Since  $\tau'$  is large at early times, it means that the terms proportional to  $\tau'$  in Eq. (58) can be neglected. This implies that  $\Theta_\ell = 0$  for  $\ell \geq 2$ . However, we need the lowest non-zero contribution from these multipoles when numerically integrating the system of ODEs. By expanding in terms of  $k/\mathcal{H}\tau' \ll 1$  and using the initial conditions, one can obtain the following expression for the higher order multipoles

$$\Theta_2 = -\frac{20k}{45\mathcal{H}\tau'} \Theta_1, \quad (60a)$$

$$\Theta_\ell = -\frac{\ell}{2\ell+1} \frac{k}{\mathcal{H}\tau'} \Theta_{\ell-1}. \quad (60b)$$

<sup>3</sup> Note that Winther has a sign error in his expression for the initial condition for  $\Psi$

These are the expressions we will use to compute the multipoles during tight coupling. Note that we  $\Theta_1 \simeq -v_b/3$  in this regime. To avoid numerical instability when multiplying this with  $\tau'$  in Eq. (58b) we derive a modified expression for  $\Theta'_1$  and  $v'_b$  in Sect. 4.2.1. Furthermore, we discuss in that section how we determine when to stop using the tight coupling approximation.

#### 4.1.4. Evolution of different modes

The evolution of a single mode is governed by Eq. (58). As the horizon grows, causal physics affects perturbations on increasingly larger scales. Initially we have  $k\eta \ll 1$ , and all modes are outside the horizon. As  $\eta$  grows, more and more modes enter the horizon. The epoch in which a mode enters the horizon affects its evolution. To study the evolution of perturbations, we focus on three different modes: one short-wavelength mode that enters the horizon during radiation domination, one large-wavelength mode that enters the horizon during matter domination, and one mode of intermediate scale.

Defining  $x_{\text{entry}}$  as  $k\eta(x_{\text{entry}}) = 1$ , we choose the wavenumbers

$$k \in \{0.001, 0.03, 0.3\}/\text{Mpc},$$

$$x_{\text{enter}} \in \{-5.182, -9.726, -12.098\}, \quad (61)$$

where the first and last element of  $x_{\text{enter}}$  correspond to modes entering the horizon when  $\Omega_m \approx 0.95$  and  $\Omega_\gamma \approx 0.95$ , respectively. The intermediate mode will enter when  $\Omega_m \approx 0.25$  and  $\Omega_\gamma \approx 0.25$ . We postpone the details regarding evolution of these modes to Sect. 4.3.

## 4.2. Implementation details

### 4.2.1. Tight coupling regime

During tight coupling, Eq. (62c) needs to be modified to avoid numerically instability at early times. Following Callin, the tight coupling regime is computed with

$$\varrho q = -[(1-R)\tau' + (1+R)\tau''](3\Theta_1 + v_b) - \frac{k}{\mathcal{H}}\Psi + (1 - \frac{\mathcal{H}'}{\mathcal{H}})\frac{k}{\mathcal{H}}(-\Theta_0 + 2\Theta_2) - \frac{k}{\mathcal{H}}\Theta'_0, \quad (62a)$$

$$v'_b = \frac{1}{1+R} \left[ -v_b - \frac{k}{\mathcal{H}}\Psi + R(q + \frac{k}{\mathcal{H}}[-\Theta_0 + 2\Theta_2 - \Psi]) \right], \quad (62b)$$

$$\Theta'_1 = \frac{1}{3}(q - v'_b), \quad (62c)$$

where we have introduced the parameter

$$\varrho = (1+R)\tau' + \frac{\mathcal{H}'}{\mathcal{H}} - 1. \quad (63)$$

In the tight coupling regime we use the initial conditions given in Eq. (59). We then solve the ODEs given in Eq. (58), but compute  $\Theta'_1$  and  $v'_b$  in the end by using Eq. (62). In this regime we use Eqs. Eq. (60a) and (60b) for  $\Theta_2$  and  $\Theta_\ell$ , respectively.

Next, we need to determine when the tight coupling approximation is valid. We use the same conditions as Callin, namely that the approximation holds as long as  $|\tau'| > 10$  and  $|k/\mathcal{H}\tau'| < 1/10$ , and that it should not be used after the start of recombination. For the latter condition, we choose  $x < -8.3$  as the time when recombination starts. If any of these three conditions fails, we switch to the full ODEs. We then use the final results from the tight coupling regime as initial conditions for the ODEs in Eq. (58), and integrate up to  $x = 0$ .

### 4.2.2. Integration of ODEs

When solving the perturbation ODEs we integrate across  $x$  for each mode  $k$ . We choose  $N_k = 200$  logarithmically spaced values of  $k \in [k_{\min}, k_{\max}]$ , where  $k_{\min} = 5 \cdot 10^{-5}/\text{Mpc}$  and  $k_{\max} = 0.3/\text{Mpc}$ . For the photon multipoles, we set  $\ell_{\max} = 7$ .

Once the ODEs have been solved for all combinations of  $x$  and  $k$ , we make splines of the integrated perturbation quantities, except for  $\Theta_\ell$  for  $\ell > 2$ , as this will be solved using the LOS integration.

## 4.3. Results

In this section we present plots showing the evolution of some perturbation quantities. For the monopole and dipole, we express plot these in terms of  $\delta_\gamma$  and  $v_\gamma$ , respectively. For easier comparison between evolution of baryons, photons and CDM, we plot CDM and baryons together in one figure, and baryons and photons together in another figure. In both cases we include all three modes in the same figure. This approach prevents cluttering, and helps us understand the interaction between the three constituents, as baryons interact with both CDM and photons.

In Figs. 13-17 we indicate the time of matter-radiation equality by a dotted black vertical line. We stress that matter-radiation equality occurs at  $x_{\text{eq}} = -8.6577$ , as opposed to the time given in table 2.3.3, since  $\mathcal{N}_{\text{eff}} = 0$  now. The time of decoupling and recombination, however, is less affected, occurring at  $x \sim -6.98$  as before. We also mark the horizon entry time for each mode, given in Eq. (61), shown as dash-dotted lines. For brevity, we will refer to modes with large  $k$  as *short modes* and modes with small  $k$  as *long modes*, as they are related to small and large scales, respectively.

We begin by studying the density evolutions in Sect. 4.3.1, and spend a considerable time interpreting these results qualitatively. This discussion provides a simpler way of understanding the mechanism behind the remaining perturbation quantities.


### 4.3.1. Density perturbations

The absolute value of the density evolution of baryons, CDM and photons is shown in Fig. 13, where the upper panel shows  $\delta_{\text{CDM}}$  and  $\delta_b$  and the lower panel shows  $\delta_b$  and  $\delta_\gamma$ . During the tight coupling regime, baryons and CDM follow each other closely, but  $\delta_b$  deviates noticeably from  $\delta_{\text{CDM}}$  during recombination for the shortest mode. As time evolves, we observe formation of overdense regions of CDM undergoing gravitational collapse, taking place around the time when the modes enter the horizon. This attracts baryons, which fall into the CDM potential wells. As baryons collapse, pressure from photons increases and pushes them outwards again. This phenomenon is apparent in the lower panel, where  $\delta_b$  and  $\delta_\gamma$  follow each other as a single fluid before decoupling. This process repeats for the two shorter modes, which have entered the horizon before recombination begins. When decoupling takes place, baryons can collapse into the gravitational wells set up by CDM. Consequently,  $\delta_b$  follows  $\delta_{\text{CDM}}$  at late times. The photons can move freely after decoupling, and their temperature remains approximately unaffected towards  $x = 0$ . There are some fluctuations in  $\delta_\gamma$  at late times, which may be caused by variations in gravitational potentials.

In general, we see more oscillations in  $\delta_b$  and  $\delta_\gamma$  for smaller modes. The large mode enters the horizon long after decoupling, and the baryon overdensity is unaffected by photons when collapse begins. Neglecting terms with factors of  $k$  in Eq. (58), we

get  $\delta_b = \delta_{\text{CDM}}$  at all times, since they have identical initial conditions.

The value of  $\delta_\gamma$  after decoupling differs between the three modes, where shorter modes have a lower final temperature. Decoupled photons are redshifted as they move out of gravitational potential wells. On small scales, where CDM have more time to cluster before decoupling, the escaping photons become more redshifted. Additionally, the actual evolution of the potential wells before decoupling is an important factor in the final photon temperatures. For the shortest mode, we see that  $\delta_{\text{CDM}}$  grows differently right after horizon entry, compared to after  $x_{\text{eq}}$ . This may be a result of matter density starting to affect the expansion rate of the Universe. We will discuss the effect of expansion in more detail in Sect. 4.3.3. We see an example of how expansion affects the overdensities, as there is a slight decrease in  $\delta_\gamma$  near  $x = 0$  with  $\delta_{\text{CDM}}$  and  $\delta_b$  flattening out. This occurs when the Universe becomes dominated by dark energy, where accelerated expansion is more effective than the accretion of matter.



**Fig. 13.** Evolution of  $\delta$  for modes with three different wavenumbers. Upper panel: Comparing  $\delta_{\text{CDM}}$  (solid line) and  $\delta_b$  (dashed line). Lower panel: Comparing  $\delta_b$  (solid line) and  $\delta_\gamma = 4\Theta_0$  (dashed line).

#### 4.3.2. Velocity perturbations

The evolution of the velocities is shown in Fig. 14, where we compare baryons and CDM in the upper panel, and baryons and photons in the lower panel. Note that we have used a linear scale for the lower panel to reduce clutter.

In Fig. 13, the oscillatory behaviour of baryons for short modes is also apparent in their velocities, where the velocity extrema coincide with density minima, as expected. Similar behaviour is observed for CDM velocities, with both baryons and CDM coinciding at early and late times, and decoupling effects in the middle for the two shorter modes. For the longest mode,

$v_{\text{CDM}}$  and  $v_b$  follow the same argument as their densities, resulting in  $v_{\text{CDM}} = v_b$ .

For the shortest mode,  $v_{\text{CDM}}$  is no longer a strictly increasing function at early times. Around  $x = -10$ ,  $v_{\text{CDM}}$  declines and becomes smaller than the intermediate mode around  $x_{\text{eq}}$ <sup>4</sup>. This shift persists all the way towards  $x = 0$ . During tight coupling we have  $v_b \sim v_\gamma$ , and near decoupling, at  $x \sim -7$ , we see that the photons decouple from the plasma. After that, the photons undergo damped oscillations, with a higher frequency for shorter modes.



**Fig. 14.** Evolution of velocities for three different wavenumbers. Upper panel:  $v_{\text{CDM}}$  (solid line) compared to  $v_b$  (dashed line). Lower panel:  $v_b$  (solid line) compared to  $v_\gamma$  (dashed line).


#### 4.3.3. Potential perturbations

The evolution of  $\Phi$  is illustrated in Fig. 15. This analysis is based on the discussion presented in (Dodelson 2020, Ch. 8). During both matter and radiation domination,  $\Phi$  can be well-approximated as being constant, with a transition period in between. As the shortest mode enters the horizon when  $x \ll x_{\text{eq}}$ , the potential decays and begins to oscillate, leading to the damped oscillations observed for photons and baryons in Fig. 13. During matter domination, for a mode that has entered the horizon, we get a constant  $\Phi$ , as the matter accretion is balanced by the Universe's expansion. The constancy of  $\Phi$  explains why all modes in Fig. 13 grow at the same rate at late times, long after decoupling.

From the transition period mentioned earlier, we can compare the changes in the growth of  $\delta_{\text{CDM}}$  at points where  $\Phi$  exhibits irregular behaviour. Most notably, the minima in  $\Phi$  at

<sup>4</sup> The reason for this behaviour has confused great minds for several light weeks. Their best estimate to this day is that it's expansion related

around  $x \sim -10$  roughly corresponds to the change in  $\delta_{\text{CDM}}$  for the short mode at the same  $x$ .




**Fig. 15.** Evolution of  $\Phi$ .


The sum of the two metric potentials,  $\Phi + \Psi$ , is plotted in Fig. 16. This quantity corresponds to the second term on the right-hand side of Eq. (58j) and is proportional to the quadrupole moment,  $\Theta_2$ , which is depicted in Fig. 17. Specifically, we have  $\Phi + \Psi \propto -\frac{\Theta_2}{k^2} e^{-2x}$ . Since  $\Theta_2(x \ll x_{\text{eq}}) \approx 0$ , the sum of the potentials is initially zero. Although the proportionality constant is very small, for larger  $x$ , late in the matter dominated era, there is a narrow range where both  $\Theta_2$  contributes and the exponent is large enough to give a non-zero contribution.

For the evolution of  $\Theta_2$ , we first begin to see growing oscillations in the short mode around  $x = x_{\text{eq}}$ . This growth is abruptly suppressed once decoupling happens, where we also had an abrupt decrease of  $v_\gamma$  in Fig. 14. Similar behaviour is seen for the remaining two modes. For these modes, oscillations start around the time of decoupling, and continues with an amplitude related to the amplitude of the oscillations of  $v_\gamma$ . The largest mode is also responsible for the largest value of  $\Theta_2$  at  $x = 0$ .

The evolution of  $\Theta_2$  is characterized by growing oscillations in the short mode around  $x = x_{\text{eq}}$ , which are suddenly suppressed upon decoupling. This abrupt decrease is also seen as a decrease in  $v_\gamma$  in Fig. 14. Similar behaviour is also observed for the remaining two modes. Oscillations for these modes start around decoupling and persist with an amplitude that is related to the amplitude of  $v_\gamma$  oscillations. The largest mode exhibits the highest  $\Theta_2$  value at  $x = 0$ .



**Fig. 16.** Evolution of the combined metric perturbations,  $\Phi + \Psi$



**Fig. 17.** Evolution of the quadrupole,  $\Theta_2$ . At  $x < -10$ , there is no apparent contribution of this quantity.

## 5. Milestone IV

We now have all the necessary constituents needed to compute the main quantity of interest in the report, the CMB power spectrum. By expanding the temperature field in spherical harmonics, we will be able to predict the angular CMB power spectrum to the observed values, and from that extract information about the underlying physics of the early Universe.

We are concerned with both the angular power spectrum of the CMB, and the matter power spectrum. When only "power spectrum" is written, we are referring to the former, and other power spectra will be referred to with appropriate prefixes.

### 5.1. Theory

#### 5.1.1. Angular power spectrum

The temperature fluctuations can be expanded in spherical harmonics as (check summation indices and momentum vector.)

$$\Theta(\mathbf{x}, \hat{\mathbf{p}}, t) = \sum_{\ell=1}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m}(\mathbf{x}, t) Y_{\ell m}(\hat{\mathbf{p}}), \quad (64)$$

where the amplitudes,  $a_{\ell m}$ , contain all the information about the temperature field, and are to be evaluated at our position in the Universe at  $t = t_0$ . However, we are unable to make prediction about particular values of these amplitudes, as inflation is a stochastic process. However, we exploit the fact that inflation predicts  $\Theta$  to be very close to Gaussian, so each  $a_{\ell m}$  is therefore drawn from a normal distribution with zero mean and a variance given as

$$\langle a_{\ell m} a_{\ell' m'} \rangle = \delta_{\ell \ell'} \delta_{m m'} C_{\ell}. \quad (65)$$

**Write something about statistics?**

We follow the approach of (Dodelson 2020, Ch. 9.5), leaving out the details of the derivation. We now omit factors of  $t$  in the expressions, as we are only concerned with the value at  $t = t_0$ , and obtain

$$C_{\ell} = 4\pi \int \mathcal{P}_{\mathcal{R}}(k) |\Theta_{\ell}^{\text{code}}(k)|^2 \frac{dk}{k}, \quad (66)$$

where we have used  $\Theta_{\ell}(\mathbf{k}) = \Theta_{\ell}^{\text{code}}(k) \mathcal{R}_{\text{ini}}$ , where  $\mathcal{R}_{\text{ini}} \propto \Psi$  comes from inflation. This rescaling accounts for the fact that we have set  $\Psi = -2/3$  as our initial condition when solving the perturbation ODEs numerically.  $\mathcal{P}_{\mathcal{R}}(k)$  is related to the primordial power spectrum,  $P(k)$ , via

$$P(k) = \frac{2\pi^2}{k^3} \mathcal{P}_{\mathcal{R}}(k), \quad (67)$$



where  $\langle \mathcal{R}_{\text{ini}}(\mathbf{k}) \mathcal{R}_{\text{ini}}^*(\mathbf{k}') \rangle = (2\pi^2) \delta(\mathbf{k} - \mathbf{k}') P(k)$ . To compute  $C_\ell$ , the last thing we need is an expression for  $\mathcal{P}_{\mathcal{R}}(k)$ . We get an expression for this from the fact that most inflationary models predict a so-called Harrison-Zel'dovich spectrum (cite), where

$$\mathcal{P}_{\mathcal{R}}(k) = A_s \left( \frac{k}{k_{\text{pivot}}} \right)^{n_s-1}, \quad (68)$$

where  $A_s$  is the amplitude of something,  $k_{\text{pivot}}$  is the something and  $n_s$  is the spectral index. **Comment about their meaning.**

Thus, the final expression we want to compute is

$$C_\ell = 4\pi \int_0^\infty A_s \left( \frac{k}{k_{\text{pivot}}} \right)^{n_s-1} \Theta_\ell^2(k) \frac{dk}{k}. \quad (69)$$

### 5.1.2. Matter power spectrum

From the Fourier components of the matter overdensity field, we can compute the matter power-spectrum

$$P_L(k, x) = |\Delta_M(k, x)|^2, \quad \text{where} \quad \Delta_M(k, x) \equiv \frac{k^2 \Phi(k, x)}{\frac{3}{2} \Omega_{m0} a^{-1} H_0^2}. \quad (70)$$

Once again, we have to take rescale  $\Delta_M$  to account for our choice of initial conditions. Taking the value today, we thus want to compute

$$P_L(k) = |\Delta_M^{\text{code}}(k)|^2 P(k) = |\Delta_M^{\text{code}}(k)|^2 \frac{2\pi^2}{k^3} \left( \frac{k}{k_{\text{pivot}}} \right)^{n_s-1}. \quad (71)$$

### 5.1.3. Line-of-sight integration

As mentioned in Sect. 4.1.2, we compute higher order multipoles from the LOS approach.

$$\Theta_\ell(k, x=0) = \int_{-\infty}^0 \tilde{S}(k, x) j_\ell[k(\eta_0 - \eta(x))] dx, \quad (72)$$

where  $j_\ell(k(\eta_0 - \eta(x)))$  is the spherical Bessel function.  $\tilde{S}(k, x)$  is the source function, which is given by

$$\begin{aligned} \tilde{S}(k, x) = & \tilde{g} \left[ \Theta_0 + \Psi + \frac{1}{4} \Pi \right] + e^{-\tau} [\Psi' - \Phi'] \\ & - \frac{1}{k} \frac{d}{dx} (\mathcal{H} \tilde{g} v_b) + \frac{3}{4k^2} \frac{d}{dx} \left[ \mathcal{H} \frac{d}{dx} (\mathcal{H} \tilde{g} \Pi) \right], \end{aligned} \quad (73)$$

where  $\Pi = \Theta_2$ .

From the source function alone we may get some insight into how different physical effects will affect the power spectrum. In Sect. 5.3.1 we will present a more detailed discussion of the various terms, by comparing the resulting  $C_\ell$  obtained from individual components in  $S(k, x)$  alone. Here, we begin by presenting an overall idea behind each term. **Dårlig avsnitt.**

The main physical idea behind the LOS approach, is that light observed in a given direction from the CMB today, is directly related to the CMB monopole along the LOS from us to infinity. The four terms in the source function represent different physical effects, with the first two being the so-called Sachs-Wolfe (SW) term and the integrated Sachs-Wolfe (ISW) effect. Since  $\tilde{g}$  is a sharply peaked at recombination, approximating it as a delta function yields

$$\begin{aligned} \Theta_\ell^{\text{today}}(k) & \approx [\Theta_0 + \Psi + \Theta_2/4]_{\text{rec}} \cdot j_\ell(k(\eta_0 - \eta_{\text{rec}})) \\ & \approx [\Theta_0 + \Psi]_{\text{rec}} \cdot j_\ell(k\eta_0), \end{aligned} \quad (74)$$

where we have used  $\eta_{\text{rec}} \ll \eta_0$ , and the fact that  $\Theta_2 \ll \Theta_0$  at recombination. The term  $\Theta_0 + \Psi$  represents an effective photon temperature. As the decoupled photons travel freely towards us today, they are gravitationally redshifted by the potential,  $\Psi$ . Hence, the SW term is an effective photon temperature with a tiny quadrupole correction. The anisotropies of the CMB power spectrum we observe today are therefore caused by temperature inhomogeneities present during recombination, as photons froze out during decoupling, and act as tracers of the physics present before they decoupled. **Last two sentences are awful.** From the approximation in Eq. (74), we see that  $\Theta_\ell \propto j_\ell(k\eta_0)$ . Since  $j_\ell(x)$  has a peak at  $x \sim \ell$ , we expect  $\Theta_\ell(k)$  to have a peak at  $k \sim \ell/\eta_0$ . Thus, following Eq. (69), we may expect perturbation features on scales  $k$  to be mapped onto angular scales of  $\ell \sim k\eta_0$  in  $C_\ell$ . **Fix explanation etc.**

The ISW represents the fact that gravitational potentials may change in time as photons move through it. For freely moving photons entering gravitational potentials, there is no net change in temperature if the potentials remain constant in time. Photons entering a gravitational potential gets an energy change that is cancelled out by the energy change it experiences when it leaves the gravitational potential again, if the potential remains constant in time. **Mention how this affects power spectrum at small and large scales.** The potentials are also weighted by a factor  $e^{-\tau}$ , so we only get a contribution from this term after recombination. Since the potentials are approximately constant during matter domination, we expect a contribution from ISW around recombination from intermediate scale modes, where the potential varies during recombination, as seen from Fig. 15. After this, we expect to get a contribution at late times, when the Universe starts accelerating. From Fig. 15, we see that the large scale modes varies most, and with  $\tau(0) = 0$ , we therefore expect the largest contribution to  $C_\ell$  to be at large scales, i.e. low  $\ell$ .

The third term in Eq. (73) is effectively a dipole term, since  $v_b \sim v_\gamma$  at early times. From our discussion of the evolution of the monopole and dipole, in Fig. 13 and 14, we can expect the dipole term to contribute on scales that had entered the horizon before decoupling, as the larger modes don't oscillate until they enter the horizon, which occurs when  $\tilde{g} \approx 0$ . The final term is a quadrupole term, and is related to the angular dependence of Thompson scattering. However, as we will see in Sect. 5.3, this term gives a very small contribution.

If the ISW contribution is ignored, for low  $l$ , we have  $(\Theta_0 + \Psi)_{\text{rec}} \approx (\Theta_0 + \Psi)_{\text{ini}}$ , since the large scale modes have not entered the horizon before decoupling. The SW term is therefore approximately given as  $\Theta_\ell \approx (\Theta_0 + \Psi)_{\text{ini}} j_\ell(k\eta_0)$ . Substituting  $x = k\eta_0$ , the integral in Eq. (69) for  $n_s = 1$  becomes

$$\begin{aligned} C_\ell & = 4\pi A_s |(\Theta_0 + \Psi)_{\text{ini}}|^2 \frac{1}{(k_{\text{pivot}} \eta_0)^{n_s-1}} \int_0^\infty dx j_\ell^2(x) x^{n_s-2}, \\ \Rightarrow \frac{\ell(\ell+1)}{2\pi} C_\ell & = A_s |(\Theta_0 + \Psi)_{\text{ini}}|^2 = \text{constant}. \end{aligned} \quad (75)$$

When plotting the CMB power spectrum, we will plot the RHS of Eq. (75) against  $\ell$  in units of  $\mu\text{K}^2$ . We have  $n_s = 0.965 < 1$ , so we have  $\ell(\ell+1)C_\ell \propto \ell^{n_s-1}$ , which will induces a small tilt at low  $\ell$ . The ISW will also result in a further deviation from a constant value.

### 5.2. Implementation details

One of the most crucial parts about integrating to obtain  $\Theta_\ell(k)$  and  $C_\ell$ , is that the quantities we're integrating are oscillating

rapidly. We must therefore ensure that we have a sufficiently high resolution, so that we accurately sample the integrands.

### 5.2.1. LOS integration

### 5.2.2. Computing the power spectrum (working title)

## 5.3. Results

### 5.3.1. CMB power spectrum


The angular power spectrum is shown in Fig. 18. We also include data from Planck Collaboration et al. (2020) for low  $\ell$  values. At higher  $\ell$ , neutrinos, helium and reionization becomes important, and the predicted value of  $C_\ell$  would deviate from that of observations (Write about how/why these things affect it.) To understand the physics behind the CMB power spectrum, we have also drawn vertical lines showing the peaks (red) and the troughs (blue) in the spectrum.

The peaks we observe in the power spectrum are a result of the oscillations of the photon baryon plasma. When photons decouple from baryons, modes at different scales will be in different states. If the plasma is fully compressed upon recombination, the photons have to escape from deep potential wells, and will therefore be redshifted. Similarly, if a mode is fully decompressed when recombination happens, the photons can escape without much energy loss. Since  $C_\ell$  depends on the absolute value of the photons multipoles, both of these cases will result in a peak of the power spectrum. The troughs therefore correspond to modes for which the temperature perturbation is approximately zero.

As seen in Sect. 4.3, the size of the modes strongly affect their behaviour before decoupling. Small scale modes will undergo several oscillations before recombination occurs. The number of oscillations a mode undergoes before decoupling decreases for larger scale modes, and for modes larger than the horizon during recombination, we expect no oscillations to take place. Thus, the peak at  $\ell = 205$  corresponds to a mode that just began to oscillate, with recombination taking place when it was at its maximum. For  $\ell < 205$ , the modes are gradually less compressed, until we reach scales of modes that had not entered the horizon before decoupling.


For modes where the oscillation is at a maximum value we obtain the peaks seen in the power spectrum.

As we saw when we studied the perturbations, the size of the modes affects its oscillatory behaviour.




**Fig. 18.** The predicted CMB power spectrum (black) compared with data from Planck at large scales. Fix y-label. Should indicate  $T_{\text{CMB0}}$

We plot  $\Theta_0$  as a function of  $x$  for modes of size  $k = \ell_{\text{peak}}/\eta_0$  and  $k = \ell_{\text{trough}}/\eta_0$ , shown in Fig. 19, where we also highlight the approximate value of  $x_{\text{rec}}$ . We see that the three modes correspond well with the peaks of the power spectrum. Similarly, we see that the troughs coincide with modes that are phase shifted with respect to the peaks, and roughly at their minima.



**Fig. 19.** Computed monopoles for of scale  $k = \ell/\eta_0$  for different values of  $\ell$ . The upper panel are  $\ell$  values where the power spectrum peaks, and the lower panel are the  $\ell$  values of the power spectrum troughs.



**Fig. 20.** The total angular power spectrum (dashed blue curve), compared to power spectra computed from individual terms in Eq. (73). For each term in the figure, the remaining terms in Eq. (73) have been set to zero.

### 5.3.2. Matter power spectrum


### 5.3.3. Photon multipoles

## 6. Conclusions


Write a short summary and conclusion in the end.

*Acknowledgements.* I thank my mom for financial support!






**Fig. 21.** Computed matter power spectrum (solid blue curve), compared with observational data from [cite](#). The equality scale,  $k_{\text{eq}}$ , is marked as the vertical black line.



**Fig. 22.** The term involved in the integrand of Eq. (69), for different values of  $\ell$ . Note that we scale each integrand by  $\ell(\ell + 1)$ , just as we did for  $C_\ell$ . This is done to see each of the terms simultaneously.



**Fig. 23.** The transfer function from Eq. (72) plotted for different values of  $\ell$ .

## References

- Betoule, M., Kessler, R., Guy, J., et al. 2014, A&A, 568, A22, doi: 10.1051/0004-6361/201423413
- Callin, P. 2006, arXiv e-prints, astro, doi: 10.48550/arXiv.astro-ph/0606683
- Dodelson, S. 2020, Modern Cosmology (Academic Press), doi: 10.1016/C2017-0-01943-2
- Galassi, M. e. a. 2009, GNU scientific library reference manual (Network Theory Ltd.)
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, A&A, 641, A6, doi: 10.1051/0004-6361/201833910
- Seljak, U., & Zaldarriaga, M. 1996, ApJ, 469, 437, doi: 10.1086/177793
- Winther, H. A. 2023, Cosmology II: Initial conditions. [https://cmb.wintherscoming.no/theory\\_initial.php#init](https://cmb.wintherscoming.no/theory_initial.php#init)

## Appendix A: Fiducial parameters

The parameter values we use in this report are

$$\begin{aligned}
 h &= 0.67, \\
 T_{\text{CMB}0} &= 2.7255 \text{ K}, \\
 \mathcal{N}_{\text{eff}} &= 3.046, \\
 \Omega_{b0} &= 0.05, \\
 \Omega_{\text{CDM}0} &= 0.267, \\
 \Omega_{k0} &= 0, \\
 \Omega_{\nu0} &= \mathcal{N}_{\text{eff}} \cdot \frac{7}{8} \left( \frac{4}{11} \right)^{4/3} \Omega_{\gamma0}, \\
 \Omega_{\Lambda0} &= 1 - (\Omega_{k0} + \Omega_{b0} + \Omega_{\text{CDM}0} + \Omega_{\gamma0} + \Omega_{\nu0}), \\
 n_s &= 0.965, \\
 A_s &= 2.1 \cdot 10^{-9}, \\
 Y_p &= 0.245, \\
 z_{\text{reion}} &= 8, \\
 \Delta z_{\text{reion}} &= 0.5, \\
 z_{\text{Hereion}} &= 3.5, \\
 \Delta z_{\text{Hereion}} &= 0.5,
 \end{aligned} \tag{A.1}$$