OCCUPATIONAL THERAPY [M.O.TH]
Qns.1) An Injury to which of the following structure results in Volkmann`s Ischemic Contracture : **Option 1 :** Median Nerve **Option 2 :** Brachial Artery **Option 3 :** Ulnar Nerve **Option 4 :** Radial Artery
Qns.2) A positive Trendelenburg`s sign is observed in paralysis of which of the following muscle ? **Option 1 :** Gluteus Maximus **Option 2 :** Gluteus Medius **Option 3 :** Piriformis **Option 4 :** Quadratus Femoris
Qns.3) Gag reflex is absent in lesion of which of the following cranial nerves ? **Option 1 :** Trigeminal **Option 2 :** Facial **Option 3 :** Glossopharygeal **Option 4 :** Vagus
Qns.4) Fibers passing through the Genu of the internal capsule are : **Option 1 :** Corticonuclear **Option 2 :** Corticospinal **Option 3 :** Auditory radiation **Option 4 :** Optic radiation
Qns.5) The receptor for pain is : **Option 1 :** Ruffinis end organ **Option 2 :** Merckels Disc **Option 3 :** Golgi body **Option 4 :** Free nerve endings
Qns.6) Cardiac muscle is able to function as a syncytium because of the structural presence of _____ **Option 1 :** Branching fibers **Option 2 :** Intercalated discs **Option 3 :** Protoplasmic bridges between cells **Option 4 :** Gap junctions
Qns.7) Somato statin inhibits the release of : **Option 1 :** GH **Option 2 :** ACTH

Option 3 : TSH
Option 4 : PTH

Qns.8) At what stage of erythropoiesis does hemoglobin appear ?
Option 1 : Pro-erythroblast
Option 2 : Early normoblast
Option 3 : Intermediate normoblast
Option 4 : Reticulocyte

Qns.9) All of the following enzymes are useful in the diagnosis of liver diseases, EXCEPT :
Option 1 : Alkaline Phosphatase
Option 2 : Aspartate Transaminase
Option 3 : Creatine Phosphokinase
Option 4 : Alanine Transaminase

Qns.10) The Amino Acid that does not participate in transamination is :
Option 1 : Lysine
Option 2 : Glutamate
Option 3 : Alanine
Option 4 : Tryptophan

Qns.11) 58 year old female with right shoulder periarthritis, has limitations in the shoulder range and muscle power, she is able to move her arm through more than 50% of full range of motion against gravity, she has a muscle power of :
Option 1 : 2
Option 2 : 3+
Option 3 : 3
Option 4 : 3−

Qns.12) Patient with fracture patella is being treated in therapy following tension band wiring. His quadriceps strength is 3 minus, the occupational therapist should prefer to give him following exercises in anti gravity position :
Option 1 : Isometric exercise
Option 2 : Isotonic exercise
Option 3 : Isotonic assistive exercise
Option 4 : Resistive exercise

Qns.13) 30 year old female with sensory involvement had been referred for occupational therapy treatment to give sensory re-education. The therapist uses therapeutic activity to fulfill the goals using following aspect of activity analysis :
Option 1 : Biomechanical
Option 2 : Sensory motor

Option 3 : Cognitive
Option 4 : Psychosocial

Qns.14) The company provides an occupational therapist, a description that includes essential tasks of job, physical requirements, cognitive aptitude, educational requirement, equipment operations and environmental assist in selecting suitable candidates for employment. The Occupational Therapist uses this knowledge to analyze work fitness for patient. This is called :
Option 1 : Work Capacity Evaluation
Option 2 : Functional Capacity Evaluation
Option 3 : Job Demand Analysis
Option 4 : Work Hardening

Qns.15) When the splints are designed to increase limited range of motion or to restore or augment function, they are classified as :
Option 1 : Mobilization splint
Option 2 : Immobilization splint
Option 3 : Restriction splint
Option 4 : Single surface splint

Qns.16) While obtaining information for Occupational therapy intervention , patients and relatives are asked to respond to Questionnaires` regarding interests, hobbies and desires during assessment procedures, it is generally collected through :
Option 1 : Inventories and checklists
Option 2 : Object history
Option 3 : Medical records
Option 4 : Observation

Qns.17) Tophi near the joints are seen in :
Option 1 : Tuberculosis
Option 2 : Gout
Option 3 : Osteoporosis
Option 4 : Osteo-sarcoma

Qns.18) Typhoid Ulcers in intestine are :
Option 1 : Small superficial ulcers
Option 2 : Longitudinal oval ulcers parallel to long axis of intestine
Option 3 : Transverse ulcers perpendicular to long axis of intestine
Option 4 : Flask shaped ulcers

Qns.19) The following bacterial organelle is used for adhesion to mucosal surface :
Option 1 : Fimbriae
Option 2 : Capsule

Option 3 : Flagella
Option 4 : Cell wall

Qns.20) Natural passive immunity is due to :
Option 1 : Natural disease
Option 2 : Vaccine
Option 3 : Placental transfer from mother to fetus
Option 4 : Hyper immune Sera

Qns.21) This anti-rheumatoid drug is classified as an Interleutcin-1 (IL-1) antagonist :
Option 1 : Infliximab
Option 2 : Leflunomide
Option 3 : Etanercept
Option 4 : Anakinra

Qns.22) Fulminant hepatic failure has been associated with this Non-Steroidal Anti Inflammatory Drug (NSAID) :
Option 1 : Diclofenac Sodium
Option 2 : Meloxicam
Option 3 : Nimesulide
Option 4 : Ketoprofen

Qns.23) Movement perceived in the absence of physical movement of an image across the retina :
Option 1 : Real motion
Option 2 : Stroboscopic motion
Option 3 : Apparent motion
Option 4 : Movement perception

Qns.24) A set of rules if followed correctly guarantees a solution to the problem :
Option 1 : Heuristics
Option 2 : Algorithm
Option 3 : Set
Option 4 : Habit

Qns.25) The highest level of ego defenses which consists of a redirection of sexual impulses to socially valued activities & goals :
Option 1 : Intellectualization
Option 2 : Regression
Option 3 : Rationalization
Option 4 : Sublimation

Qns.26) A theory of motivation which states that we are motivated to seek goals which give us good emotional feeling & to avoid those resulting in displeasure :

Option 1 : Opponent-process theory
Option 2 : Incentive theory
Option 3 : Drive theory
Option 4 : Optimal-level theory

Qns.27) Pronation of the non weight bearing Calcaneous on the fixed Talus and the lower leg, is composed of the following coupled motions, EXCEPT :
Option 1 : Abduction
Option 2 : Eversion
Option 3 : Plantar flexion
Option 4 : Dorsiflexion

Qns.28) During knee extension ,motion of the femoral condoyle on a fixed tibia, is as follows :
Option 1 : Posterior rolling with simultaneous posterior gliding
Option 2 : Anterior rolling with simultaneous anterior gliding
Option 3 : Posterior rolling with simultaneous anterior gliding
Option 4 : Anterior rolling with simultaneous posterior gliding

Qns.29) In all second class levers, the mechanical advantage of the lever will always be :
Option 1 : Greater than one
Option 2 : Lesser than one
Option 3 : Equal to one
Option 4 : None of those given as options

Qns.30) The overall ratio of Gleno-humeral to Scapulo-thoracic motion during arm elevation is :
Option 1 : 1 : 1
Option 2 : 2 : 1
Option 3 : 3 : 1
Option 4 : 4 : 1

Qns.31) The Lumbar pelvic rhythm during bending over to touch one`s toes with knees straight is composed of :
Option 1 : Anterior tilting of pelvis followed by Lumbosacral flexion
Option 2 : Lumbosacral flexion followed by anterior tilting of pelvis
Option 3 : Posterior tilting of pelvis followed by Lumbosacral flexion
Option 4 : Lumbosacral flexion followed by posterior tilting of pelvis

Qns.32) Transfer of Learning refers to a person`s ability to carry out :
Option 1 : Same task in a different environment
Option 2 : Same task in same environment
Option 3 : Varying task in a varying environment
Option 4 : Varying task in same environment

Qns.33) According to Brunnstorm stages of motor recovery after CVA, if ``Gross grasp and Hook grasp are possible, with No release`` then the hand is in following stage of recovery :

Option 1 : 2
Option 2 : 3
Option 3 : 4
Option 4 : 5

Qns.34) The Minnesota Rate of Manipulation Test consists of :

Option 1 : Four horizontal rows of openings with 80 disks
Option 2 : Four horizontal rows of openings with 60 disks
Option 3 : Six horizontal rows of openings with 80 disks
Option 4 : Six horizontal rows of openings with 60 disks

Qns.35) When a child plays with toys, similar to those used by other children near him but, plays beside rather than with the children, it is known as the following type of play :

Option 1 : Onlooker
Option 2 : Solitary independent
Option 3 : Parallel
Option 4 : Associative

Qns.36) According to Piagets` stages of cognitive development, `concrete operational stage` is seen during the following years of a child :

Option 1 : Birth to 2 years
Option 2 : 2 to 6 years
Option 3 : 6 to 12 years
Option 4 : At 16 years

Qns.37) Which of the following groups correctly describes clinical features of Parkinson`s disease ?

Option 1 : Asymmetric onset ,Lead pipe rigidity, intentional tremors
Option 2 : Symmetric onset, cog wheel rigidity, resting tremors
Option 3 : Asymmetric onset, cog wheel rigidity, resting tremors
Option 4 : Asymmetric onset, clasp knife rigidity, intentional tremors

Qns.38) Which nutritional deficiency is associated with parasthesia, sensory ataxia, weakness, spasticity, extensor planter responses and diminished ankle jerks ?

Option 1 : Vitamin B_{12}
Option 2 : Vitamin D
Option 3 : Vitamin B_1
Option 4 : Vitamin B_2

Qns.39) ``Brocks syndrome`` refers to recurrent or fixed atelectasis of _____

Option 1 : Right upper lobe

Option 2 : Right middle lobe
Option 3 : Left upper lobe
Option 4 : Left lower lobe

Qns.40) Which one of the following joints is commonly involved in osteoarthritis ?
Option 1 : Ankle
Option 2 : Elbow
Option 3 : Intertarsal joints
Option 4 : First carpometacarpal joint

Qns.41) Which disease has enthesitis as an important manifestation ?
Option 1 : Rheumatoid arthritis
Option 2 : SLE [systemic lupus erythematosis]
Option 3 : Ankylosing spondylitis
Option 4 : Gouty arthritis

Qns.42) All of the following can be signs and symptoms of congestive heart failure, EXCEPT :
Option 1 : Nocturnal cough
Option 2 : Weight gain and peripheral edema
Option 3 : Distension of jugular veins
Option 4 : Clubbing

Qns.43) Immediate management of a patient with multiple fractures and blood loss includes infusion of _____
Option 1 : Blood
Option 2 : Dextran
Option 3 : Normal saline
Option 4 : Ringer`s lactate

Qns.44) Best vein for infusion of total Parenteral Nutrition is :
Option 1 : Brachial vein
Option 2 : Femoral vein
Option 3 : Subclavian vein
Option 4 : Saphenous vein

Qns.45) Thiersch graft is which type of graft ?
Option 1 : Partial thickness
Option 2 : Pedicle
Option 3 : Full thickness
Option 4 : Patch

Qns.46) All of the following are possible Complications of Distal End Radius Fractures, EXCEPT :
Option 1 : Carpal Tunnel Syndrome
Option 2 : Sudeck`s Osteo-dystrophy

Option 3 : Non union
Option 4 : Mal-union

Qns.47) ``Pointing Index Sign`` is seen after injury to which nerve :
Option 1 : Ulnar nerve
Option 2 : Median nerve
Option 3 : Radial nerve
Option 4 : Axillary nerve

Qns.48) Kienbock`s Disease is due to Ischemic Necrosis of :
Option 1 : Lunate
Option 2 : Capitate
Option 3 : Calcaneum
Option 4 : Capitullum

Qns.49) Which Is **not** a Cannabis Preparation :
Option 1 : Charas
Option 2 : Cocaine
Option 3 : Ganja
Option 4 : Bhang

Qns.50) Fear of heights / high places is termed as :
Option 1 : Acrophobia
Option 2 : Xenophobia
Option 3 : Algophobia
Option 4 : Claustrophobia

Qns.51) Children with severe mental retardation have an IQ in the range of :
Option 1 : 50 to 70
Option 2 : 35 to 50
Option 3 : 20 to 35
Option 4 : Less than 20

Qns.52) Cluster B personality disorder includes all, EXCEPT :
Option 1 : Histrionic personality disorder
Option 2 : Antisocial personality disorder
Option 3 : Borderline personality disorder
Option 4 : Dependant personality disorder

Qns.53) In industrial settings, displays with controls, dials and counters are used for :
Option 1 : Qualitative readings
Option 2 : Tracking
Option 3 : Comparison
Option 4 : Quantitative readings

Qns.54) For optimal functioning body temperature should be maintained

between :
Option 1 : 35° to 38°
Option 2 : 36.5° to 40°
Option 3 : 38° to 42°
Option 4 : 40° to 45°

Qns.55) A 45 year old chronic Rheumatoid Arthritis patient has limitations in vocational and avocational activities. She is independent with her self care activities. Her functional classification as per revised criteria of American college of Rheumatology is :
Option 1 : CLASS I
Option 2 : CLASS II
Option 3 : CLASS III
Option 4 : CLASS IV

Qns.56) MET is a multiple of metabolic equivalents of energy at rest in Cardiopulmonary Conditions to indicate O_2 consumption. It is measured as :
Option 1 : 3.5 ml of O_2 /Kg/min of body weight
Option 2 : 3.5 ml of O_2 /min/Kg of body weight
Option 3 : 3.5 L of O_2 /Kg/min of body weight
Option 4 : 3.5 L of O_2 /min/Kg of body weight

Qns.57) Following activities require an approximate metabolic cost of energy at 1.5 to 2 METs, EXCEPT :
Option 1 : Auto repair
Option 2 : Auto driving
Option 3 : Typing
Option 4 : Deskwork

Qns.58) 40 year old male laborer, diagnosed as case of HIV, is referred to Occupational Therapy for tertiary Prevention Program. Occupational Therapist should concentrate on the following :
Option 1 : Prevent opportunistic infections
Option 2 : Education on abstinence & safe sex
Option 3 : Interpersonal skills training
Option 4 : Continued Support for participation in Occupations

Qns.59) One of the following Occupational therapy treatment approaches is **NOT** used in geriatric practice :
Option 1 : Prevention approach
Option 2 : Accommodation approach
Option 3 : Restriction approach
Option 4 : Restoration approach

Qns.60) For a patient with Ulnar drift and volar subluxation of the wrist,

the most appropriate joint protection technique to be advised is :
Option 1 : Keep pulling the wrist in radial deviation
Option 2 : Avoid turning taps and jar covers towards the ulnar side of the hand
Option 3 : Use splint to avoid these movements
Option 4 : Use adaptive devices

Qns.61) The anti-deformity position for neck burns is :
Option 1 : Neutral
Option 2 : Slight flexion
Option 3 : 5-10° of extension
Option 4 : Slight rotation

Qns.62) Following exercises are indicated in Klinert`s protocol for management of flexor tendon injuries during first three weeks in the post operative phase :
Option 1 : Passive flexion, active extension
Option 2 : Active flexion, passive extension
Option 3 : Passive flexion, passive extension
Option 4 : Active flexion, active extension

Qns.63) A 65 year old female with mastectomy and axillary node dissection is referred for therapy intervention on the second day of surgery. Following activity should be contraindicated to her :
Option 1 : Pouring liquids into cup
Option 2 : Light dusting on tabletop
Option 3 : Reach shelves at overhead level
Option 4 : Pick up a phone for communication

Qns.64) A 30 year old data entry operator has a residual visual impairment and is ready to resume work. The occupational therapist would recommend the following to him, EXCEPT :
Option 1 : Software to navigate with and between the documents, using key controls
Option 2 : Captioning on multimedia software
Option 3 : Optical character recognition (OCR)
Option 4 : Voice convertor/speech synthesizer or refreshable Braille display

Qns.65) Total active motion (TAM) is a measure of the small joints of the hand, and is measured as :
Option 1 : Total range of motion of MP, PIP and DIP flexion, minus available extension of DIP
Option 2 : Range of motion of MP,PIP, DIP flexion added to active extension of MP,PIP and DIP joints
Option 3 : Total of MP, PIP, and DIP flexion and active extension ranges

of DIP joint
Option 4 : Total of MP, PIP and DIP flexion, minus incomplete active extension of each of the three joints

Qns.66) A person with 30 % loss of humeral length should be recommended the following :
Option 1 : Above elbow prosthesis
Option 2 : Below elbow prosthesis
Option 3 : Elbow disarticulation prosthesis
Option 4 : Shoulder disarticulation prosthesis

Qns.67) Post traumatic amnesia of 1-24 hours corresponds to following severity of traumatic brain injury :
Option 1 : Mild
Option 2 : Moderate
Option 3 : Severe
Option 4 : Very severe

Qns.68) Following traditional sensory-motor approach uses a treatment that directly emphasizes the development of skilled movements for task performance :
Option 1 : Neuro-Developmental Approach
Option 2 : Proprioceptive Neuromuscular Facilitation Approach
Option 3 : Brainstorm`s Approach
Option 4 : Rood`s Approach

Qns.69) Following are the primitive reflexes associated with brainstem, EXCEPT :
Option 1 : Crossed Extension Reflex
Option 2 : Asymmetric Tonic Neck Reflex
Option 3 : Symmetric Tonic Neck Reflex
Option 4 : Tonic Labyrinthine Reflex

Qns.70) Following are the `Manual Rating Scales` for spasticity and hyper-tonicity, EXCEPT :
Option 1 : Ashworth scale
Option 2 : Tardian scale
Option 3 : Myotonometer
Option 4 : Preston`s hypertonicity scale

Qns.71) According to Modified Brunnstorm stages of motor recovery of arm and hand, a patient is an excellent candidate for fine- tuning motor control via blocks, when he is in following stage of recovery :
Option 1 : Stage 2
Option 2 : Stage 3
Option 3 : Stage 4
Option 4 : Stage 5

Qns.72) Central cord syndrome is characterized by prominent cellular destruction in the centre of the cord as compared to the periphery of the cord. The presentation in the central cord syndrome is thus :

Option 1 : Greater sensory and motor loss in lower extremities versus upper extremities

Option 2 : Greater sensory and motor loss in upper extremities versus lower extremities

Option 3 : Greater sensory and motor loss in spinal muscles versus extremities

Option 4 : Greater sensory and motor loss in both extremities versus spinal muscles

Qns.73) For a client with Parkinson`s who is on levodopa, the best period for therapeutic intervention to promote self-care activities is :

Option 1 : ``On`` period

Option 2 : ``Off`` period

Option 3 : ``In`` period

Option 4 : ``Out`` period

Qns.74) Activities that require frustration tolerance and attention to detail, should be avoided in patients exhibiting :

Option 1 : Paranoia

Option 2 : Hostility and aggression

Option 3 : Seductive behavior

Option 4 : Anxiety

Qns.75) According to Beck`s level of cognition, `The foundations of our cognitive constructs` is provided by :

Option 1 : Automatic thoughts

Option 2 : Beliefs

Option 3 : Core schema

Option 4 : None of the options provided

Qns.76) A patient has been depressed for a long time, the depression is just lifting, she complains to the treating therapist that she is fed up of life, the most important precaution to take in case of this patient is :

Option 1 : Counsel the patient

Option 2 : Report to the treating team

Option 3 : Do not leave the patient unsupervised anytime

Option 4 : Increase the dose of antidepressants

Qns.77) Following is a standardized, performance based task that involves imitation of leather lacing :

Option 1 : Comprehensive Occupational therapy evaluation (COTE)

Option 2 : Azima battery

Option 3 : Allen`s cognitive level test (ACL)
Option 4 : Comprehensive evaluation of basic skills

Qns.78) A method of approaching the terminal behavior gradually , using a series of steps that lead to the goal is called :
Option 1 : Forward chaining
Option 2 : Backward chaining
Option 3 : Schedule of reinforcement
Option 4 : Shaping

Qns.79) The following frame of reference is applicable to the patient population of all ages and diagnoses :
Option 1 : Acquisitional
Option 2 : Model of Human Occupation
Option 3 : Psychodynamic
Option 4 : Sensory Integration

Qns.80) ``Wernicke`s Encephalopathy`` is a common Neuro-psychiatric complication in which of the following chronic substance abuse --- ?
Option 1 : Opioid
Option 2 : Alcohol
Option 3 : Cannabis
Option 4 : Cocaine

Qns.81) Autonomic dysreflexia is usually seen in patients with lesion at T6 and above. The symptoms include all of the following EXCEPT :
Option 1 : Hypertension
Option 2 : Sweating
Option 3 : Pupil constriction
Option 4 : Hypothermia

Qns.82) An individual who shows weakness in pinch, clumsiness in hand functions, hyperesthesia`s or parasthesia in median nerve distribution of the hand and nocturnal pain may be diagnosed as :
Option 1 : Guyon`s Canal Entrapment
Option 2 : Carpal Tunnel Syndrome
Option 3 : deQuervain`s Tenosynovitis
Option 4 : None of those given as options

Qns.83) In case of spinal cord injuries at level C5- C6, expected functional outcome for eating will be :
Option 1 : Dependent
Option 2 : Independent
Option 3 : Totally assisted
Option 4 : Modified independence

Qns.84) When partial weight bearing is advised after hip surgery, the

following percentage of body weight is borne by operated leg for functional tasks :

Option 1 : 15
Option 2 : 30
Option 3 : 50
Option 4 : 75

Qns.85) A 69 year old right handed retired sales manager with an 8 to 10 years history of Dupyutren`s Contracture has been referred to Occupational Therapy after reconstruction of contractures including Z-plasty. All of the following are the problem areas EXCEPT :

Option 1 : Limited active and passive range of motion
Option 2 : Dense scar tissue
Option 3 : Sensation
Option 4 : Edema

Qns.86) Tenodesis grasp is seen in patients with the following level of SCI :

Option 1 : C1-C2
Option 2 : C3-C4
Option 3 : C6-C7
Option 4 : C8

Qns.87) In quadriparesis, tenodesis hand grasp is present at all cervical levels ,EXCEPT at :

Option 1 : C5 level
Option 2 : C6 level
Option 3 : C7 level
Option 4 : C8 level

Qns.88) Stress management techniques like autogenic training, bio-feedback, time management and verbalization function involve intervention at the following nervous system level :

Option 1 : Hypothalamus
Option 2 : Cortex
Option 3 : Reticular activating system
Option 4 : Peripheral Nervous system

Qns.89) Patients with spinal cord injury are rarely associated with problems of :

Option 1 : Fertility
Option 2 : Erection
Option 3 : Ejaculation
Option 4 : Vaginal lubrication

Qns.90) The basic accounting method for organizing data in reference to cost is

Option 1 : Balance sheet
Option 2 : Chart of accounts
Option 3 : Direct cost
Option 4 : Indirect cost

Qns.91) ``To assure third party reimbursement`` (insurance coverage) for a patient under home care, while documenting activities, therapist has to follow constraints and can recommend following activities ,EXCEPT :
Option 1 : Bathing
Option 2 : Laying Table
Option 3 : Cleaning Closet
Option 4 : Gardening

Qns.92) Drugs prescribed for the following conditions are known to cause impotency :
Option 1 : Hypertension and depression
Option 2 : Hypertension and common cold
Option 3 : Allergic reaction and common cold
Option 4 : Allergic reaction and depression

Qns.93) In patient with lower extremity weakness and poor endurance in upper extremity, following type of transfer technique is used;
Option 1 : Stand pivot transfer
Option 2 : Bent pivot transfer
Option 3 : Sliding board transfer
Option 4 : Dependent transfer

Qns.94) Following type of budget is used for the purchase of major equipments or improvements of the facility :
Option 1 : Departmental budget
Option 2 : Capital budget
Option 3 : Revenue expenses & budget
Option 4 : Developmental budget

Qns.95) The following is the managerial style, in which the manager has low involvement with people and contributes minimally towards organizational purpose :
Option 1 : Country Club Style
Option 2 : Authoritarian Style
Option 3 : Participatory Style
Option 4 : Laissez -faire style

Qns.96) Which one of the following is useful index for surveillance of Iodine deficiency :
Option 1 : Prevalence of goiter
Option 2 : Prevalence of Cretinism

Option 3 : Serum T4 levels
Option 4 : Urinary Iodine secretion

Qns.97) ``Physical quality of Life Index`` consolidates following three indicators ,EXCEPT :
Option 1 : Infant Mortality Rate
Option 2 : Life Expectancy at age one
Option 3 : Maternal Mortality Rate
Option 4 : Literacy Rate

Qns.98) The sampling technique, in which a part of the information is collected from the whole sample, and the part is collected from the subsample is called as _____ sampling
Option 1 : Multiphase
Option 2 : Multistage
Option 3 : Cluster
Option 4 : Systematic

Qns.99) The standard deviation of a standard normal curve is taken as :
Option 1 : Zero (0)
Option 2 : One
Option 3 : Two
Option 4 : Three

Qns.100) The statistical test which is used as a test of ``goodness of fit`` is :
Option 1 : Mc Nemars test
Option 2 : Chi square test
Option 3 : Standard error of mean
Option 4 : Student T test