

yes i know we need a title

1 Context

The beginning of XXIth century sees a drastic increase of applications coming as an assistance to numerous human activities, among which medicine is one of the main domains. And the use of technology to the healthcare system is the focus today, with technologies ranging from robots to assist in surgery to gadgets that measure sugar levels and help diabetic people. Among those areas, an important one deals with pain, which is a complex subject to master. The main objectives will consist in being able to properly measure pain levels is of a great importance, as it will help medical doctors to provide a precise treatment and also allow pharmaceutical companies to have a better feedback for the development of treatments, the last being pharmaceutical or not. But pain assessment is not a simple task and has been a challenge even for humans. Many scales and methods have been created to tackle such a problem [3, 4, 5, 6, 7, 8] with limited efficiency. It is still so far difficult to have an accurate assessment of pain, notably chronic pain. As a matter of fact, as pain is a subjective feeling, the most used measurement is the auto evaluation, but even if it gives fairly acceptable results, it still falls short on some aspects like simulated pain [1], or, most commonly, when the patient has difficulties to communicate, as is the case of infants and some elderly people [2]. And the creation of a method capable of assessing precisely the level of pain felt by patients can help the present diagnosis system overcome a complicated barrier.

1.1 state of the art

1.2 methodological appro

2 objectives

2.1 expected results

2.2 scientific and technological challenges

3 organization

3.1 general organization

3.2 planning

T0 + 6 months: gathering and study of the data
T0 + 12 months: assessment based on the facial expression
T0 + 18 months: assessment of emotions from natural language analyses
T0 + 24 months: assessment by combining the 2 methods
T0 + 30 months: formalization, validations publications and theses
T0 + 36 months: theses and defense

References

- [1] CL Gwen, C Marian, Stewart Bartlett, and Kang Lee. Faces of pain: automated measurement of spontaneous all facial expressions of genuine and posed pain. 2007.
- [2] Patrick Lucey, Jeffrey F Cohn, Iain Matthews, Simon Lucey, Sridha Sridharan, Jessica Howlett, and Kenneth M Prkachin. Automatically detecting pain in video through facial action units. *IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics)*, 41(3):664–674, 2011.
- [3] Donna Lee Wong, M Hockenberry-Eaton, D Wilson, ML Winkelstein, and P Schwartz. Wong-baker faces pain rating scale. *Home Health Focus*, 2(8):62, 1996.
- [4] M McCaffery and C Pasero. Pain: clinical manual . st louis, mo: Mosby, 1999.
- [5] RK Portenoy and RM Tanner. Visual analog scale and verbal pain intensity scale: from pain management: theory and practice. *Oxford University Press, Inc*, 1:996, 1996.

- [6] Ronald Melzack. The mcgill pain questionnaire: major properties and scoring methods. *Pain*, 1(3):277–299, 1975.
- [7] Bradley S Galer and Mark P Jensen. Development and preliminary validation of a pain measure specific to neuropathic pain the neuropathic pain scale. *Neurology*, 48(2):332–338, 1997.
- [8] Richard H Gracely and Donna M Kwilosz. The descriptor differential scale: applying psychophysical principles to clinical pain assessment. *Pain*, 35(3):279–288, 1988.