
A. Rate by GBD region
B. Share
C. Change, 2010–2023
SUB-SAHARAN AFRICA
Western Sub-Saharan Africa
Central Sub-Saharan Africa
Eastern Sub-Saharan Africa
Southern Sub-Saharan Africa
SOUTH ASIA
South Asia
SE ASIA, E ASIA & OCEANIA
Oceania
Southeast Asia
East Asia
LATIN AMERICA & CARIBBEAN
Tropical Latin America
Andean Latin America
Caribbean
Central Latin America
N AFRICA & MIDDLE EAST
North Africa and Middle East
C EUROPE, E EUROPE & C ASIA
Central Asia
Eastern Europe
Central Europe
HIGH-INCOME
Southern Latin America
High-income Asia Pacific
High-income North America
Western Europe
Australasia
49·7
13·4
16·7
9·2
3·1
1·9
1·1
2·1
Only regional increase:
malaria, N Africa & Middle East
≤1
10
100
1 000
0
25
50
−100
−50
0
50
Age-standardised DALYs per 100 000, 2023
(log scale; | = malaria ≤1)
Share of global
NTD + malaria DALYs (%)
Change in age-standardised rate, 2010–2023
(%, 95% UI; × = eliminated)
NTDs excl. malaria (A); NTDs and malaria combined (C)
Malaria