Global, regional and national inequalities in neglected tropical disease and malaria burden in 204 countries and territories, 1990–2023, and the projected consequences of external financing cuts: an econometric modelling study

Vincent Uwumboriyhie Gmayinaam[1], Abraham Norman Nortey, Patrick Elorm Dzissem[1], Mustapha Immurana[2], Alfred Kwesi Manyeh[1]

[1]Centre for Neglected Tropical Diseases, Institute of Health Research, University of Health and Allied Sciences, Ho, Volta Region, Ghana *(V U Gmayinaam MPH, P E Dzissem MPH, M Immurana Bsc. PhD, A K Manyeh, Msc. PhD)*

Corresponding author: (Dr, Mustapha Immurana, PMB 31, Ho, Volta Region, Ghana., mimmurana@uhas.edu.gh)

ORCID iDs:

Vincent Uwumboriyhie Gmayinaam, https://orcid.org/0000-0001-6997-1903
Patrick Elorm Dzissem,

Emails:

V.U.G: vugmayinaam@uhas.edu.gh
P.E.D: pedzissem@uhas.edu.gh
M.I: mimmurana@uhas.edu.gh
A.K.M: akmanyeh@uhas.edu.gh

Summary

BackgroundNeglected tropical diseases (NTDs) and malaria are concentrated in countries that depend heavily on external health financing, which fell by about a fifth in 2025. We quantified how this burden is distributed, how socioeconomic inequality in it changed, and the health losses that financing cuts could cause.

MethodsIn this ecological study, we analysed Global Burden of Disease (GBD) 2023 disability-adjusted life-years (DALYs) for 204 countries and territories, 1990–2023, by super-region, region, and country. We measured spatial clustering, absolute and relative inequality by Socio-demographic Index (SDI) and Theil decomposition. Panel models for 153 low- and middle-income countries, 2007–2022, related external financing share to DALY rates; two scenario models projected excess DALYs under 5–60% cuts.

FindingsIn 2023, NTDs and malaria caused 70·8 million DALYs (51·6 million from malaria). Sub-Saharan Africa had 15·1% of the world population but 80·2% of these DALYs; western and central sub-Saharan Africa alone had 63·1%. Age-standardised rates fell 21·6% (95% UI 7·8–36·3) from 2010. The absolute SDI gap narrowed (NTD slope index of inequality −1966 to −632 DALYs per 100 000, 1990–2023), but relative concentration in low-SDI countries persisted (malaria concentration index −0·79 in 1990 and −0·81 in 2023). Disability accounted for 62·4% of NTD DALYs. A 10-percentage-point higher external share was associated with a 13·4% (95% CI 6·4–20·0) lower NTD rate between countries but not within countries (−0·64%, −2·39 to

1·14). A 19–26% cut projected 0·41 million (95% UI 0·33–0·50) excess NTD and 1·85 million (1·51–2·21) excess malaria DALYs a year, over 90% in sub-Saharan Africa.

InterpretationFalling averages have not moved burden away from the poorest, most donor-dependent countries, where projected losses also concentrate. Protecting service continuity there is a priority; the projections are conditional scenarios, not causal estimates.

Funding

This study was self-funded.

Research in context

Evidence before this study

We searched PubMed from inception to Oct 21, 2026, without language restrictions, for studies of NTD or malaria burden, inequality, spatial analysis, development assistance for health, and funding cuts, and reviewed WHO and IHME reports; 76 of 704 unique records were eligible (appendix). GBD 2019 and 2021 analyses showed combined NTD and malaria DALYs falling by about 18% between 1990 and 2021, with burden concentrating in low-SDI countries and relative inequality widening as absolute inequality narrowed. None used GBD 2023, tested geographical clustering, or decomposed inequality between and within regions. Cross-country panels associated aid volume with lower malaria, NTD, and child mortality and projected 14–23 million additional deaths by 2030 from current cuts, but none separated between-country from within-country associations, examined dependence on external financing, or estimated NTD DALYs under the cuts.

Added value of this study

Using GBD 2023, we show that 63·1% of global NTD and malaria DALYs occur in western and central sub-Saharan Africa, that burden hot spots were unchanged between 2010 and 2023, and that the absolute socioeconomic gap narrowed while relative concentration in the poorest countries did not. Most NTD burden is disability, which mortality-based evaluations omit. Thirty-four countries combining high NTD burden and high donor dependence hold over half of malaria DALYs. The association of external financing with NTD burden seen between countries was absent within countries over time. We provide country-level estimates of NTD and malaria DALYs at risk under specified cuts.

Implications of all the available evidence

The 2025 reductions in development assistance fall where NTD and malaria burden and financing dependence are greatest, which supports protecting service continuity in those countries. Aggregate financing shares should not be read as disease-specific effects; estimating the consequences of the cuts needs programme disbursement data linked to coverage and outcomes.

Introduction

NTDs and malaria remain central to Sustainable Development Goal 3.3. In 2023, 1·495 billion people required NTD interventions, and in 2024 malaria caused an estimated 282 million cases and 610 000 deaths.[1,2] Combined DALYs fell by about 18% between 1990 and 2021, to 71·6 million,[3] and successive GBD rounds show burden concentrating in low-SDI countries and western sub-Saharan Africa.[4–6] Between-country inequality narrowed in absolute terms but widened in relative terms,[7–11] and NTD

treatment coverage remains uneven within countries.[12,13] None of these studies used GBD 2023, tested whether burden clusters geographically, or decomposed inequality between and within regions.

External financing underpinned much of this progress. Cross-national panels associate development assistance for health with lower malaria and child mortality,[14–18] and evaluations attribute large gains to the US President's Malaria Initiative and insecticide-treated nets.[19,20] Development assistance for health fell by an estimated 22% between 2024 and 2025.[21] Recent studies project 14–23 million additional deaths by 2030 from USAID and broader aid cuts,[22,23] and models address HIV, tuberculosis, and malaria.[24–26] These analyses model aid volume and mortality. Yet most NTD burden is non-fatal, and because aid flows towards high-burden countries,[27,28] between-country associations can differ from within-country change. WHO already reports disruption of NTD programmes.[29] No study has estimated NTD DALYs at risk from the cuts.

We analysed NTD and malaria burden in 204 countries and territories using GBD 2023, at super-region, regional, and national level. We quantified spatial clustering, absolute and relative socioeconomic inequality from 1990 to 2023, and the fatal and non-fatal composition of burden; compared between-country and within-country associations of external financing dependence with burden; and projected excess DALYs under specified financing cuts.

Methods

Study design and data

We conducted an ecological study of 204 countries and territories in seven GBD super-regions and 21 regions, reported according to GATHER and STROBE (appendix section 1).[30,31] We linked GBD 2023 estimates of DALYs, years lived with disability (YLDs) and years of life lost (YLLs), World Bank World Development Indicators (WDI), WHO indicators from the UN SDG database, IHME Financing Global Health estimates, and Natural Earth boundaries (Table S1; Figure 1).[21,32,33] Outcomes were age-standardised DALY rates per 100 000 for NTDs excluding malaria (the GBD NTD-and-malaria aggregate minus malaria; 19 causes) and for malaria; counts were all-age. Percentage changes with uncertainty intervals (UIs) were taken from GBD. GBD does not publish changes for NTDs excluding malaria or country-level changes for malaria alone; we computed these from point estimates, without intervals.

Financing exposure and covariates

The primary exposure was external health expenditure as a share of current health expenditure, per 10 percentage points, which measures dependence; the secondary exposure was log(1 + external health expenditure per capita), which measures volume.[33] Both were lagged by 2 years. Models adjusted for log domestic government health expenditure per capita (lagged 2 years), log lag-distributed income per capita, universal health coverage (UHC) effective coverage, and basic sanitation and drinking-water coverage, selected a priori.[15,17,34]

Missing data

GBD outcomes and covariates were complete. Gaps in WDI series (15·1% for lagged external share) were linearly interpolated within countries, with last observations carried forward for up to 2 years; remaining incomplete records were excluded. The analysis samples, restricted to countries outside the GBD high-income super-region (hereafter low- and middle-income countries), comprised 153 countries (2287 country-years) for NTDs and 77 malaria-endemic countries (1187 country-years) for malaria,

2007–2022. Sample derivation and characteristics are in Tables S25–S26; complete-case, balanced-panel, and multiple-imputation analyses assessed sensitivity (Tables S10a–b).

Descriptive and inequality analyses

We summarised burden by super-region, region, and country, ranked countries by count and rate, and identified countries whose rates rose between 2010 and 2023. Socioeconomic inequality was measured in 1990, 2010, and 2023, ranking countries by SDI of the same year and weighting by population. Relative inequality was measured by the concentration index and absolute inequality by the slope index of inequality (SII), the modelled difference in rate between the highest-SDI and lowest-SDI ends of the distribution.[35–37] Country bootstrap resampling (2000 replicates) gave 95% CIs, with paired resampling for changes over time. Theil T was decomposed into components between super-regions, between regions within super-regions, and within regions.[38,39] GBD rates were compared with WHO-reported people requiring NTD interventions and malaria incidence by Spearman correlation.

Spatial autocorrelation

Global Moran's I and local indicators of spatial association were computed on log(1 + rate) with six-nearest-neighbour weights and 9999 permutations; local clusters were reported at a Benjamini–Hochberg false-discovery rate below 5%.[40–42] Analyses were repeated for 1990 and 2010 and with alternative weights (Table S7).

Financing models

The primary analysis estimated within-country associations with two-way (country and year) fixed effects and Driscoll–Kraay standard errors, which allow for the heteroskedasticity, serial correlation, and cross-sectional dependence found in diagnostic tests (Table S3).[43–45] NTD rates were log-transformed and malaria rates log(1 + rate)-transformed. We compared this model with a 2023 cross-section of 145 countries, pooled, random-effects, first-difference and country-trend models, spatial panel models, and a correlated random-effects (Mundlak) model that separates between-country from within-country associations in one sample.[46,47] Estimates are percentage differences per 10-percentage-point higher external share, interpreted as conditional associations rather than causal effects.[48] Malaria mortality was an alternative outcome. Geographically weighted regression and machine-learning prediction are described in the appendix.

Scenarios of financing cuts

Two models projected annual excess DALYs if each country's external health expenditure fell by 5–15% (mild), 19–26% (reference, bracketing the 22% fall in development assistance for health in 2025) or 40–60% (severe) in 153 low- and middle-income countries.[21] The accounting model assumed that the fall in each country's all-age DALY rate between 2010 and 2023, summed over the 19 NTD causes or for malaria, was attributable to external financing in proportion to its share of health expenditure and would reverse in proportion to the cut. The econometric model applied the preferred fixed-effects coefficient to the cut-induced fall in external share; for malaria, the change was computed on the (1 + rate) scale and converted to the implied change in rate. Uncertainty was propagated through 4000 Monte Carlo draws of GBD estimates, cut size, and coefficients.[49] DALYs were valued at one times GDP per capita (0·5 and 3 times in sensitivity analyses), and the rise in domestic government health expenditure needed to replace lost external funds was calculated by super-region.

Statistical software and reproducibility

Analyses used Python (pandas, statsmodels, linearmodels, PySAL, scikit-learn, and LightGBM).[50] Code, data and execution instructions are in the appendix and the study GitHub repository (https://github.com/VincentUwumboriyhieGmayinaam/NTD-Malaria-Burden-Financing). An AI assistant (Claude, Anthropic) helped write and run the analysis code under author supervision (see Acknowledgments). Tests were two-sided at $\alpha=0{\cdot}05$.

Ethics and role of the funding source

This study was based exclusively on secondary analyses of publicly available, aggregated data obtained from the Global Burden of Disease 2023 study, the World Bank's World Development Indicators, the World Health Organization through the United Nations Sustainable Development Goals database, the Institute for Health Metrics and Evaluation's Financing Global Health database, and Natural Earth. All data were available at the country or geographical level and contained no individual-level identifiable information. The study involved no direct recruitment of human participants, collection of primary data, or access to identifiable personal health records. Consequently, formal ethical approval and individual informed consent were not required for this secondary analysis, subject to applicable institutional requirements.

This study was self-funded by the authors. No funder had any role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

Global, regional and national burden

In 2023, NTDs and malaria caused 70·83 million DALYs (95% UI 41·3–113·8 million); country estimates attributed 51·56 million to malaria and 19·24 million to NTDs (Table 1). Sub-Saharan Africa had 15·1% of the population (1208·6 million) but 80·2% of combined DALYs (56·78 million), 95·1% of malaria DALYs (49·03 million) and 40·3% of NTD DALYs (7·75 million). Western sub-Saharan Africa alone accounted for 49·7% of global combined DALYs (35·17 million) and central sub-Saharan Africa for 13·4% (9·49 million), with age-standardised rates of 5556·5 and 5006·5 per 100 000, against 314·2 in southern sub-Saharan Africa; malaria contributed 4828·5 and 4104·3 of those rates (Table S16; Figure S2). Outside Africa, Oceania had the highest rate (1694·8 per 100 000). For NTDs, South Asia held the largest regional share of DALYs (25·0%), followed by western (20·7%) and eastern sub-Saharan Africa (12·3%; Figure 2).

Three countries, India (4·00 million DALYs), Nigeria (2·10 million) and China (1·30 million), accounted for 38·5% of NTD DALYs, and Nigeria (16·63 million) and the Democratic Republic of the Congo (6·46 million) for 44·8% of malaria DALYs (Table S18). Rates were highest in South Sudan for NTDs (2227·1 per 100 000) and Sierra Leone for malaria (8237·2). Among the 19 NTDs, other NTDs (5·14 million DALYs), dengue (3·01 million) and schistosomiasis (2·07 million) contributed most (Figure 3; Table S2). Single countries held most of the burden of several causes: India 53·3% of lymphatic filariasis DALYs, China 75·6% of food-borne trematodiasis, Brazil 58·2% of Chagas disease and the Democratic Republic of the Congo 42·3% of onchocerciasis (Table S20).

Disability accounted for 62·4% of NTD DALYs (12·01 million YLDs), from 54·6% in sub-Saharan Africa to 94·5% in Central Europe, Eastern Europe, and Central Asia, whereas premature death accounted for 95·2% of malaria DALYs (Table S17; Figure S3). GBD country rates correlated closely with WHO-

reported people requiring NTD interventions (Spearman ρ = 0·81) and malaria incidence (ρ = 0·94; Table S21).

Temporal trends, 1990–2023

The global age-standardised rate fell from 1714·0 per 100 000 in 1990 to 1290·8 in 2010 and 1012·5 in 2023, a fall of 21·6% (95% UI 7·8–36·3) since 2010 (Table 1; Figure 4). Since 2010, rates fell most in South Asia (43·8%, 29·2–52·9) and Southeast Asia, East Asia, and Oceania (38·2%, 7·1–56·3), and by 34·1–39·9% in each of the four sub-Saharan African regions; sub-Saharan Africa's rate fell from 5·7 to 3·8 times the global rate between 1990 and 2023. Malaria rates rose only in North Africa and the Middle East (26·3%, 0·1–46·4) and fell to zero in Central Asia, Southern Latin America, and Western Europe (Table S16). Nationally, 78 of 204 countries had a significant decline in the combined rate and four a significant increase: Nicaragua (105·0%, 46·2–178·8), Fiji (92·5%, 62·7–132·2), Monaco (88·8%, 16·9–242·3) and American Samoa (56·2%, 20·4–91·4). Malaria rates were higher in 2023 than in 2010 in 13 countries, including Burundi, Madagascar, Zambia, and Kenya (Table S19).

Spatial clustering and socioeconomic inequality

Burden clustered geographically: Moran's I was 0·71 for NTDs and 0·79 for malaria (both p<0·0001; Table 2), and 0·66–0·82 across alternative weights (Table S7). High–high clusters (48 countries for NTDs and 42 for malaria) lay almost entirely in sub-Saharan Africa (Figure 5) and were stable: all 47 NTD high–high countries of 2010 remained so in 2023, with Somalia added, and Comoros and Somalia joined the malaria cluster (Table S21; Figure S4).

Burden was concentrated in low-SDI countries, and relative concentration changed little over time. The malaria concentration index was −0·79 in 1990 (95% CI −0·88 to −0·63) and −0·81 in 2023 (−0·89 to −0·65); the NTD index moved from −0·51 (−0·64 to −0·36) to −0·43 (−0·54 to −0·28), a change distinguishable from zero only for 2010–2023 (0·05, 0·02 to 0·08). Absolute inequality narrowed: the SII fell from −1966 (−3053 to −1198) to −632 (−877 to −412) DALYs per 100 000 for NTDs and from −3806 (−6754 to −1974) to −2484 (−4271 to −1131) for malaria (Table S15a; Figure S1). Most inequality lay between super-regions (79·3% for malaria and 64·0% for NTDs in 2023), but the share of NTD inequality between countries within the same region rose from 12·6% in 1990 to 21·1% in 2023 (Table S15b).

Financing dependence and burden

Median external expenditure share was 23·0% (IQR 14·6–39·1) in sub-Saharan Africa, 11·5% (1·2–18·4) in Southeast Asia, East Asia, and Oceania and 6·4% (3·5–6·9) in South Asia. Thirty-four countries, 31 in sub-Saharan Africa, were in the highest tertiles of both NTD burden and external share; they held 27·2% of NTD DALYs (5·24 million) and 52·3% of malaria DALYs (26·95 million) (Figure 6). Seventeen high-burden countries, including Brazil, Peru, and Indonesia, had low or intermediate external shares.

Between countries, a 10-percentage-point higher external share was associated with a 13·4% (95% CI 6·4–20·0) lower NTD rate in the 2023 cross-section (Table 3). Within countries, the association was −0·64% (−2·39 to 1·14) in the preferred model and ranged from −1·48% to 0·17% across within-country estimators, all intervals including zero (Figure 7A). In the correlated random-effects model, the between-country association was −4·99% (−11·25 to 1·71) and the within-country association −0·64% (Table 3; Table S23). For malaria, the preferred within-country estimate was −6·32% in (1 + rate) (−10·85 to −1·56), but intervals included zero with country-clustered errors (−17·14 to 5·91) and in other within-country estimators, and the between-country association was positive (18·25%, −13·46 to 61·58). Malaria mortality gave a similar within-country estimate (−6·30%, −10·96 to −1·38; Table S24), as did alternative

missing-data strategies and extension to 2023 (Tables S10b and S22). Higher UHC, sanitation, and drinking-water coverage were each associated with lower NTD burden (Table S4). Local external-share coefficients from multiscale geographically weighted regression were significant in 36 of 145 countries, none in sub-Saharan Africa (Figure 7C; Table S6b).

Projected health losses under development-assistance cuts

Under the reference scenario, the accounting model projected 406 600 excess NTD DALYs a year (95% UI 334 500–501 800) and 1·85 million excess malaria DALYs (1·51–2·21 million), 2·2% and 3·6% of 2023 burden; severe cuts projected 908 700 (708 000–1 140 400) and 4·10 million (3·25–5·07 million; Table 4; Figure 8A–B). The econometric model projected 24 500 excess NTD DALYs (−42 100 to 90 300) and 1·22 million malaria DALYs (0·27–2·29 million). Sub-Saharan Africa accounted for 91% of excess NTD and 96% of excess malaria DALYs in the reference accounting scenario. The largest NTD projections were for the Democratic Republic of the Congo (109 900; 83 600–145 100), Ethiopia (34 700) and Tanzania (32 000), and the largest per-capita projections for South Sudan (246 per 100 000) and the Democratic Republic of the Congo (104; Table S8; Figure 8C). Reference-scenario losses were valued at US$0·55 billion (0·45–0·72) for NTDs and US$2·11 billion (1·65–2·55) for malaria a year. Replacing lost external funds would require domestic government health expenditure to rise by 8·7% in sub-Saharan Africa (US$3·44 billion; 3·9–19·4% across scenarios), 1·8% in South Asia, and 0·5% in North Africa and the Middle East (Table S14; Figure 8D).

Discussion

NTDs and malaria caused 70·8 million DALYs in 2023, 63·1% of them in western and central sub-Saharan Africa. Since 1990 the absolute socioeconomic gap narrowed by about two-thirds for NTDs and one-third for malaria, but burden remained as concentrated in low-SDI countries as before and hot spots did not move. Thirty-four countries combining high NTD burden with high donor dependence held over half of malaria DALYs. The inverse association between external financing and NTD burden seen across countries was not detectable within countries over time, and the malaria association depended on the estimator. These findings locate where need and financing vulnerability coincide; they do not establish the size of losses attributable to withdrawal.

Our 2023 total is close to the 71·6 million DALYs reported for 2021 from GBD 2021;[3] because each GBD round re-estimates the full series, the difference does not measure change. The regional pattern agrees with GBD 2019 and 2021 analyses identifying western sub-Saharan Africa as the highest-burden region,[8,10] and stable high–high clusters extend malaria-only geospatial evidence[27] to NTDs. National concentration matters for planning: three countries held 38·5% of NTD DALYs and two held 44·8% of malaria DALYs, and single countries held most of the burden of lymphatic filariasis, food-borne trematodiasis, and Chagas disease, which aggregate rankings conceal. The post-2015 plateau in malaria incidence in Africa[51] and higher malaria rates in 13 countries in 2023 than in 2010 show that recent progress has been neither uniform nor secure.

Absolute and relative inequality moved differently. Narrowing absolute gaps reflect falling rates in low-SDI countries, but unchanged concentration indices show that these countries still carry the same disproportionate share of burden; GBD 2021 analyses found relative inequality widening to 2021,[8,10] and our estimates to 2023 show at most stabilisation. Targets framed in absolute terms can therefore be met while the relative disadvantage of the poorest countries persists.[36] The rising within-region share of NTD inequality suggests that residual NTD burden increasingly reflects differences between neighbouring countries, consistent with uneven subnational treatment coverage,[12,13] whereas malaria inequality remains

largely between super-regions. Associations of NTD burden with sanitation and water coverage are consistent with evidence on water, housing, and cash transfers,[52–54] but do not identify intervention effects.

The NTD financing association attenuated from 13·4% between countries to 0·64% within countries, and the Mundlak model reproduced this contrast in one sample. Earlier panels associated aid volume with lower malaria and child mortality,[14–16,18] and recent analyses associated USAID and total aid with lower NTD and malaria mortality.[22,23] Our exposure measures dependence rather than volume, and our outcome includes disability, which makes up 62·4% of NTD DALYs and which mortality analyses cannot capture. For malaria, within-country associations with DALYs and deaths were negative, in the direction of programme evaluations,[19,20] whereas the between-country association was positive, consistent with aid allocation to high-burden countries.[27,28] This contrast is why between-country coefficients should not be converted into projected losses. The weak NTD association may reflect an exposure that does not isolate NTD funding, which received 0·6% of health aid in 2003–2007,[55] aggregation of diseases with different response times,[56] and a fixed 2-year lag; these are hypotheses that this study cannot test.

The scenario models bracket structural uncertainty. The accounting model assumes that donor-financed spending produced a matching share of health gains; the econometric model transfers a weak historical association; their NTD projections differ 17-fold. Neither represents the nonlinear rebound that follows interrupted mass drug administration[56] or the resurgence that followed weakened control in 68 of 75 malaria resurgence events,[57] and the programme-specific malaria projections of Symons and colleagues[26] address a different counterfactual. Domestic replacement is uncertain. Reductions in health aid have not historically been offset by domestic spending,[58] aid has had no significant effect on overall domestic public health spending,[59] poorly prepared donor transitions disrupt services,[60] and higher tax revenue is the main route to greater public health spending.[61]

This ecological study cannot establish individual-level effects or locate underserved communities within countries.[62] GBD outcomes are modelled; close agreement with WHO series supports the ordering of countries, not absolute levels. Financing samples were smaller than the descriptive sample, missing WDI values were interpolated, and fixed effects do not remove time-varying confounding. Country-level changes for NTDs excluding malaria and for malaria alone lack uncertainty intervals, and spatial-panel models used balanced subsets of 130 and 71 countries (appendix).

NTD and malaria burden remains concentrated where poverty and donor dependence are greatest, and the 2025 cuts fall on those countries. These data support protecting service continuity there, with priorities adapted to each disease's geography. Measuring the burden attributable to the cuts requires linking programme disbursements to coverage and disease-specific outcomes.

CONTRIBUTORS (Required)

Detail each author's specific contributions (which needs to match your signed Author statement forms); confirm all authors had full access to the data and final responsibility for submission; identify at least two authors who have accessed and verified the underlying data, including at least one from the academic team where applicable.

DECLARATION OF INTERESTS

All other authors declare no competing interests.

DATA SHARING

All data analysed in this study were obtained from publicly accessible secondary data repositories. Disease burden, sociodemographic, and universal health coverage estimates were retrieved from the Global Burden of Disease (GBD) 2023 study through the Institute for Health Metrics and Evaluation (IHME) (https://vizhub.healthdata.org/gbd-results/). Economic and health-system indicators were obtained from the World Bank's World Development Indicators database (https://databank.worldbank.org/source/world-development-indicators). Sustainable Development Goal indicators were obtained from the United Nations SDG Global Database, incorporating data reported by the World Health Organization (https://unstats.un.org/sdgs/dataportal). Development assistance for health estimates were obtained from IHME's Financing Global Health database (https://www.healthdata.org/data-tools-practices/data-sources), and geographical boundary data were sourced from Natural Earth (https://www.naturalearthdata.com/). The analytical dataset was constructed by harmonising and linking these sources using country identifiers and calendar years. All original datasets are accessible through their respective repositories, subject to applicable data-use conditions. The harmonised analytical dataset and analytical code are held in the study GitHub repository (https://github.com/VincentUwumboriyhieGmayinaam/NTD-Malaria-Burden-Financing), which will be made publicly available on publication.

ACKNOWLEDGMENTS

Declaration of generative AI and AI-assisted technologies: during this work the authors used Claude (Anthropic), a generative AI assistant, to retrieve published estimates from the GBD 2023 results tool and literature records from PubMed; to write and run the analysis code for data linkage and the descriptive, inequality, spatial, panel and scenario analyses; to draft the figures, tables and supplementary appendix; and to edit the manuscript, including formatting it to journal style. The authors specified the research questions and study design, reviewed and verified the code, results, citations and text, and take full responsibility for the content of the publication. All analysis code is available in the study GitHub repository.

REFERENCES

1 WHO. Global report on neglected tropical diseases 2025. Geneva: World Health Organization, 2025. https://www.who.int/publications/i/item/9789240114043 (accessed Oct 21, 2026).

2 WHO. World malaria report 2025: addressing the threat of antimalarial drug resistance. Geneva: World Health Organization, 2025. https://www.who.int/publications/i/item/9789240117822 (accessed Oct 21, 2026).

3 Zheng J-X, Liu Y, Guan S-Y, et al. Global, regional, and national burden of neglected tropical diseases and malaria in the general population, 1990–2021: systematic analysis of the Global Burden of Disease Study 2021. J Adv Res 2025; 79: 769–81.

4 GBD 2023 Causes of Death Collaborators. Global burden of 292 causes of death in 204 countries and territories and 660 subnational locations, 1990–2023: a systematic analysis for the Global Burden of Disease Study 2023. Lancet 2025; 406: 1811–72.

5 Lin Y, Fang K, Zheng Y, Wang HL, Wu J. Global burden and trends of neglected tropical diseases from 1990 to 2019. J Travel Med 2022; 29: taac031.

6 Murray CJL, Ortblad KF, Guinovart C, et al. Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet 2014; 384: 1005–70.

7 Emadi M, Delavari S, Bayati M. Global socioeconomic inequality in the burden of communicable and non-communicable diseases and injuries: an analysis on Global Burden of Disease Study 2019. BMC Public Health 2021; 21: 1771.

8 Feng L, Mo G, Liu Z, Liu Q. Global, regional, national burden, trends and health inequality of neglected tropical diseases and malaria from 1990 to 2021. BMC Infect Dis 2025; 25: 1649.

9 He R, Zhu W, Hui S, et al. Inequalities in disease burden and care quality of neglected tropical diseases and malaria, 1990–2021: findings from the 2021 Global Burden of Disease Study. PLoS One 2025; 20: e0329475.

10 Lin Y, Long Q, Sun J, et al. Global trends, health inequalities, and projections in the burden of neglected tropical diseases and malaria from 1990 to 2021. Sci Rep 2025; 15: 20958.

11 Stolk WA, Kulik MC, le Rutte EA, et al. Between-country inequalities in the neglected tropical disease burden in 1990 and 2010, with projections for 2020. PLoS Negl Trop Dis 2016; 10: e0004560.

12 Khaki JJ, Minnery M, Giorgi E. Using ESPEN data for evidence-based control of neglected tropical diseases in sub-Saharan Africa: a comprehensive model-based geostatistical analysis of soil-transmitted helminths. PLoS Negl Trop Dis 2025; 19: e0012782.

13 Lo NC, Heft-Neal S, Coulibaly JT, Leonard L, Bendavid E, Addiss DG. State of deworming coverage and equity in low-income and middle-income countries using household health surveys: a spatiotemporal cross-sectional study. Lancet Glob Health 2019; 7: e1511–20.

14 Akachi Y, Atun R. Effect of investment in malaria control on child mortality in sub-Saharan Africa in 2002–2008. PLoS One 2011; 6: e21309.

15 Bendavid E, Bhattacharya J. The relationship of health aid to population health improvements. JAMA Intern Med 2014; 174: 881–87.

16 Hsiao AJ, Emdin CA. The association between development assistance for health and malaria, HIV and tuberculosis mortality: a cross-national analysis. J Epidemiol Glob Health 2014; 5: 41–48.

17 Mishra P, Newhouse D. Does health aid matter? J Health Econ 2009; 28: 855–72.

18 Yan I, Korenromp E, Bendavid E. Mortality changes after grants from the Global Fund to Fight AIDS, Tuberculosis and Malaria: an econometric analysis from 1995 to 2010. BMC Public Health 2015; 15: 977.

19 Bhatt S, Weiss DJ, Cameron E, et al. The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. Nature 2015; 526: 207–11.

20 Jakubowski A, Stearns SC, Kruk ME, Angeles G, Thirumurthy H. The US President's Malaria Initiative and under-5 child mortality in sub-Saharan Africa: a difference-in-differences analysis. PLoS Med 2017; 14: e1002319.

21 Apeagyei AE, Bisignano C, Elliott H, et al. Tracking development assistance for health, 1990–2030: historical trends, recent cuts, and outlook. Lancet 2025; 406: 337–48.

22 Cavalcanti DM, de Oliveira Ferreira de Sales L, da Silva AF, et al. Evaluating the impact of two decades of USAID interventions and projecting the effects of defunding on mortality up to 2030: a retrospective impact evaluation and forecasting analysis. Lancet 2025; 406: 283–94.

23 da Silva AF, Anderle RVR, Barreix Sibils G, et al. Impact of two decades of humanitarian and development assistance and the projected mortality consequences of current defunding to 2030: retrospective evaluation and forecasting analysis. Lancet Glob Health 2026; 14: e690–701.

24 Hallett TB, Menzies NA, Resch S, et al. The case for optimal investment in combating HIV, tuberculosis, and malaria: a global modelling study. Lancet 2025; 406: 261–70.

25 Stover J, Sonneveldt E, Tam Y, et al. Effects of reductions in US foreign assistance on HIV, tuberculosis, family planning, and maternal and child health: a modelling study. Lancet Glob Health 2025; 13: e1669–80.

26 Symons TL, Lubinda J, McPhail M, et al. Estimating the potential malaria morbidity and mortality avertable by the US President's Malaria Initiative in 2025: a geospatial modelling analysis. Lancet 2025; 405: 2231–40.

27 Namubiru C. Institutional quality, aid flows, and malaria burden: a geospatial analysis of sub-Saharan Africa. Malar J 2025; 24: 332.

28 Sempungu JK, Choi M, Lee EH, Lee YH. The burden of malaria in sub-Saharan Africa and its association with development assistance for health and governance: an exploration of the GBD 2019 study. Glob Public Health 2023; 18: 2229892.

29 WHO Regional Office for Africa. Sustaining momentum in NTD programmes amidst a reduced funding landscape: a regional assessment. Brazzaville: WHO Regional Office for Africa, 2025. https://espen.afro.who.int/sites/default/files/content/document/Sustaining%20Momentum%20in%20NTD%20Programmes%20Amidst%20a%20Reduced%20Funding%20Landscape_A%20Regional%20Assessment_0.pdf (accessed Oct 21, 2026).

30 Stevens GA, Alkema L, Black RE, et al. Guidelines for accurate and transparent health estimates reporting: the GATHER statement. Lancet 2016; 388: e19–23.

31 von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Lancet 2007; 370: 1453–57.

32 GBD 2023 Disease and Injury and Risk Factor Collaborators. Burden of 375 diseases and injuries, risk-attributable burden of 88 risk factors, and healthy life expectancy in 204 countries and territories, including 660 subnational locations, 1990–2023: a systematic analysis for the Global Burden of Disease Study 2023. Lancet 2025; 406: 1873–922.

33 World Bank. World Development Indicators. https://databank.worldbank.org/source/world-development-indicators (accessed Oct 7, 2026).

34 WHO. Ending the neglect to attain the Sustainable Development Goals: a road map for neglected tropical diseases 2021–2030. Geneva: World Health Organization, 2020.

35 Kakwani N, Wagstaff A, van Doorslaer E. Socioeconomic inequalities in health: measurement, computation, and statistical inference. J Econom 1997; 77: 87–103.

36 Mackenbach JP, Kunst AE. Measuring the magnitude of socio-economic inequalities in health: an overview of available measures illustrated with two examples from Europe. Soc Sci Med 1997; 44: 757–71.

37 Wagstaff A, Paci P, van Doorslaer E. On the measurement of inequalities in health. Soc Sci Med 1991; 33: 545–57.

38 Shorrocks AF. The class of additively decomposable inequality measures. Econometrica 1980; 48: 613–25.

39 Theil H. Economics and information theory. Amsterdam: North-Holland, 1967.

40 Anselin L. Local indicators of spatial association—LISA. Geogr Anal 1995; 27: 93–115.

41 Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc Series B Stat Methodol 1995; 57: 289–300.

42 Moran PAP. Notes on continuous stochastic phenomena. Biometrika 1950; 37: 17–23.

43 Driscoll JC, Kraay AC. Consistent covariance matrix estimation with spatially dependent panel data. Rev Econ Stat 1998; 80: 549–60.

44 Hoechle D. Robust standard errors for panel regressions with cross-sectional dependence. Stata J 2007; 7: 281–312.

45 Wooldridge JM. Econometric analysis of cross section and panel data, 2nd edn. Cambridge, MA: MIT Press, 2010.

46 Bell A, Jones K. Explaining fixed effects: random effects modeling of time-series cross-sectional and panel data. Polit Sci Res Methods 2015; 3: 133–53.

47 Mundlak Y. On the pooling of time series and cross section data. Econometrica 1978; 46: 69–85.

48 Imai K, Kim IS. On the use of two-way fixed effects regression models for causal inference with panel data. Polit Anal 2021; 29: 405–15.

49 Briggs AH, Weinstein MC, Fenwick EAL, Karnon J, Sculpher MJ, Paltiel AD. Model parameter estimation and uncertainty analysis: a report of the ISPOR-SMDM Modeling Good Research Practices Task Force Working Group–6. Med Decis Making 2012; 32: 722–32.

50 Rey SJ, Anselin L. PySAL: a Python library of spatial analytical methods. Rev Reg Stud 2007; 37: 5–27.

51 Weiss DJ, Dzianach PA, Saddler A, et al. Mapping the global prevalence, incidence, and mortality of Plasmodium falciparum and Plasmodium vivax malaria, 2000–22: a spatial and temporal modelling study. Lancet 2025; 405: 979–90.

52 Ahmed A, Aune D, Vineis P, Pescarini JM, Millett C, Hone T. The effect of conditional cash transfers on the control of neglected tropical disease: a systematic review. Lancet Glob Health 2022; 10: e640–48.

53 Grimes JET, Croll D, Harrison WE, Utzinger J, Freeman MC, Templeton MR. The relationship between water, sanitation and schistosomiasis: a systematic review and meta-analysis. PLoS Negl Trop Dis 2014; 8: e3296.

54 Tusting LS, Bottomley C, Gibson H, et al. Housing improvements and malaria risk in sub-Saharan Africa: a multi-country analysis of survey data. PLoS Med 2017; 14: e1002234.

55 Liese BH, Schubert L. Official development assistance for health—how neglected are neglected tropical diseases? An analysis of health financing. Int Health 2009; 1: 141–47.

56 Borlase A, Le Rutte EA, Castaño S, et al. Evaluating and mitigating the potential indirect effect of COVID-19 on control programmes for seven neglected tropical diseases: a modelling study. Lancet Glob Health 2022; 10: e1600–11.

57 Cohen JM, Smith DL, Cotter C, et al. Malaria resurgence: a systematic review and assessment of its causes. Malar J 2012; 11: 122.

58 Dieleman JL, Hanlon M. Measuring the displacement and replacement of government health expenditure. Health Econ 2014; 23: 129–40.

59 Patenaude BN. The relationship between development assistance for health and public health financing in 134 countries between 2000 and 2015. Health Policy Plan 2021; 36: 369–83.

60 Huffstetler HE, Bandara S, Bharali I, et al. The impacts of donor transitions on health systems in middle-income countries: a scoping review. Health Policy Plan 2022; 37: 1188–202.

61 Reeves A, Gourtsoyannis Y, Basu S, McCoy D, McKee M, Stuckler D. Financing universal health coverage—effects of alternative tax structures on public health systems: cross-national modelling in 89 low-income and middle-income countries. Lancet 2015; 386: 274–80.

62 Morgenstern H. Ecologic studies in epidemiology: concepts, principles, and methods. Annu Rev Public Health 1995; 16: 61–81.

FIGURE LEGENDS

Figure 1: Analytical workflow. Data acquisition, linkage and quality checks produced a country-year panel for descriptive, inequality, spatial and panel analyses. Panel estimates informed sensitivity analyses and funding-cut scenarios

Figure 2: Geography of NTD and malaria burden, 2023. (A) Age-standardised DALY rate per 100 000 for NTDs excluding malaria and (B) malaria, by country, with GBD super-region boundaries; bars show super-region age-standardised rates. Stepped classes chosen to span the log-distributed rates. Hatched = no data. Equal Earth projection.

Figure 3: Cause-specific burden by GBD super-region, 2023. (A) All-age DALY rate per 100 000 by cause and super-region (log colour scale); labels show the super-region share of global DALYs where ≥50%. (B) Global DALYs by cause (log scale).

Figure 4: Trends in age-standardised DALY rates, 1990–2023. Small multiples by super-region on a shared log scale; shaded band marks 2010–2023, with GBD-published percentage change (95% UI) in the combined NTD and malaria rate.

Figure 5: Spatial clustering and socioeconomic inequality. (A, B) LISA clusters (FDR q<0·05; Equal Earth projection) with global Moran's I; (C) concentration curves ranked by SDI with concentration indices (95% CI); (D) Theil T decomposition into between- and within-super-region inequality.

Figure 6: NTD burden and external health financing. (A) Bivariate tertile map (Equal Earth projection) of age-standardised NTD DALY rates and external expenditure share in the sample outside the GBD high-income super-region. (B) Country-level association, with marker area proportional to NTD DALYs and reference lines at the medians.

Figure 7: Donor financing and burden: estimator comparison, predictive validation and spatial heterogeneity. (A) Percentage change in age-standardised DALY rate per 10-percentage-point external share across between-country, within-country panel and spatial-panel estimators (preferred: two-way FE with Driscoll–Kraay SE). (B) Out-of-sample RMSE of machine-learning and panel models under four validation schemes. (C) Multiscale GWR local coefficients for external share (Equal Earth projection) (significant countries coloured).

Figure 8: Projected consequences of development-assistance cuts. (A, B) Excess annual DALYs in eligible financing-scenario locations under three cut scenarios from the accounting (filled) and econometric (open) models, median and 95% UI; (C) countries with the largest per-capita excess NTD burden (reference scenario); (D) increase in domestic government health spending needed to replace lost external funds, by super-region

SUPPLEMENTARY MATERIAL

Additional material uploaded separately to Editorial Manager, as a single PDF file, with numbered pages, the title of the article and a list of contents.

CHECKLIST (to be deleted upon submission)

☐ Reports conforms to STROBE guidelines.

☐ Refer to the journal's Information for Authors on word count requirements.

☐ Include line numbers in your manuscript.

☐ Tables and figures should be uploaded separately (include permissions for any reproduced figures/tables). As a guide, 3–6 tables/figures (see Information for Authors for the respective journal). Figure titles and short captions should be supplied in the main manuscript after the References section.

☐ Author statement and Conflict of Interest (ICMJE) statement forms should be uploaded as separate documents from the main manuscript to Editorial Manager. Use as many author forms as you need.